Down-Regulation of GABAA Receptor via Promiscuity with the Vasoactive Peptide Urotensin II Receptor. Potential Involvement in Astrocyte Plasticity

Laurence Desrues, Thomas Lefebvre, Céline Lecointre, Marie-Thérèse Schouft, Jérôme Leprince, Vincent Compère, Fabrice Morin, François Proust, Pierrick Gandolfo, Marie-Christine Tonon, et al.

► To cite this version:

Laurence Desrues, Thomas Lefebvre, Céline Lecointre, Marie-Thérèse Schouft, Jérôme Leprince, et al.. Down-Regulation of GABAA Receptor via Promiscuity with the Vasoactive Peptide Urotensin II Receptor. Potential Involvement in Astrocyte Plasticity. PLoS ONE, 2012, 7 (5), pp.e36319. 10.1371/journal.pone.0036319 . hal-01962706

HAL Id: hal-01962706

<https://hal-normandie-univ.archives-ouvertes.fr/hal-01962706>

Submitted on 26 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Down-Regulation of GABA_A Receptor *via* Promiscuity with the Vasoactive Peptide Urotensin II Receptor. Potential Involvement in Astrocyte Plasticity

Laurence Desrues^{1,2,3}, Thomas Lefebvre^{1,2,3}, Céline Lecointre^{1,2}, Marie-Thérèse Schouft^{1,2}, Jérôme Leprince^{1,2}, Vincent Compère^{1,2,3}, Fabrice Morin^{1,2}, François Proust^{1,2,4}, Pierrick Gandolfo^{1,2}, Marie-Christine Tonon^{1,2}, Hélène Castel^{1,2*}

1 Inserm U982, Laboratory of Neuronal and Neuroendocrine Communication and Differentiation, Astrocyte and Vascular Niche, University of Rouen, Mont-Saint-Aignan, France, **2** Institute of Research and Biomedical Innovation (IRIB), Normandy University PRES, University of Rouen, Mont-Saint-Aignan, France, **3** Department of Anesthesiology and Critical Care, Rouen University Hospital, Rouen, France, **4** Department of Neurosurgery, Rouen University Hospital, Rouen, France

Abstract

GABA_A receptor (GABA_AR) expression level is inversely correlated with the proliferation rate of astrocytes after stroke or during malignancy of astrocytoma, leading to the hypothesis that GABA_AR expression/activation may work as a cell proliferation repressor. A number of vasoactive peptides exhibit the potential to modulate astrocyte proliferation, and the question whether these mechanisms may imply alteration in GABA_AR-mediated functions and/or plasma membrane densities is open. The peptide urotensin II (UII) activates a G protein-coupled receptor named UT, and mediates potent vasoconstriction or vasodilation in mammalian vasculature. We have previously demonstrated that UII activates a PLC/PIPs/Ca²⁺ transduction pathway, *via* both G_q and G_{i/o} proteins and stimulates astrocyte proliferation in culture. It was also shown that UT/G_q/IP₃ coupling is regulated by the GABA_AR in rat cultured astrocytes. Here we report that UT and GABA_AR are co-expressed in cerebellar glial cells from rat brain slices, in human native astrocytes and in glioma cell line, and that UII inhibited the GABAergic activity in rat cultured astrocytes. In CHO cell line co-expressing human UT and combinations of GABA_AR subunits, UII markedly depressed the GABA current ($\beta_3\gamma_2 > \alpha_2\beta_3\gamma_2 > \alpha_2\beta_1\gamma_2$). This effect, characterized by a fast short-term inhibition followed by drastic and irreversible run-down, is not relayed by G proteins. The run-down partially involves Ca²⁺ and phosphorylation processes, requires dynamin, and results from GABA_AR internalization. Thus, activation of the vasoactive G protein-coupled receptor UT triggers functional inhibition and endocytosis of GABA_AR in CHO and human astrocytes, *via* its receptor C-terminus. This UII-induced disappearance of the repressor activity of GABA_AR, may play a key role in the initiation of astrocyte proliferation.

Citation: Desrues L, Lefebvre T, Lecointre C, Schouft M-T, Leprince J, et al. (2012) Down-Regulation of GABA_A Receptor *via* Promiscuity with the Vasoactive Peptide Urotensin II Receptor. Potential Involvement in Astrocyte Plasticity. PLoS ONE 7(5): e36319. doi:10.1371/journal.pone.0036319

Editor: Maria A. Deli, Biological Research Centre of the Hungarian Academy of Sciences, Hungary

Received: July 21, 2011; **Accepted:** April 2, 2012; **Published:** May 1, 2012

Copyright: © 2012 Desrues et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by Inserm, University of Rouen and the Conseil régional de Haute-Normandie. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: helene.castel@univ-rouen.fr

† These authors contributed equally to this work.

Introduction

Integrated brain function and dysfunction arise from the complex interactions between a network of multiple cell types including neurons, c and the microvascular endothelial cells comprising the cerebral vasculature [1,2,3]. This micro-environment is a dynamic structure referred as neurovascular unit where polarized astrocytes have a pivotal role [4], rapidly transducing synaptic information [2,3,4,5]. In pathological conditions including stroke, the astroglial reactivity is characterized by proliferation, hypertrophy, process extension, increased synthesis of intermediate filaments, as well as expression of bioactive molecules and their receptors [6,7,8].

GABA_A receptors (GABA_AR) are believed to be pentameric heterooligomers mainly constructed from homologous subunit types α_{1-6} , β_{1-3} , γ_{1-3} , δ and ϵ [9,10,11]. The GABA_AR is expressed in neurons but also in glial cells in culture [12], brain

slices [13], acutely isolated hippocampal slices [13], membrane fractions of the mature rodent brain [14] and also *in vivo* in healthy brain [15]. In pathological conditions, a significant decrease of benzodiazepine sites associated to the GABA_AR has been demonstrated in patients with ischemic cerebrovascular [15,16,17], Parkinson [18] and Alzheimer [19,20] diseases. It was also observed a reduced chloride conductance [21], a decrease in receptor mediated inhibitory post-synaptic potentials [22] and a marked down-regulation of the GABA_AR expression at the cell surface along with a fast time course [15,23,24]. In reactive and malignant astrocytes, mRNA levels of GABA_AR have been shown to remain constant before diminution of functional GABA_AR [15,25]. Thus, the disappearance of GABA_AR expression is correlated with higher glial proliferation rate after stroke or during malignancy of astrocytoma [15,25,26], leading to the hypothesis that GABA_AR expression/activation works as a repressor of cell proliferation. Investigations on alterations in GABA_AR-mediated

functions, receptor densities or modulation in astrocytes remain unchallenged. It has been demonstrated that simultaneous activation of different postsynaptic receptors induces cross-modulation of their activation properties and receptor membrane insertion/deletion. Thus, as many neurotransmitters and vasoactive peptides are released by endothelium and astrocytes, and their receptors are expressed by astrocytes, there is a potential for complex signaling within the neurovascular unit, involving receptor cross-talks.

Urotensin II (U1I) and its paralog urotensin II-related peptide URP, are highly efficient vasoactive peptides, which share a fully conserved C-terminal cyclic CFWKYC core corresponding to the molecular pharmacophore [26,27,28]. The biological actions of U1I and URP are mediated through activation of a G protein-coupled receptor named UT. It is now clearly established that activation of native U1I receptors or UT-transfected cell lines is associated with an increase in polyphosphoinositide (PIPs) turnover promoting a cytosolic calcium concentration ($[Ca^{2+}]_c$) rise [29,30,31]. U1I and UT are expressed in the mammalian cardiovascular system namely in the myocardium, vascular smooth muscle cells and endothelial cells [32,33,34,35], affecting cell proliferation [35,36] or neoangiogenesis [37], stimulating collagen synthesis and cardiac hypertrophy [34].

In the brain, U1I mRNA is particularly abundant in motoneurons of the medulla oblongata and spinal cord [38,39] while UT mRNA is widely expressed in various regions of the central nervous system including the olfactory system, hippocampus, amygdala, hypothalamus, or cerebellum [27]. However, immunohistochemical studies revealed that UT is expressed in astroglial processes *in vivo* [40] and in cultured rat cortical astrocytes [41]. In this cell type, we have previously demonstrated that U1I activates a PLC/PIPs/ Ca^{2+} transduction pathway, *via* both G_q and $G_{i/o}$ proteins and stimulates cell proliferation [41,42]. Moreover, a functional interaction between GABA_AR and UT suggested a cross-talk between these two receptors, involved in astrocyte activity [43]. In this study we demonstrate that activation of UT receptor induces a long-term inhibition of GABA_AR-mediated chloride currents, a process potentially relevant for astrocyte proliferation.

Results

GABA_AR and UT functional coupling in rat cerebellar astrocytes

We and others have previously demonstrated that astrocytes express *in vitro* and *in vivo* UT mRNA and/or protein [38,40,41]. As shown in Figure 1Aa and 1b, UT is extensively expressed in astroglial processes, but also in the sparse mature granule cells present in co-cultures. In order to investigate a potential cross-talk between GABA_AR and UT in both astrocytes and neurons, patch-clamp recording of the GABA_AR agonist isoguvacine (Iso, 10^{-4} M) responses was carried out on membrane potentials and currents of astrocytes in mono-culture, and astrocytes or neurons in co-culture (Figure 1B and 1C). We established that flat and proliferating astrocytes in mono-culture ($n = 31$) (Figure 1B) were rarely responding (only 12% of cells are responding to Iso). In contrast, astrocytes cultured with cerebellar granule neurons present a slowly proliferating stellate astrocytic morphology ($n = 78$) (Figure 1B) and are more responding to the GABA_AR agonist (48% of tested cells). In co-cultured astrocytes, local perfusion of rat U1I ($rU1I$, 10^{-7} M, 40 s) provoked a marked inhibition of the amplitude of the depolarization and chloride current (voltage clamp; -60 mV) evoked by the GABA_AR agonist isoguvacine by $24.2 \pm 7.5\%$ and $33 \pm 8\%$ ($n = 5$), respectively

(Figure 1C). In contrast, $rU1I$ did not affect the Iso-evoked depolarization and current in cerebellar granule neurons (Figure 1C). In astrocytes, $rU1I$ induced a dose-dependent inhibition of the current with an EC_{50} value of $43.6 \pm 23.7\%$ pM (Figure 1D). It can be noticed that in some cells, U1I tested at 10^{-8} M, activated a small inward current. These data indicate that in astrocytes, unlike in neurons, $rU1I$ efficiently and markedly down regulated the GABAergic activity when UT and GABA_AR are co-expressed.

In postnatal day 7 (P7) cerebellar slices, we investigated the topographic cellular and subcellular distribution of UT in the different layers. UT protein immunoreactivity was specifically distributed in particular zones of the cerebellar cortex, in the Purkinje cell layer (PCL), on fibers irradiating from the thin molecular cell layer (ML) toward the external granule cell layer (EGL), and on isolated cells residing in the internal granule cell layer (IGL) (Figure 2). In particular, UT receptor staining was present on NeuN-positive mature granule cell bodies (Figure 2A and 2A') and also in Purkinje neuronal cell bodies and ramifications, as revealed by the strong immunofluorescence of UT in calbindin-positive Purkinje cell soma and dendrites (Figure 2B and 2B'). Doublecortin (DCX) is a distinctive marker of granule cells during the period of radial descent along the Bergmann glia into the cerebellar layers [44]. The DCX immunoreactivity appeared as a diffuse labeling in the IGL and densely in the ML, contiguously associated to UT-positive components but not co-localized (Figure 2C and 2C'). Moreover, UT was strongly co-localized with GFAP expressed by astrocytes of the white matter (*not shown*), and on glial cells (Figure 2D) including Bergmann cells [45]. At higher magnification, the double immunofluorescence analysis revealed that Bergmann cell bodies and fibers were surrounded by yellow co-localizing signals of UT varicosities and of GFAP-positive filaments (Figure 2D'). In addition, immunohistochemical analysis showed labeling of the γ_1 subunit in Purkinje cells, and faint staining in fibers of the ML and in granule cell bodies of the IGL, co-localizing with UT (Figure 2E and 2E'). The γ_2 subunit immunoreactivity was also mainly detected in Purkinje cells and fibers of the EGL (Figure 2F and 2F'). Thus, it appears that UT and GABA_AR subunits are colocalized in cerebellar Purkinje and glial cell fibers *in situ*.

UT mediates inhibition of γ -composed GABA_AR complexes

The molecular basis for the observed directional decrease in GABA_AR function has been investigated in CHO co-expressing human UT and $\alpha\beta$ and/or γ GABA_AR subunits. GABA_AR can be made from several different subunit families (α_1 – α_6 , β_1 – β_3 , γ_1 – γ_3 , ρ_{1-3} , δ , ϵ , π , and θ), which come together in various combinations to form the pentameric receptor [46]. Most receptors are thought to contain α , β and the third subunit type varying, being often the γ_2 . Expression of unitary subunits has produced conflicting results, but some subunits expressed alone [47,48] or as binary combinations, for instance $\alpha_1\gamma_2$ or $\beta_2\gamma_2$, appear to be able to produce GABA-gated ions channels [49,50,51]. In order to establish a direct functional link between GABA_AR and UT, the effect of graded concentrations of $rU1I$ was studied on the current evoked by Iso on CHO-UT, expressing different subunit combinations of GABA_AR (Figure 3A). It is observed that $rU1I$ induced inhibition of GABA_AR complexes which systematically contained γ_1 or γ_2 subunit (Figure 3A and 3B), and the normalized data fit yielded various EC_{50} values and efficacies summarized in supplementary information (Table S1). In particular, $rU1I$ was less effective on the $\alpha_2\beta_3\gamma_1$ and $\alpha_2\beta_1\gamma_1$ and totally ineffective on the $\alpha_2\beta_3$ and $\alpha_2\beta_1$ complexes (Figure 3A and 3B, Table S1).

Figure 1. U1I-induced depression of GABA_AR in UT-expressing cerebellar astrocytes. (Aa, Ab) Double immunofluorescence labeling of UT (green) and the specific astrocyte marker GFAP (red, Aa), or the mature neuron marker NeuN (red, Ab) in astrocyte-neuron co-culture from P7 rat cerebellum. Astrocytes, recognized by strong GFAP staining show UT immunoreactivity (arrows), whereas few weaker UT-stained cells express NeuN (arrowheads), and were likely attributed to mature granule cells (arrowheads, Ab). Nuclei (blue) were counterstained with DAPI. Scale bars, 50 μ m. (B) Phase contrast photomicrograph of astrocytes in mono-culture, or astrocytes and neurons in co-culture at 3 days *in vitro*. (C) Membrane depolarizations and currents evoked by the GABA_AR agonist isoguvacine (Iso, 10^{-4} M, 2 s for membrane potential and 5 s for chloride current) in astrocytes and cerebellar granule neurons before, during rU1I (10^{-7} M, 40 s) application and after 2-min washout. Right, normalized amplitudes deduced by the mean Iso-evoked depolarization or current obtained before rU1I application. (D) Concentration-response relationship of Iso-evoked

currents from astrocytes yielding an EC_{50} value of $43.6 \pm 23.7 \times 10^{-12}$ M. Data are mean $\pm$ SEM of 4 to 6 cells. *, $P < 0.05$; ** $P < 0.01$ compared with the corresponding control Iso-evoked current.
doi:10.1371/journal.pone.0036319.g001

As a control, the effect of the GABA_AR allosteric inverse modulator DMCM, α UII and other uterotensinergic modulators, were tested on the $\alpha_2\beta_3\gamma_2$ GABA_AR function in the absence of

UT. Our data demonstrated that, as expected, DMCM induced inhibition of the current in most tested cells, and α UII and its paralog URP failed to affect the amplitude of the current

Figure 2. Co-localization of UT with γ subunits in neuron and glial components in rat cerebellum. (A, A') Double-fluorescence staining for UT (green) and NeuN (red) showing the presence of UT in both mature (arrowhead, merge, A') and unidentified cells (arrows, merge, A') in the IGL. (B) Co-staining of UT and the marker of Purkinje cells, calbindin (red), in Purkinje cell soma and dendrites (arrowhead, B'). (C) Staining for UT and the marker of migrating neuroblasts doublecortin DCX (red) depicting a diffuse labeling in the ML. (C') UT immunopositive fibers contiguous to DCX-expressing migrating granule cells (merge, yellow, arrowhead). (D, D') Staining for UT and GFAP (red) in glial fibers (merge, yellow, arrowhead) of the ML. (E, F) Distribution of UT and the γ_1 (E) and γ_2 (F) GABA_AR subunits (red), in Purkinje cells (merge, arrowhead) and few extents of glia (merge, arrow) in the ML and IGL. Nuclei (blue) were counterstained with DAPI. Scale bars, 50 μ m (A–F); 20 μ m (A'–F'). EGL, external granule cell layer; IGL, internal granule cell layer; ML, molecular layer; PCL, Purkinje cell layer. (A'–F') images of digitally zoomed regions corresponding to the white boxes in A–F.
doi:10.1371/journal.pone.0036319.g002

A

B

Figure 3. Effect of hUII on different GABA_AR subunit combinations. (A) Typical Iso-evoked currents at the holding potential of -60 mV, in the whole-cell configuration, on CHO stably expressing human UT (CHO-UT) and transiently transfected with cDNAs encoding $\alpha_2\beta_3\gamma_2$, $\alpha_2\beta_1\gamma_2$, $\alpha_2\beta_3\gamma_1$, $\alpha_2\beta_1\gamma_1$, $\beta_3\gamma_2$, $\beta_1\gamma_2$, $\alpha_2\beta_3$ or $\alpha_2\beta_1$ subunits of the GABA_AR. Iso (10^{-4} M) was repeatedly applied for 2 s at 2 min intervals and increasing concentrations of hUII (10^{-13} to 10^{-7} M) were bath perfused in the vicinity of cells. (B) Corresponding concentration-response curves for hUII on $\alpha_2\beta_1\gamma_2$ and $\alpha_2\beta_3\gamma_2$, $\alpha_2\beta_1\gamma_1$ and $\alpha_2\beta_3\gamma_1$, $\beta_1\gamma_2$ and $\beta_3\gamma_2$, $\alpha_2\beta_1$ and $\alpha_2\beta_3$ receptor subunits. Data are normalized to the control Iso response immediately prior to lower hUII concentration application. Data are mean $\pm$ SEM of 3 to 23 cells. doi:10.1371/journal.pone.0036319.g003

Figure 4. Pharmacological characterization of the UT-mediated inhibition of the GABA_AR currents. (A) Whole-cell current response to Iso (10^{-4} M, 2 s) recorded in the absence or presence of the benzodiazepine site inverse agonist DMCM (10^{-5} M), hUII and URP (10^{-8} M, each), or UT antagonists [Orn⁵]-URP and palosuran (10^{-6} M, each) in CHO expressing α_2 , β_3 and γ_2 subunits. Below, summary of the various experimental conditions ($n = 3-18$). (B) Comparison of the inhibitory effect of hUII and DMCM on CHO-UT-GABA_AR, as summarized in bar graphs ($n = 25$). Bottom row, plot of the positive correlation ($r^2 = 0.8$) of hUII-induced inhibition as function of the DMCM-evoked current decrease ($n = 28$). (C) Comparison of the inhibitory effect of hUII and URP on CHO-UT-GABA_AR as summarized in bar graphs ($n = 12-54$). (D, E) Effect of [Orn⁵]-URP and palosuran in the absence or presence of hUII versus the effect of hUII alone. Right, summary of the various

experimental conditions ($n=7-54$). Data are mean $\pm$ SEM from 3 to 54 cells. ns, non significant, *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$ compared with the corresponding control Iso-evoked current.
doi:10.1371/journal.pone.0036319.g004

(Figure 4A). The rapid development in recent years of several UT antagonists has led to the synthesis of [Orn⁵]-URP which has been previously characterized in our laboratory [52,53] and palosuran, with one such high affinity toward human UT [54,55]. Thus, the specific rat UT antagonist [Orn⁵]-URP (10^{-6} M), and the primate UT specific antagonist palosuran (10^{-6} M), did not modulate the Iso-evoked current.

To assess the role of γ subunit into $\alpha\beta$ binary complexes in the U1I-induced GABA_AR regulation, we tested the effect of DMCM, as this allosteric modulator exhibits negative effect on GABA current in cells expressing γ_2 subunit [56,57] or null effect when γ is not present [56]. We analyzed the effect of *h*U1I (10^{-8} M) concomitantly to that of DMCM (10^{-6} M) on CHO-UT co-expressing $\alpha_2\beta_3\gamma_2$ GABA_AR. The amplitude of the *h*U1I-induced inhibition was plotted as a function of the DMCM-induced inhibition of chloride currents recorded from twenty five CHO-UT-GABA_AR expressing cells. A linear regression analysis confirmed the significance of a correlation as represented on the scatter plot ($r^2=0.8$, $P<0.001$) (Figure 4B), suggesting that γ_2 regulates the degree of UT-mediating inhibition of the GABAergic activity.

The effects of the different uterine ligands were then tested on CHO-UT-GABA_AR composed of the γ_2 subunit. We found that *h*U1I and URP (10^{-8} M, each) induced a marked current inhibition by $29.24\pm1.90\%$ ($n=54$) and $22.9\pm3.5\%$ ($n=12$), respectively, that persisted during washout for *h*U1I, but slightly recovered during washout for URP (Figure 4C). In order to examine whether UT antagonists might counteract the U1I-induced decrease of the GABAergic activity, [Orn⁵]-URP and palosuran were tested. [Orn⁵]-URP (10^{-6} M) significantly, but weakly inhibited the Iso-evoked current, blocked the effect of the acute application of *h*U1I, but failed to counteract the prolonged effect of the peptide on the same cell (Figure 4D). Surprisingly, palosuran mimicked the *h*U1I-induced inhibition of the Iso-evoked current, but abolished the irreversible action of *h*U1I during washout (Figure 4E). This indicates that [Orn⁵]-URP or palosuran may keep their antagonist profile toward UT, but exhibit “agonistic activity” in regard to the GABA_AR function.

To test whether activation of the UT/[Ca²⁺]_c signaling pathway may be closely linked to the modulation of GABA_AR, changes in [Ca²⁺]_c evoked by *h*U1I, URP, [Orn⁵]-URP and palosuran were measured by continuous Ca²⁺ fluorescence imaging in CHO-UT. When applied to the bath solution, *h*U1I and URP evoked a significant and rapid increase of the amplitude of the baseline [Ca²⁺]_c by 322% and 341%, respectively (Figure 5A and 5B). The effect of *h*U1I was irreversible, only partially recovering after 32-min washout (Figure 5A), as compared with URP whose effect totally recovered after 16-min washout (Figure 5B). It is also observed that [Orn⁵]-URP (10^{-6} M) behaved as a partial agonist (265% of [Ca²⁺]_c increase) but prevented the sustained effect of *h*U1I on [Ca²⁺]_c (Figure 5C). In contrast, palosuran (10^{-6} M) failed to evoke a [Ca²⁺]_c rise and completely blocked the *h*U1I-induced [Ca²⁺]_c increase (Figure 5D).

Effect of UT on GABA_AR pharmacology and gating properties

According to the action of UT on GABA_AR, we asked the question whether the pharmacology and gating properties of GABA_AR were affected by UT. Thus, we found that the specific positive allosteric GABA_AR modulator pentobarbital (10^{-5} M)

directly activated a chloride current, and reversibly potentiated the Iso-evoked current by $196.28\pm12.33\%$ (Figure S1A). In addition, SR95531 (10^{-5} M) and picrotoxin (10^{-4} M) induced attended current inhibition by $76.92\pm10.35\%$, and by $56.76\pm4.33\%$, respectively (Figure S1A). This indicates that pharmacological characteristics of the GABA_AR are not altered by the presence of UT. Next, to determine whether activation of UT might modify the conductance and selectivity of the GABA_AR-channel complex, the current-voltage (I-V) relationship was studied on CHO-UT co-expressing $\alpha_2\beta_3\gamma_2$ GABA_AR subunits. The voltage-dependence of the cell response to Iso (10^{-4} M) was investigated in the absence or presence of *h*U1I and the amplitude of the current was measured at different holding potentials (Figure S1B). Local perfusion of *h*U1I (10^{-8} M) decreased the slope of the I-V curve but did not significantly shift the chloride reversal potentials (EqCl[−], control, 4.84 mV; *h*U1I, 3.36 mV), closed to the theoretical EqCl[−] value calculated from the Nernst equation, based on the external and internal chloride concentrations used during recording (see Materials and Methods section). It is observed that *h*U1I-induced inhibition of the Iso-evoked current recorded at -60 mV ($26.93\pm6.35\%$) and $+60$ mV ($37.06\pm11.25\%$) was very similar and did not significantly depend on the holding potential (Figure S1B).

The subunit composition determines the GABA sensitivity and the pharmacological properties of the GABA_AR [9] as well as the time course of the GABA response referred as desensitization and deactivation of chloride current [58,59,60]. To clearly assess the mechanism of UT-mediating inhibitions of the GABA current, *h*U1I was applied on CHO-UT-GABA_AR on the fast component of current desensitization. As shown in Figure 6, the current evoked by Iso showed a slow decay during continuous agonist ejection as observed by an apparent desensitization of 59% in control, and 77% under *h*U1I perfusion. We then followed the time-dependent recovery from desensitization in the absence or presence of *h*U1I. Recovering of 50% receptors from this long exposure occurred with mean time constants of 16.2 s in control and 82.2 s in the presence of *h*U1I, demonstrating that the progressive recovery is delayed in the presence of the peptide (Figure 6). Thus, when coexpressed with UT, the pharmacological profile of GABA_AR is not altered, but U1I rather affects macroscopic $\alpha_2\beta_3\gamma_{2L}$ receptor current desensitization, and markedly slows the recovering process. Since recovery from desensitization does not involve membrane voltage [60], it is suggested that UT-mediating GABA_AR desensitization is mainly due to conformational changes of the ligand-bound receptor chloride channel, paralleled to a mechanism known to develop from the closed but fully bound conducting state of the receptor [61].

Mechanisms promoting fast short-term and long-term U1I-induced GABA_AR current inhibition

In CHO-UT-GABA_AR, the Iso-activated currents were measured during a 28-min recording period from the initial application of the GABA_AR agonist. A 1-min application of *h*U1I (10^{-8} M) provoked a fast and significant decrease of the current ($23.44\pm2.47\%$, $n=10$) followed by a progressive run-down, reaching $84.61\pm5.92\%$ ($n=4$) inhibition after 24-min washout (Figure 7A). The second large intracellular loop of several GABA_AR subunits possesses numerous consensus phosphorylation sites [62] and effective phosphorylation mechanisms have been shown to be involved in either short- or long-term regulation of

Figure 5. Role of specific UT ligands on cytosolic calcium in CHO-UT. (A, B) hUII (A) or URP (B) (10^{-8} M, each) provoked a robust increase of $[Ca^{2+}]_c$, which remained stable (A) or recovered to the basal line level (B) during washout. (C, D) Effect of the UT antagonists [Orn⁵]-URP (10^{-6} M, C) or palosuran (10^{-6} M, D), before and during hUII application. Right, bar graphs represent the percent increase of the $[Ca^{2+}]_c$ during drug perfusion or during the washout period. Percent values were obtained by normalizing signals evoked during and after treatments to the value measured before

ligand application. Data are mean $\pm$ SEM from 9 to 25 cells. ns, non significant; ** $P < 0.01$; *** $P < 0.001$ compared with the corresponding control Iso-evoked current. In each type of experiment, three different cells have been selected as representative examples. doi:10.1371/journal.pone.0036319.g005

inhibitory synaptic transmission [63,64]. To further explore the mechanisms sustaining the h U1I-induced current long-term depression, we hypothesized that various UT-couplings to G proteins, relay $[Ca^{2+}]_c$ rise resulting in kinase or phosphatase activation, likely responsible for changes in GABA_AR subunits phosphorylation state. G-protein activation requires the exchange of bound GDP (resting state) with GTP (activated state). This is a common and necessary step of all G-protein mediated actions and is independent of G-protein type or the second messenger system involved. Therefore, blockade of this exchange will result in inability of the ligand-bound receptor to exert its action. Thus, G protein specific blockade with GDP β S (10^{-4} M) did not significantly prevent the fast and long-term inhibition of the current induced by h U1I (Figure 7B). Intracellular dialysis with a cocktail of kinase and phosphatase inhibitors (KIC; phosphatase inhibitor cocktail, quercetin 10^{-5} M and staurosporine 10^{-5} M), failed to alter the fast h U1I inhibitory effect but attenuated the run-down phenomenon (Figure 7C). It is also observed that h U1I reduced the peak current amplitude after 5-s perfusion, but evoked a peak $[Ca^{2+}]_c$ increase only after a 10-s delay (Figure 7D). Consistent with this observation, intracellular BAPTA (10^{-4} M) dialysis reduced the long-term current inhibition by only $39.29 \pm 10.16\%$ ($n = 9$) (Figure 7E). Together, these observations tend to show that G proteins do not transduce UT-induced current inhibition and that calcium transient and phosphorylation mechanisms do not play a promoting role, but participate in the

run-down of the GABA_AR current. To test a hypothetical role of U1I in the dynamin-dependent GABA_AR endocytosis, the dynamin inhibitory peptide DIP, which competitively blocks binding of dynamin to amphiphysin [64], has been introduced in the intrapipette solution. As shown in Figure 7F, when cells were dialyzed with DIP (10^{-5} M), h U1I retained its ability to induce a fast and highly reversible inhibition of the Iso-evoked current, but failed to reduce the current amplitude with time recording.

Inhibition of the GABAergic activity involves GABA_AR internalization and requires specific UT receptor domains in CHO and human astrocytes

To assess whether the C-terminus fragment of UT, corresponding to the cytoplasmic C-tail of the receptor contributes to the U1I-induced run-down of the GABA_AR activity, we constructed four truncated mutants in which the last 19 (UT^{HA}₃₇₀), 38 (UT^{HA}₃₅₁), 57 (UT^{HA}₃₃₂) or 70 (UT^{HA}₃₁₉) residues were removed (Figure 8A). They are all present at the plasma membrane (Figure S2) and functionally expressed except the UT^{HA}₃₁₉ truncated form of UT (Figure S3). When UT^{HA}₃₇₀ is coexpressed with GABA_AR, the h U1I-induced fast short-term inhibition was totally abolished, whereas the long-term inhibition was delayed, and significantly altered (Figure 8B). In the presence of shorter truncated forms, h U1I totally failed to alter the GABA_AR activity, suggesting that the most distal C-terminus part of UT is involved in the functional cross-talk with GABA_AR (Figure 8B). Moreover, the peptidomi-

Figure 6. U1I-induced fast current inhibition and GABA_AR desensitization. (A, B) Examples of currents recorded from CHO-UT-GABA_AR during a long desensitizing pulse (25 s) of Iso (10^{-4} M), in the absence (black line) or presence (green line) of h U1I (10^{-8} M, 1 min). (A) Exponential fit to the desensitizing current phases were shown overlaid on the currents. Bar graphs corresponding to the average desensitization constant parameter τ in the absence (τ_{Ctrl}) or presence (τ_{hU1I}) of h U1I ($n = 5$). (B) Prolonged Iso (30 s) application eliciting current desensitization followed by a time course of the recovery from desensitization, in the absence (control) or presence of h U1I. Graph represents the Iso-evoked current expressed as a fraction of the peak control current induced by the long Iso application to the current amplitude elicited by each short pulse, and plotted against interpulse intervals. Data are mean $\pm$ SEM from 3 to 8 cells. *, $P < 0.05$; **, $P < 0.01$; *** $P < 0.001$ compared with the corresponding control Iso-evoked current. doi:10.1371/journal.pone.0036319.g006

Figure 7. Intracellular mechanisms of UT-triggering GABA_AR inhibition. (A) Traces of Iso (10^{-4} M, 2 s)-evoked current amplitude time-course on CHO-UT-GABA_AR, in control (above row) or during a 1-min application of hUII (10^{-8} M, bottom row). Corresponding average time course of the Iso-evoked current, in control or during and after hUII application. (B, C) Current traces before (1), during (2) a 1-min hUII application and after 20-min washout (3), in the absence or presence of GDP β S (B, 10^{-4} M, 15 min of dialysis) or the cocktail of kinase and phosphatase inhibitors (C, KIC,

15 min of dialysis). Note that the KIC composition consists in phosphatase inhibitor cocktail at 5 mg/ml (sodium vanadate, sodium molybdate, sodium tartrate and imidazole), Quercetin (10 μ M) and staurosporine (10 μ M). In the bottom rows are represented the corresponding average time course of the Iso-evoked current, in the absence or presence of GDP β S (B) or KIC (C). (D) Representative $[Ca^{2+}]_i$ (Fura-2 AM) imaging field before, during *h*U1I application and during washout, and time-course of the fluorescence ratio 340/380. Numbers above each curve indicate the corresponding fluorescent image. The bottom row shows simultaneous currents evoked by repetitive Iso ejections, the time scale has been enlarged to show that the current inhibition occurs before *h*U1I-induced $[Ca^{2+}]_i$ rise. (E, F) Current traces before (1), during (2) a 1-min *h*U1I application and after 20 min washout (3), in the absence or presence of the rapid Ca^{2+} chelator BAPTA (10^{-3} M, E) or the dynamin inhibitory peptide DIP (10^{-5} M, F). In the bottom rows are represented the corresponding average time course of the Iso-evoked current, in the absence or presence of BAPTA (E) or DIP (F). Data are mean $\pm$ SEM from 3 to 21 cells. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ compared with the corresponding control Iso-evoked current. doi:10.1371/journal.pone.0036319.g007

metic UT^{c-myc}_{319–389} (Figure 8A) completely inhibited the fast and long-term effects of *h*U1I on the GABA-evoked current (Figure 8C). Collectively, these results indicate that the C-terminus of UT relays the inhibition of the GABA_AR function, and may counteract a potential *h*U1I-induced internalization process.

Our data thus suggest that UT activation likely regulates GABA_AR endocytosis. We first established the subcellular localization of both UT and GABA_AR in cultured CHO transiently transfected with cDNAs encoding recombinant human UT and the $\alpha_2\beta_3$ ^{HA}- γ_2 -tagged ($\alpha_2\beta_3$ ^{HA} γ_2) GABA_AR subunits and then, internalization of GABA_AR was followed by labeling the surface receptors with antibodies directed against the β_3 ^{HA} co-expressed with $\alpha_2\gamma_2$ GABA_AR subunits and/or UT. In the absence of ligand, the immunoreactivity for β_3 exhibited membrane localization (green) as enlightened by the intensity profiles (Figure 9Aa). In contrast, treatment with *h*U1I (10^{-8} M), Iso (10^{-4} M) or the two agonists, drastically promoted GABA_AR endocytosis by $40.29 \pm 4.14\%$, $39.31 \pm 2.84\%$ and $34.71 \pm 3.19\%$, respectively (Figure 9B), as seen by the increase of red punctuates in the cell soma (Figure 9Ab–8Ad). When GABA_AR was expressed alone, *h*U1I failed to induce GABA_AR internalization whereas Iso or Iso combined with *h*U1I remained able to provoke GABA_AR removal from the plasma membrane (Figure 9B).

In order to confirm the involvement of U1I/UT in the internalization process of GABA_AR in CHO, we investigated the plasma membrane expression of the γ_2 ^{HA} GABA_AR subunit, as well as UT^{c-myc} after exposure to *h*U1I by measuring the amount of surface immunolabelled receptors by ELISA. Typical bioluminescence and fluorescence values obtained from CHO expressing either UT^{c-myc}- $\alpha_2\beta_3$ or UT^{c-myc}- $\alpha_2\beta_3\gamma_2$ ^{HA}, and UT^{c-myc} or UT^{c-myc}- $\alpha_2\beta_3\gamma_2$ ^{HA}-UT_{319–389}YFP, respectively, were shown in Figure 10A. In CHO-UT^{c-myc}- $\alpha_2\beta_3\gamma_2$ ^{HA}, a 30 min treatment with *h*U1I (10^{-8} M) led to approximately 40% loss of γ_2 ^{HA} subunit from the cell surface, without modifying cell membrane amounts of UT^{c-myc} (Figure 10B). In CHO-UT^{c-myc}- $\alpha_2\beta_3\gamma_2$ ^{HA} cotransfected with the cDNA encoding the UT_{319–389}YFP peptidomimetic fragment, *h*U1I incubation failed to remove the γ_2 ^{HA} subunit from the plasma membrane (Figure 10B), establishing that the C-terminus part of UT played a major role in the U1I-induced GABA_AR internalization in a recombinant system. The physiological relevance of such mechanism was thus assessed in native human astrocytes and in the human glioma U87 cell line. Flow cytometry analysis showed that normal and tumoral glial cells in culture expressed β_3 subunit (Figure 11A) and UT (Figure 11B), and that one population (around 8%) of astrocytes and U87 exhibited both receptors at the plasma membrane (Figure 11). *h*U1I (10^{-8} M, 30 min) provoked more than 65% GABA_AR disappearance from the surface, whereas UT internalized in native astrocytes (Figure 11A) but was still present at the cell surface in U87 (Figure 11B). We then examined the cell surface stability of GABA_AR following UT activation in the absence or presence of the coexpressed UT_{319–389}YFP on U87 glioma cell line. ELISA-based assay revealed that the UT_{319–389} fragment reduced the β_3 -associated GABA_AR subunit internalization from 20% to approximately 10% (Figure 11C), with an efficacy of

transfection of around 32%, as controlled by flow cytometry (*data not shown*). As already shown in CHO, UT levels at the cell surface remained unchanged after exposition to *h*U1I, in the absence or presence of the UT C-terminus (Figure 11C).

Discussion

The question of the regulation of the GABA_AR plasticity and cross-modulation in extrasynaptic glial cells by neurovascular factors, and its functional consequence were not explored. In rodent brain, mRNA encoding the vasoactive receptor UT are detected in the olfactory system, hippocampus, amygdala, tegmental nuclei, or cerebellum [27,65,66] and U1I binding sites are restricted to few areas including the cerebellar cortex [39,67]. We and others have also shown that UT receptor expression may be specific to glial cells *in vivo* [40], and to astrocytes in culture [41], as well as to brain microvascular entities (*unpublished data*).

In the present study, to gain insight into the regulation of the GABAergic activity by the vasoactive peptide U1I in astrocyte, the cellular and subcellular distribution and feature of the UT receptor, have been investigated in the cerebellum *in situ*, and in astrocyte-neuron co-cultures. We showed intense UT labeling in Purkinje cell bodies and ramifications as well as in glial Bergmann-GFAP positive long processes where it co-localizes with γ_1 and, to a lesser extent, with γ_2 subunits of the GABA_AR, and in astrocytes co-cultured with granule neurons. Together, this corroborates previous studies establishing expression of α_5 , γ_1 and γ_3 subunits in the PCL [68,69], and of γ_1 mRNA in Bergmann glia [70,71]. We confirmed the coexpression of GABA_AR subunits with UT in native human astrocytes and in the U87 glioma cell line. It is interesting to note that a majority of glial cells expresses at least the β_3 GABA_AR subunit and also UT in the cytosolic compartments, but that only around 10% of cells corresponding to a common subpopulation, show these receptors at the plasma membrane. This is in a good agreement with the 12% responding rat cultured astrocytes to isoguvacine in mono-culture. In fact, the existence of two populations of astrocytes, exhibiting depolarized membrane potentials (around -30 mV) in a majority of cells and hyperpolarized membrane potentials (around -80 mV) in a minority (*data not shown*) was observed in our study, as already shown in cultured astrocytes [72], independent on patch-clamp recording conditions. Here, we determined that this hyperpolarized subpopulation represents astrocytes specifically responding to the GABA_AR activation. Since it was suggested that GABA acts as an anti-proliferating neurotransmitter in ventricular and subventricular zones [73] and in cortical progenitor cells [74] and that down regulation of functional GABA_AR is correlated with the proliferation rate of reactive or malignant astrocytes [15,25], we propose that rat and human cell subpopulation expressing GABA_AR likely correspond to quiescent astrocytes in culture.

These colocalized features of UT and GABA_AR in astrocytes prompted us to investigate a potential functional cross regulation between the two receptors, likely involved in astrocyte plasticity. It has been shown that astrocytic GABA response is specific of early

Figure 8. Receptor sequences involved in UT regulation of the GABA_AR activity. (A) Schematic diagrams mixed with sequence alignments of the HA epitope-tagged human UT, C-terminus truncated UT^{HA}₃₇₀, UT^{HA}₃₅₁, UT^{HA}₃₃₂, UT^{HA}₃₁₉ mutants, and peptidomimetics corresponding to the entire C-terminus cytosolic fragment of UT (UT^{c-myc}₃₁₉₋₃₈₉). (B and C) Traces of Iso (10⁻⁴ M, 2 s)-evoked current before (1), during (2) a 1-min hU1 (10⁻⁸ M) application and after 22-min washout (3). (B) Currents recorded from CHO coexpressing GABA_AR and UT^{HA} (Control), UT^{HA}₃₇₀, UT^{HA}₃₅₁, UT^{HA}₃₃₂ or UT^{HA}₃₁₉. Corresponding average time course of the current, in the absence or presence of UT truncated mutants. (C) Current traces recorded from CHO-UT-GABA_AR, in the absence or presence of UT^{c-myc}₃₁₉₋₃₈₉. Corresponding average time course of the Iso-evoked current, in the absence or presence of UT^{c-myc}₃₁₉₋₃₈₉. In B, significance was only annotated above the time course graph during hU1 perfusion and after 18-min washout, for clarity. Data are mean ± SEM from 3 to 13 cells. ns, non significant; *, *P* < 0.05; **, *P* < 0.01 compared with the corresponding control Iso-evoked current.

doi:10.1371/journal.pone.0036319.g008

culture period which is maintained by interaction with neurons [75]. We demonstrate that UII down regulates the Iso-evoked depolarization and chloride current amplitudes recorded from astrocytes co-cultured with granule neurons. This UII-induced GABA_AR current inhibition is shown to be a very high affinity

process, specific of astrocytes, which hardly recovered during washout. In CHO co-expressing human UT and αβ and/or γ GABA_AR subunits, we found that UII was *i*) very potent on β_{1/3}γ_{1/2L} GABA_AR subunit complexes, *ii*) less potent on α₂β_{1/3}γ_{1/2L} complexes and *iii*) inactive on αβ binary complexes. These results

Figure 9. UT activation mediating GABA_AR internalization. (Aa–Ad) CHO-UT transiently transfected with cDNA encoding $\alpha_2\beta_3^{HA}\gamma_2$ GABA_AR subunits. Internalization was controlled through translocation of β_3^{HA} subunit (red) in control (Aa) or after 60 min of *h*UII (10^{-8} M, Ab), Iso (10^{-4} M, Ac) or *h*UII+Iso (Ad) incubation. Fluorescence intensity plots of green and red fluorescences corresponding to the localization of GABA_AR (β_3^{HA}) at the plasma membrane and in the cytosol, respectively, across the regions delimited by the white line scans. A.u., arbitrary unit; scale bars, 25 μ m. (B) Bar graphs of the fraction of fluorescence at the plasma membrane on CHO-UT-GABA_AR or CHO-GABA_AR in the different conditions. Each bar corresponds to mean $\pm$ SEM percent obtained from 3 to 18 cells. ns, non significant; ***, $P < 0.001$ versus control in CHO-UT-GABA_AR; ###, $P < 0.001$ versus control in CHO-GABA_AR. doi:10.1371/journal.pone.0036319.g009

Figure 10. UII-induced GABA_AR loss from the plasma membrane through the C-terminus fragment of UT in CHO. The effect of hUII on the proportion of GABA_AR and UT at the cell surface of CHO was assessed by ELISA. (A) CHO transiently transfected with cDNA encoding UT^{c-myc} and $\alpha_2\beta_3$, or $\alpha_2\beta_3\gamma_2^{\text{HA}}$ GABA_AR subunits (left), or UT^{c-myc}, and $\alpha_2\beta_3\gamma_2^{\text{HA}}$ GABA_AR subunits cotransfected with the cDNA encoding UT₃₁₉₋₃₈₉YFP (right). Background bioluminescence (left) and fluorescence (right) were measured after anti-HA antibody and colorimetric alkaline phosphatase substrate incubation, in the absence or presence of 30 min of hUII (10^{-8} M, left), or directly on a fluorescent plate reader (right). (B) CHO transiently transfected with cDNA encoding UT^{c-myc} and $\alpha_2\beta_3\gamma_2^{\text{HA}}$ GABA_AR subunits (left), or cotransfected with the cDNA encoding UT₃₁₉₋₃₈₉YFP, and immunodetected with anti-HA (left) or anti-c-myc (right) antibodies. Percentage of cell surface γ_2^{HA} GABA_AR subunit (left) or UT^{c-myc} (right) are represented as the proportion of receptor at the plasma membrane (non permeabilized cells) to the total expressed receptor (permeabilized cells). One hundred percent correspond to values in the absence of 30 min treatment with hUII (10^{-8} M, 37°C). Each bar corresponds to mean $\pm$ SEM percent obtained from 5 to 7 independent experiments, in triplicates. ns, non significant; *, $P < 0.05$; ***, $P < 0.001$. doi:10.1371/journal.pone.0036319.g010

thus establish that UII exhibits a very high affinity directional inhibition toward the GABA_AR specifically composed of the γ subunit. This observation can be paralleled to the high affinity binding sites for UII determined on recombinant UT-expressing

cells [76,77,78,79], and also on astrocytes [41]. The time course of the UT-evoked current inhibition can be distinguished by two phases, *i.e.* a short-term decrease detected immediately after and during UII administration, followed by a progressive run-down of

Figure 11. U11-evoked GABA_AR internalization in native human astrocytes and glioma. (A, B) Flow cytometric analysis of the β_3 GABA_AR subunit and UT expression in native human astrocytes (A) and human U87 glioma cell line (B). Cells were stained with the anti-human β_3 subunit or anti-human UT in permeabilized or non permeabilized conditions (membrane receptor). The black lines depict results from control staining with only secondary antibodies. The β_3 GABA_AR subunit or UT cell surface expression was evaluated in the absence or presence of hU11 (10⁻⁸ M, 30 min) by flow cytometry. Data obtained in A and B illustrate two representative experiments showing β_3 (magenta line) and UT (yellow line) mean fluorescence in the cytosol and at the plasma membrane of a minority of non permeabilized human astrocytes (A) or U87 (B) in culture. The exposure to hU11 induced internalization of β_3 in both cell types and of UT only in U87 glioma. (C) U87 glioma cell line expressing UT and GABA_AR composed of β_3 subunit, and transfected with the cDNA encoding UT₃₁₉₋₃₈₉YFP, and immunodetected with anti- β_3 (left) or anti-UT (right) antibodies. Percentage of

cell surface β_3 subunit (left) or UT (right) are represented as the proportion of receptor at the plasma membrane (non permeabilized cells) to the total expressed receptor (permeabilized cells). One hundred percent correspond to values in the absence of 30 min treatment with hU1I (10^{-8} M, 37°C). Each bar corresponds to mean $\pm$ SEM percent obtained from at least 3 independent experiments, in triplicates. ns, non significant; *, $P < 0.05$; ***, $P < 0.001$.

doi:10.1371/journal.pone.0036319.g011

the current, leading to about 80% GABA_AR current disappearance. However, URP which exhibits the same conserved biologically active cyclic sequence than U1I, triggers a reversible inhibition. This is in accordance with the U1I-induced long-lasting in the one hand, and the URP-evoked transient, on the other hand, increase in $[Ca^{2+}]_c$ in CHO-UT. The specific long-lasting phenomenon might be attributable to the slow dissociation rate of U1I, as already described for rat and human U1I on UT transfected cells, skeletal muscle myoblasts and astrocytes [41,54,79]. This could account for the sustained and washout-resistant contractile responses induced by U1I on primate arteries, $[Ca^{2+}]_c$ increase in rat cortical astrocytes [42], and GABA_AR current inhibition in our native and recombinant systems. We propose a mechanism whereby U1I interacts reversibly with the classical binding site, but also with a secondary exosite in a wash resistant manner, resulting in persistent activation of UT and consequently, in a long-term inhibition of the GABA_AR. Such process has already been demonstrated *in vivo* and cell culture, for exogenous agonists of M1 muscarinic and β_2 adrenergic receptors [80,81]. Together, our previous work suggesting that GABA negatively controls UT-mediated signaling transduction in astrocytes [43], corroborated by the effect of benzodiazepines on U1I-induced neurotransmitter release [82], supports at most the existence of a negative cross-talk coupling between UT and the closely associated GABA_AR, leading to a high affinity functional receptor complex in astrocytes. This functional complex may exhibit new pharmacological profile. Accordingly, we demonstrated that $[Orn^3]$ -URP acts as a partial agonist and competitive antagonist on both GABA currents and $[Ca^{2+}]_c$. Surprisingly, palosuran as a specific primate UT antagonist [55], counteracts the U1I-evoked $[Ca^{2+}]_c$ increase but mimics the effect of U1I by inhibiting the chloride current. Thus, palosuran behaving as an antagonist of the UT-mediated $[Ca^{2+}]_c$ transduction signaling can be considered as a partial UT “agonist” toward the GABA_AR effector pathway, then suggesting a different UT pharmacology when co-expressed with GABA_AR.

Here we found that initiation of UT and GABA_AR functional interaction is independent on G protein, calcium and phosphorylation mechanisms, but that U1I-induced current run-down partially requires calcium and kinase/phosphatase activities. In this context, the Ca^{2+} /calmodulin requirement for membrane fusion in endocytic pathways [83], as well as the clathrin-mediated internalization depending on kinase and phosphatase activities [84,85,86], support here a possible role of UT in a delayed calcium/kinase dependent GABA_AR endocytosis. GABA_AR internalization is primarily thought to occur *via* a clathrin- and dynamin-dependent mechanism [87]. Here DIP, known to block endocytosis by disrupting the interaction between dynamin and amphiphysin, did not interfere with the U1I-induced fast short-term but totally abolished the long-term inhibition of the current, supporting a role of UT in the dynamin-dependent GABA_AR internalization. We then show that the GABA_AR agonist triggered removal of β_3 or γ_2 GABA_AR subunit from CHO plasma membrane, or from native human astrocytic and glioma cell surface. Together, the interesting point resides in the ability of the U1I/UT system in the promotion of the marked GABA_AR internalization in the absence of co-activation of GABA_AR. In addition, 30 min incubation with U1I failed to internalize UT in

CHO and U87, but led to UT loss from human astrocyte surface, a discrepancy unexplained but needing further investigations. Thus, a constitutive tight promiscuity between UT and GABA_AR might be responsible for the high affinity effect of U1I on GABA_AR disappearance from the plasma membrane.

Here we produced and expressed truncated UT receptors in order to identify which specific receptor determinants are involved in the GABA_AR modulation. The deletion of up to 57 residues of the UT C-terminus, did not affect the expression and ability of truncated mutants to stimulate the $[Ca^{2+}]_c$, as already observed for rat UT truncated mutants [88]. We demonstrate that the entire UT C-terminus totally abolished the U1I-mediated current inhibition and that the most distal part likely relays the fast and long-term inhibitions of the GABA_AR function. In rat, the last 19 residues of the C-terminus contain motifs that are not crucial for UT internalization [88]. In fact, the serine cluster localized upstream rather displays consensus motifs for PKC and casein kinase I important for rat UT internalization. In humans, the distal UT C-terminus (351–389) exhibits serine residues and a combination of two polyproline motifs (Figure 8A), the last one possibly interacts with SH3 domain proteins [89]. Our present data in CHO co-expressing human UT and $\alpha_2\beta_3$ and γ_2 GABA_AR and in U87 transfected with the cDNA encoding the UT_{319–389} peptidomimetic, indicate that the U1I-induced internalization of GABA_AR formed from γ_2 or β_3 subunit, requires at least in part, the C-terminus fragment of UT. Therefore, the question of whether UT and GABA_AR physically associate directly or whether SH3 proteins relayed GABA_AR down regulation, has to be elucidated. Together, the functional cross-modulation between UT and GABA_AR is mediated through the most distal part of the UT C-terminus, which would directly interact with γ subunits, or recruit intermediate proteins implicated in GABA_AR inhibitory transactivation.

Therefore, our observations suggest a model in which UT and GABA_AR are closely associated to depress the GABAergic activity (Figure 12). When UT and $\alpha\beta\gamma$ GABA_AR subunits are co-expressed, as in native and tumoral glial cells, U1I efficiently activates its receptor, leading to a fast short-term decrease of the chloride current, independently of G proteins, calcium, phosphorylation and endocytosis processes, and involving the last 19 amino acids of the UT C-terminus. During washout, a long-term inhibition referred as run-down, develops *via* a dynamin-dependent internalization requiring the 351–370 sequence of UT, and calcium- and phosphorylation-dependent endocytic mechanisms. This directional cross-talk between UT and the GABA_AR leads to the extinction of the GABA_AR expression at the plasma membrane that would play a key role in the induction of cell proliferation (Figure 12).

Materials and Methods

Animals

Wistar rats (Dépré, Saint-Doulchard, France) were kept in a temperature-controlled room ($21 \pm 1^\circ\text{C}$), under an established photoperiod (lights on 07.00–19.00 h) with free access to food and tap water. The work described in this article was carried out in accordance with the Directive 2010/63/EU of the European parliament and of the council of 22nd September 2010 on the

Figure 12. Schematic model depicting the mechanism of UT-mediated GABA_AR down-regulation. U11 efficiently activates the G protein-coupled receptor UT, leading to a fast short-term decrease of the chloride current not sustained by G proteins, calcium, phosphorylation and endocytosis processes. This rapid effect involves the distal 19 C-terminal amino acids of UT and the presence of γ subunits within of the GABA_AR complex (1). During the washout period, a long-term inhibition develops via a dynamin-, calcium- and phosphorylation-dependent endocytic mechanisms, requiring at least in part the 351–370 sequence of UT and GABA_AR γ subunits (2). It is hypothesized that the directional cross-talk between UT and GABA_AR, and the extinction of the latter at the plasma membrane, may relay transition from quiescent to proliferant astrocytes. doi:10.1371/journal.pone.0036319.g012

protection of animals used for scientific purposes, published in the Official Journal of the European Union L276/33 (20.10.2010) and authorized by the French Ethical Committee. These experiments were conducted under the supervision of authorized investigators (H. Castel; authorization no. 76.98 from the Ministère de l'Alimentation, de l'Agriculture et de la Pêche) and were approved by the local animal ethic committee of Normandy, approval number N/02-09-09/03/09-12.

Primary Cell Culture of astrocytes and astrocyte-neurone co-culture

Primary cultures of astrocytes were prepared as previously described [90]. Briefly, cerebellum from 7-day-old (P7) Wistar rats were collected in DMEM/Ham-F12 (2:1, v/v) culture medium supplemented with 2 mM glutamine, 1% insulin, 5 mM HEPES, 0.4% D(+)-glucose and 1% of the antibiotic-antimycotic solution. The tissues were disaggregated mechanically with a syringe equipped with a 1-mm gauge needle, and filtered through a 100- μ m pore size mesh filter (Falcon, Becton Dickinson, Grenoble, France). Dissociated cells were resuspended in culture medium supplemented with 10% heat-inactivated FBS and seeded in 150-cm² culture flasks (Falcon) at a density of 20×10^6 cells/flask. Cells were incubated at 37°C in a humidified atmosphere (5% CO₂) and the medium was changed twice a week. When cultures were confluent, the flasks were gently shaken on an orbital shaker at 250 g for 2 h. Dislodged cells were discarded and a second step of purification was performed at 250 g for 14–16 h. Remaining adhesive cells were collected by trypsinization, centrifuged (800 g, 10 min) and plated in 150-cm² flasks. Suspended astrocytes were harvested and seeded in 24-well poly-L-lysine-coated plates. The purity of the cultures was previously assessed by counting the percentage of astrocytes immunostained with GFAP antibodies. The enriched cultures contained >99% astrocytes [41].

For astrocyte-neuron co-culture, granule cell cultures were prepared from cerebella of P7 Wistar rats as described previously [91]. Isolated cells were plated on 14-mm culture dishes coated with poly-L-lysine (5 mM) at a density of 1.5×10^6 cells/dish and incubated at 37°C in a humidified atmosphere (5% CO₂) for 1 to 10 days before use. Culture medium consisted of DMEM/Ham's F12 (75%/25%) supplemented with 10% FBS, 2 mM glutamine, 5 μ g/ml insulin, 25 mM KCl and 1% of antibiotic-antimycotic solution. Co-cultures are obtained by seeding granule cells (1.5×10^6 cells/ml) on cerebellar astrocytes plated in 24-well plates after 12 hours. Co-cultures are maintained in the specific medium for neuron and incubated at 37°C in a humidified atmosphere (5% CO₂) for several days.

Human native astrocytes and glioma cell line

The human cell line from glioblastoma U87 was obtained from the American Type Culture Collection (LGC Standards, Molsheim, France). U87 cells were maintained in DMEM containing 10% FBS and 1% sodium pyruvate. NHA-Astrocytes (Lonza, Walkersville, MD, USA) were cultivated in DMEM culture medium supplemented with 2 mM glutamine, 5 mM HEPES, 1% non essential amino acids, 1% sodium pyruvate, B27, 25 ng/ml EGF, 1% of the antibiotic-antimycotic solution and 10% FBS. All cells were incubated at 37°C in a humidified atmosphere containing 5% CO₂. Culture media were replaced every three days.

CHO recombinant cell line and plasmid transfection

CHO-K1 cell lines were obtained from American Type Culture Collection (Manassas, VA, USA). The human UT stable CHO (CHO-UT) was generously provided by Dr Christophe Dubessy (Inserm, Rouen University, France) et generated by CHO electroporation with 20 μ g pIRES-neo2-UT DNA and 500 μ g

sterile sonicated salmon sperm DNA using the EasyJect One electroporation system (Equibio, Angleur, Belgium), followed by repetitive rounds of limiting dilution of cells in G-418 (400 $\mu\text{g}/\text{ml}$) for selection. Normal or stable CHO was grown in Ham-F12 medium supplemented with 10% FBS, 1% antibiotic-antimycotic solution and 2 mM glutamine, at 37°C in a humidified incubator with an atmosphere of 5% CO_2 . The stable CHO-UT medium was supplemented with the antibiotic G-418 (40 $\mu\text{g}/\text{ml}$).

For transfection, cells were trypsinized (0.05%), tritured in HamF-12 containing 10% FBS media, pelleted by centrifugation, resuspended in 100 μl solution V for nucleofection by an Amaxa Nucleofector Device (Köln, Germany; set to program U-016). Experiments were performed on normal CHO or on the stable CHO-UT cells and transiently transfected with combinations of cDNA encoding $\alpha_2\beta_3\gamma_2$, $\alpha_2\beta_3\gamma_1$, $\alpha_2\beta_1\gamma_2$, $\alpha_2\beta_3\gamma_1$, $\alpha_2\beta_3$, $\beta_3\gamma_2$ (4 μg of cDNA total/transfection) GABA_AR subunits. Cells were seeded on 14-mm poly-L-lysine-coated glass bottom insert dishes at 5×10^5 cells in a volume of 0.5 ml/dish. Cells were incubated overnight at 37°C in a humidified incubator (5% CO_2) during 16 h before electrophysiological or immunocytochemical experiments.

Recombinant receptors

For UT epitope-tagged with HA (UT^{HA}), or c-myc (UT^{c-myc}), human UT receptor cDNA inserted into pcDNA3.1 (Ressource Center, MI, USA) was amplified and PCR products were subcloned using the EcoRI and XhoI sites of pCMV-HA or pCMV-c-myc. Mutant UT receptor cDNAs were constructed by oligonucleotide-directed mutagenesis (Expand High Fidelity PCR System; Roche) using the human UT receptor cDNA inserted into pcDNA3.1 as a template. Two sets of forward and reverse oligonucleotides were used (Table S2) to introduce stop codons in frame of Leu370, Phe351, Gly332 and Thr319 (to generate UT₃₇₁, UT₃₅₁, UT₃₃₂ and UT₃₁₉, respectively). PCR products were subcloned using the EcoRI and XhoI sites of pCMV-HA after digestion by the same restriction enzymes. Mutagenesis was confirmed by automated nucleotide sequencing.

GABA_A receptor cDNA clones; α_2 , β_1 , β_3 , γ_1 and γ_{2L} engineered into the expression vector pCDM8 (α_1 , β_1 , γ_2) or pcDNA/Amp (β_3 , γ_1), were generously provided by Dr Wingrove (Merk Sharp and Dohme, Harlow, UK). To obtain β_3^{HA} or γ_2^{HA} epitope-tagged subunits, β_3 or γ_2 cDNA was amplified and the PCR product was subcloned using the SalI and NotI sites of pCMV-HA (Table S2) after digestion by the same restriction enzymes.

In order to generate mini-peptides corresponding to the C-terminus of UT, cDNA encoding the UT^{c-myc}_{319–389} or UT_{319–389}YFP fragment was amplified by PCR. The 5' and 3' oligonucleotides incorporated SalI and NotI or EcoRI and BamHI sites, respectively, to facilitate subcloning into pCMV-c-myc or pEYFP-N1 (BD Biosciences Clontech, Mississauga, ON, Canada), and incorporated initiation and stop codons where appropriate (Table S2).

Immunocytochemistry on co-culture

Astrocyte-neuron co-cultured on glass coverslips were washed three times in PBS, fixed in 4% paraformaldehyde at 4°C for 20 min, and washed three times in PBS. Cells were permeabilized in PBS containing 0.1% Triton X-100 (10 min) and pre-incubated with normal goat and/or normal donkey antiserum (1:50, Santa-Cruz, Tebu bio, Le Perray en Yvelines, France) for 1 h. Then, cells were incubated at 20°C for 1 h with a goat anti-UT (1:200), a mouse anti-NeuN (1:200, Santa-Cruz) or a rabbit anti-GFAP (1:1000, Dako, Trappes, France). Specificity of the UT immuno-

labelling on astrocytes has already been demonstrated [40]. After several rinses in PBS, cells were incubated at 20°C for 2 h with Alexa 488-conjugated donkey anti-goat and anti-Alexa 594-conjugated donkey anti-rabbit IgGs diluted 1:300 (Invitrogen, Boulogne Billancourt, France).

Receptor cell surface internalization

For double-immunofluorescence, non-permeabilized living CHO cells expressing UT and/or $\alpha_2\beta_3^{\text{HA}}\gamma_2$ GABA_A receptors were washed two times in PBS, and incubated with monoclonal mouse anti-c-myc and rabbit polyclonal anti-HA (1:200, Santa-cruz) for 1 h on ice in DMEM. Excess antibody was removed and cells were incubated with the different receptor agonists for 30 min at RT. After several rinses in PBS, cells were fixed in 4% paraformaldehyde at 4°C for 20 min, washed three times in PBS and then incubated at RT for 2 h with appropriate secondary antibodies, i.e. Alexa-488- and 594-conjugated donkey anti-mouse IgGs diluted 1:300 (Invitrogen). After washing, coverslips were mounted in Eukitt (VWR International, Strasbourg, France).

All preparations were examined using a confocal laser-scanning microscope (Leica, Heidelberg, Germany) equipped with a Diaplan optical system, a UV laser (excitation wavelength 405 nm) and argon/krypton ion (excitation wavelengths 488/594 nm) laser.

Immunohistochemistry on cerebellar sections

Following decapitation, the cerebellum of 7-day-old (P7) Wistar rats was extracted and immersed in iced PBS. Cerebella of P7 rats were sectioned transversely into 180 μm -thick slices on a vibrating blade microtome (VT1000S, Leica Instruments). The slices were subsequently washed in PBS (pH 7.4) and postfixed in a 4% PFA solution for 20 min. Thereafter, free-floating sections were rinsed and non-specific binding was blocked by 10% normal donkey serum, 0.1% BSA and 0.3% Triton X-100 in PBS for 1 h. The tissue sections were incubated overnight at 4°C with a goat antiserum directed against rat UT (1:200, Santa Cruz), a mouse anti-calbindin (1:400, Sigma-Aldrich, Saint-Quentin Fallavier, France), a mouse anti-GFAP (1:1000, Dako), a mouse anti-NeuN raised in mouse (1:200, Santa-Cruz), an anti-doublecortin (DCX) raised in goat (1:400, Santa-Cruz), or anti- γ_1 and anti- γ_2 GABA_AR subunits (1:200, generous gift from Dr Sieghart, Brain Research Institute, Vienna, Austria). The sections were rinsed three times with PBS and incubated for 2 h at RT with Alexa 488- or 594-conjugated donkey anti-rabbit, donkey anti-goat or donkey anti-mouse (1:200, Invitrogen). After washing, slices were incubated with 4,6-diamidino-2-phenylindole (DAPI, 1:10000, Sigma-Aldrich) for 5 min to label nuclei. Finally, the sections were rinsed in PBS, and mounted with mowiol. To study the specificity of UT and other marker immunoreactivities, the following controls were performed (1) substitution of each antiserum by PBS, and evaluation of the level of fluorescence given by each type of secondary antibody, (2) systematic mono-immunolabeling of each protein marker. The preparations were examined on a Leica SP2 upright confocal laser scanning microscope (DM RXA-UV) equipped with Acousto-Optical Beam Splitter (AOBS) system. For confocal images, Alexa-488 and Alexa-594 were excited respectively at 488 and 594 nm.

Electrophysiology

The conventional whole-cell configuration of the patch-clamp technique was used to study the GABA-gated currents in astrocytes and granule neurons, UT stably transfected CHO, CHO-UT, and CHO transiently transfected with diverse variants of UT and GABAAR subunits. After 24-h transfection, cell culture coverslips were placed in a small chamber (1.5 ml) on a stage of a

right microscope DMLFSA (Leica, Heidelberg, Germany) and superfused continuously with the following bath solution containing (in mM): NaCl, 150; KCl, 2.5; HEPES, 5; CaCl₂, 2; MgCl₂, 1; glucose, 10 (pH 7.4 adjusted with NaOH). The patch pipettes were fabricated from 1.5 mm (outer diameter) soft glass tubes on a two-step vertical pipette puller (List-Medical, L/M-3P-A, Darmstadt, Germany). Patch electrodes had a final resistance of 4–6 MΩ when filled with an internal pipette solution containing (in mM): KCl, 130; MgCl₂, 2; CaCl₂, 0.5; EGTA, 5; HEPES, 10; ATP, 1; GTP, 0.1 (pH 7.4 adjusted with KOH). ATP and GTP were added to the internal solution used to fill electrode just before recording. All recordings were obtained at RT with cells voltage-clamped at −70 mV. The GABA_A receptor agonist isoguvacine was prepared in the extracellular solution and was applied to cells by pneumatic pressure ejection. To prevent desensitization, isoguvacine was more often ejected during 5 s at 2-min intervals. ATP, GTP, or guanosine 5′-O-(2-thiodiphosphate) (GDPβS) were administered through the patch pipette solution. We investigated the effect of competitive inhibition of GDP-GTP exchange by including GDPβS in the pipette solution (in addition to the normal amount of GTP). For GDPβS and the KIC, immediately following patch rupture, GABA current recordings were performed and the experiment was commenced after an equilibration period of 15 min.

All peptide ligands, inhibitors or allosteric modulators of the GABA_A receptor function were applied via gravity through a plastic tubing positioned in the vicinity of the cell body in order to maintain a sustained perfusion. In these conditions, drugs could reach the cell of interest after 30 s of perfusion. Isoguvacine was applied focally by pneumatic pressure ejection from a micropipette.

All current signals were amplified from an Axopatch 200A Amplifier (Axon Instruments, Union City, CA, USA) and filtered at 2 kHz (3 dB, four-pole, low-pass Bessel filter). Data acquisition and analysis were performed through a digidata 1200 interface using the pClamp 8 suite programs (Axon Instruments, Union City, CA, USA) and/or the Origin 4.1 analysis software (Microcal Software, Northampton, MA, USA).

The decrease of the chloride current (normalized current, *I*) was defined as $(I_{\text{Iso}} - I_{\text{UII}}) / I_{\text{Iso}}$ where I_{Iso} and I_{UII} is the current response in the presence of various concentrations of UII and I_{Iso} is the control GABA_AR current. Concentration response curves were generated and the data were fitted by a non-linear regression analysis using Microcal Origin Software. Dose-response curves were fitted using a nonlinear square-fitting program to the equation: $F(x) = B_{\text{max}} / [1 + (EC_{50}/x)^n]$, where *x* is the drug concentration, EC_{50} is the concentration of drug eliciting a half-maximal response and *n* is the Hill coefficient.

Cell calcium imaging

For cell calcium imaging, Fura-2 AM (5 mM, Molecular Probe; Fisher, Cergy-Pontoise, France) was dissolved in 20% pluronic F-127 (w/v, DMSO) and then added to culture medium at final concentrations of 5 μM and 0.02%, with 2.5 mM probenecid (Sigma-Aldrich) respectively. Cells were incubated in the dye solution for 1 h in an humidified atmosphere (37°C, 5% CO₂) and then rinsed in the standard extracellular solution used for patch-clamp experiments. For simultaneous measurements of intracellular calcium and chloride-evoked currents in CHO, patch-clamp electrodes were filled with an internal solution containing: KCl, 140; MgCl₂, 4; Fura-2-pentoK, 0.25; HEPES, 10; ATP, 1; GTP, 0.1 (pH 7.4 adjusted with KOH). Fluorescence images were acquired with the right microscope DMLFSA (Leica) equipped with a digital CCD camera Coolsnap HQ (Photometrics, Roper scientific, Evry, France). A high-speed scanning polychromatic light source was used for alternate excitations at wavelengths of

340 and 380 nm. The fluorescence intensities at both wavelengths (F340 and F380) were measured every 500 ms. Image acquisition and analysis were obtained with a MetaFluor/Metamorph Imaging System (Roper scientific). The ratio between the two images was proportional to the $[Ca^{2+}]_i$ in the cell under study. Before an experiment, the bath ground level of fluorescence (attributable to autofluorescence and camera noise) was determined and systematically subtracted.

Cell surface expression of receptors by ELISA

Receptor surface expression was assessed by ELISA 24 h post-transfection of CHO with cDNA encoding UT^{HA} or UT^{HA}-truncated mutants (supplementary Fig. S2) or encoding UT^{c-myc} and α₂β₃γ₂^{HA} GABA_AR subunits, or U87 glioma cell line before and after UII treatment, after quantification of HA, c-myc, β₃ subunit or UT immunoreactivity. Cells were plated at 50 000 cells/well in 96-well plates coated with 0.1 mg/ml poly-L-ornithine (Sigma-Aldrich). After transfection of CHO cells with human UT^{c-myc} and α₂β₃γ₂^{HA} and UT_{319–389}YFP cDNA (Supplementary Table S2), cells were serum starved for 2 h before exposure to UII 10^{−8} M during 30 min at 37°C. Cells were fixed with 4% PFA in PBS for 5 min at RT, washed in PBS, and then permeabilized in 0.05% triton X100 (only for permeabilized cells) and non specific binding were blocked with PBS containing 1% FBS for 30 min at RT. The first rat anti-HA monoclonal antibody (0.5 μg/ml, 3F10 clone, Roche, Meylan, France or 1 μg/ml, Santa-Cruz), mouse anti-c-myc monoclonal antibody (1.33 μg/ml, 9E10 clone, Roche), rabbit anti-UT antibody (1 μg/ml, Tebu, Santa-Cruz) or rabbit anti-β₃ antibody (1:200, Abcam, Paris, France), were added for 1 h30 at RT. Incubation with goat anti-rat (Thermo scientific, Fisher, Brebières, France), goat anti-mouse (Santa Cruz) or goat anti-rabbit (Tebu, Santa Cruz) conjugated alkaline phosphatase diluted at 1:1000 in PBS/FBS was carried out for 30 min at RT. The cells were washed four times with PBS, a colorimetric alkaline phosphatase substrate was added (Super-Signal ELISA, Thermo scientific, Fisher) and the resulting color reaction was measured using a Viktor multilabel plate reader (PerkinElmer, Courtaboeuf, France). Background absorbance from samples transfected with non-tagged receptors or from cells without first antibodies were systematically measured. Results are expressed as the percentage of membrane receptor corresponding to the proportion of receptor at the plasma membrane (non permeabilized cells) to the total receptor (permeabilized cells), and normalized to the values obtained in the absence of UII. All experiments were done at least three times in triplicates.

Receptor expression by Flow cytometry

Human astrocytes and glioma U87 grown in 75-cm² flasks until confluence were washed in PBS, detached and spun down at 4°C. The cell pellet was washed, re-suspended in PBS containing 1% BSA and incubated with 10 μg/mL non-immune rabbit IgGs for 30 min. For total receptor detection, cells were permeabilized in PBS containing 1% BSA and 0.1% saponin for 30 min. Then, cells were incubated with antibodies directed against rabbit anti-β₃ subunit (1:100, Abcam) or anti-UT (1:100, Santa Cruz), diluted in PBS containing 1% BSA and 0.1% saponin at RT (22°C) for 30 min. Non permeabilized cells were directly incubated with rabbit IgGs (10 μg/ml, 30 min), and exposed to the first antibodies for 30 min at 4°C. In negative controls, the primary antibodies were omitted. Cells were incubated with an FITC- or PerCP-conjugated goat anti-rabbit secondary antibody (Santa Cruz) diluted 1:200 and measured in the FL1 (530 nm) or FL3 (670 nm) channel. Cells were analyzed on a FACScalibur flow cytometer (BD Biosciences) operated with the Cell Quest™ software.

Samples in which the primary antibodies were omitted were used as control to normalize mean fluorescence levels. Dead cells and debris were excluded from the analysis by gating living astrocytes or glioma cells from size/structure density plots. Data were displayed on a logarithmic scale in arbitrary units corresponding to the mean fluorescence intensity. Each histogram plot was recorded from at least 10,000 gated events.

Calcium mobilization assay

CHO cells stably transiently transfected with the human UT^{HA} or their truncated forms were plated at a density of 4×10^4 cells/well in flat clear bottom black 96-well plates. After 24 h in culture, cells were incubated at 37°C with 40 μ l of 2 μ M Fluo-4 AM dye containing 20% pluronic acid for 40 min in a 5% CO₂ atmosphere. Cells were washed twice with modified HBSS, and the effects of graded concentrations of U1I on [Ca²⁺]_c were measured with a fluorometric imaging plate reader FlexStation II (Molecular Devices, Sunnyvale, CA) during 150 s with an excitation wavelength of 480 nm and an emission wavelength of 525 nm. After 18 s recording in basal conditions, 50 μ l of graded concentration of U1I (four-fold final concentration) was added to the incubation medium with a built-in eight-channel pipettor to assess activity. After subtraction of mean fluorescence background, baseline was normalized to 100%. Fluorescence peak values were determined for each concentration of U1I, and potency (IC₅₀) and efficacy (E_{max}) were calculated with the Prism 4.0 software using a logistic equation. Results were expressed as mean $\pm$ SEM.

Chemicals

Rat U1I (rU1I, pQHGTAPCEFWKYCI), human U1I (hU1I, ETPDCFWKYCV), URP and [Orn⁵]URP were synthesized by the solid phase methodology on a Pioneer PerSeptive Biosystem peptide synthesizer (Applera France, Courtabœuf, France) using the standard manufacturer's procedures as previously described [32]. All peptides were purified on a 2.2 $\times$ 25-cm Vydac C₁₈ column (Alltech, Templemars, France) (>98% pure) and characterized by MALDI-TOF MS on a Voyager DE-PRO mass spectrometer (Applera France). B27, DMEM, glutamine, HEPES, non essential amino acids, sodium pyruvate and the antibiotic-antimycotic solution were purchased from Invitrogen (Fischer, Illkirch, France). EGF was obtained from Abcys (Les Ulis, France). ATP, GTP, guanosine 5'-[β thio]triphosphate (GDP β S), the GABA_A receptor agonist isoguvacine, pentobarbital, poly-D-ornithine, methyl 6,7-dimethoxy-4-ethyl- β -carboline-3-carboxylate (DMCM), a phosphatase inhibitor cocktail 2 (sodium vanadate, sodium molybdate, sodium tartrate and imidazole), quercetin, staurosporine, picrotoxin, SR95531, Ham-F12, insulin, D(+)-glucose, Tri-reagent, probenicid, and bovine serum albumin (BSA) were obtained from Sigma. FBS was from Eurobio or Lonza (France). Pluronic acid and Fura-2 pentapotassium (Fura-2 AM pentoK) salt and Fura-2 acetoxymethyl ester (Fura-2 AM) were from Molecular Probes (Leiden, Netherlands). The dynamin inhibitory peptide (DIP) was obtained from Tocris Bioscience (Ellisville, MI, USA).

Statistics

All data are presented as mean $\pm$ SEM. Statistical comparisons were assessed with One-way ANOVA followed by Mann and

Whitney, Newman-Keuls or Friedman post hoc tests, as relevant, $P < 0.05$ was taken as significance.

Supporting Information

Figure S1 Pharmacological and gating properties of hU1I-induced regulation of GABA_AR. (A) Iso-evoked current in the absence or presence of pentobarbital (10^{-5} M), SR95531 (10^{-5} M) and picrotoxin (10^{-4} M, 2 s) in CHO-GABA_AR. Right, summary of the effects of modulators on the GABAergic activity. (B) Current-Voltage (I-V) relationship of the Iso-evoked current, in the absence or presence of hU1I (10^{-8} M). Data are mean $\pm$ SEM from 5 to 9 cells. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ compared with the control Iso-evoked current. Ns, non significant. (PPT)

Figure S2 Expression of the UT C-terminus truncated mutants. (A) Confocal microscope images of CHO expressing UT^{HA}, UT₃₁₉^{HA}, UT₃₃₂^{HA}, UT₃₅₁^{HA}, UT₃₇₀^{HA} (green). (B) Expression of the different UT^{HA} mutants expressed as receptors in whole cells (permeabilized) or only at the cell plasma membrane (non-permeabilized) using anti-HA antibody. Data are mean $\pm$ SEM from a representative experiment in triplicate. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ compared to control. Mock, empty pCMV-HA vector. (PPT)

Figure S3 Functional expression of the UT C-terminus truncated mutants. (A) Dose-response curves of the mean of maximum amplitude of [Ca²⁺]_c transients induced by hU1I in CHO expressing UT^{HA}, UT₃₁₉^{HA}, UT₃₃₂^{HA}, UT₃₅₁^{HA}, UT₃₇₀^{HA}. The results are expressed as percentages of the corresponding control values in the absence of hU1I. (B) Corresponding table summarizing EC₅₀ values and percentage of efficacy of the effect of hU1I on each UT construction. Data are mean $\pm$ SEM from 3 independent experiments in duplicate. The Pearson coefficient r^2 close to 1 is used for significance. (PPT)

Table S1 EC₅₀ and maximum inhibitory effects of hU1I on different GABA_AR subunit combinations. Data are mean $\pm$ SEM from 3 to 23 independent experiments. ND, not determined; r^2 , Pearson coefficient. (PPT)

Table S2 Primer sequences and restriction enzymes used for the different UT and GABA_AR subunit constructions. (PPT)

Acknowledgments

We thank Dr L. Prézeau for critical reading of the manuscript. We are grateful to Mrs Huguette Lemonnier and Mr Sébastien Arthaud for technical assistance.

Author Contributions

Conceived and designed the experiments: HC PG MCT VC. Performed the experiments: LD TL CL MTS JL FM. Analyzed the data: HC LD TL. Contributed reagents/materials/analysis tools: JL FP. Wrote the paper: HC LD.

References

1. Takano T, Tian GF, Peng W, Lou N, Libionka W, et al. (2006) Astrocyte-mediated control of cerebral blood flow. *Nature neuroscience* 9: 260–267.
2. Lo EH, Rosenberg GA (2009) The neurovascular unit in health and disease: introduction. *Stroke* 40: S2–3.
3. Ohab JJ, Fleming S, Blesch A, Carmichael ST (2006) A neurovascular niche for neurogenesis after stroke. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 26: 13007–13016.

4. Wang DD, Bordey A (2008) The astrocyte odyssey. *Prog Neurobiol* 86: 342–367.
5. Abbott NJ, Ronnback L, Hansson E (2006) Astrocyte-endothelial interactions at the blood-brain barrier. *Nature reviews Neuroscience* 7: 41–53.
6. Eddleston AL (1993) Immunogenetics of autoimmune chronic active hepatitis. *Gastroenterologia Japonica* 28 Suppl 4: 57–62; discussion 76–80.
7. Silver J, Miller JH (2004) Regeneration beyond the glial scar. *Nature reviews Neuroscience* 5: 146–156.
8. Seifert G, Schilling K, Steinhauser C (2006) Astrocyte dysfunction in neurological disorders: a molecular perspective. *Nature reviews Neuroscience* 7: 194–206.
9. Sieghart W (2006) Structure, pharmacology, and function of GABAA receptor subtypes. *Adv Pharmacol* 54: 231–263.
10. Nutt DJ, Stahl SM (2009) Searching for perfect sleep: the continuing evolution of GABAA receptor modulators as hypnotics. *J Psychopharmacol* 24: 1601–1612.
11. D'Hulst C, Atack JR, Kooy RF (2009) The complexity of the GABAA receptor shapes unique pharmacological profiles. *Drug Discov Today* 14: 866–875.
12. Von Blankenfeld G, Trotter J, Kettenmann H (1991) Expression and Developmental Regulation of a GABAA Receptor in Cultured Murine Cells of the Oligodendrocyte Lineage. *Eur J Neurosci* 3: 310–316.
13. Fraser DD, Duffy S, Angelides KJ, Perez-Velazquez JL, Kettenmann H, et al. (1995) GABAA/benzodiazepine receptors in acutely isolated hippocampal astrocytes. *J Neurosci* 15: 2720–2732.
14. Bureau M, Laschet J, Bureau-Heeren M, Hennuy B, Minet A, et al. (1995) Astroglial cells express large amounts of GABAA receptor proteins in mature brain. *J Neurochem* 65: 2006–2015.
15. Tateishi N, Shimoda T, Manako J, Katsumata S, Shinagawa R, et al. (2006) Relevance of astrocytic activation to reductions of astrocytic GABAA receptors. *Brain Res* 1089: 79–91.
16. Moriaki H, Matsumoto M, Hashikawa K, Oku N, Ishida M, et al. (1998) Iodine-123-iodamazenil and iodine-123-iodoamphetamine SPECT in major cerebral artery occlusive disease. *J Nucl Med* 39: 1348–1353.
17. Muller V, Saur D, Klutmann S, Weiller C, Rother J, et al. (2002) Experience with 123I-iodamazenil SPECT in acute cerebral infarction. *Nucl Med Commun* 23: 1191–1196.
18. Kawabata K, Tachibana H, Sugita M, Fukuchi M (1996) [Impairment of benzodiazepine receptor in Parkinson's disease evaluated by 123I-iodamazenil SPECT]. *Kaku Igaku* 33: 391–397.
19. Ohyama M, Senda M, Ishiwata K, Kitamura S, Mishina M, et al. (1999) Preserved benzodiazepine receptors in Alzheimer's disease measured with C-11 flumazenil PET and I-123 iodamazenil SPECT in comparison with CBF. *Ann Nucl Med* 13: 309–315.
20. Rissman RA, Mishizen-Eberz AJ, Carter TL, Wolfe BB, De Blas AL, et al. (2003) Biochemical analysis of GABA(A) receptor subunits alpha 1, alpha 5, beta 1, beta 2 in the hippocampus of patients with Alzheimer's disease neuropathology. *Neuroscience* 120: 695–704.
21. Verheul HB, de Leeuw FE, Scholten G, Tulleken CA, Lopes da Silva FH, et al. (1993) GABAA receptor function in the early period after transient forebrain ischaemia in the rat. *Eur J Neurosci* 5: 955–960.
22. Luhmann HJ, Mittmann T, van Luijckelaar G, Heinemann U (1995) Impairment of intracortical GABAergic inhibition in a rat model of absence epilepsy. *Epilepsy Res* 22: 43–51.
23. Li H, Siegel RE, Schwartz RD (1993) Rapid decline of GABAA receptor subunit mRNA expression in hippocampus following transient cerebral ischemia in the gerbil. *Hippocampus* 3: 527–537.
24. Schwartz-Bloom RD, Sah R (2001) gamma-Aminobutyric acid(A) neurotransmission and cerebral ischemia. *J Neurochem* 77: 353–371.
25. Labrakakis C, Patt S, Hartmann J, Kettenmann H (1998) Functional GABA(A) receptors on human glioma cells. *Eur J Neurosci* 10: 231–238.
26. Labarrere P, Chatenet D, Leprince J, Marionneau C, Loirand G, et al. (2003) Structure-activity relationships of human urotensin II and related analogues on rat aortic ring contraction. *J Enzyme Inhib Med Chem* 18: 77–88.
27. Sugo T, Murakami Y, Shimomura Y, Harada M, Abe M, et al. (2003) Identification of urotensin II-related peptide as the urotensin II-immunoreactive molecule in the rat brain. *Biochem Biophys Res Commun* 310: 860–868.
28. Douglas SA, Dhanak D, Johns DG (2004) From 'gills to pills': urotensin-II as a regulator of mammalian cardiorenal function. *Trends Pharmacol Sci* 25: 76–85.
29. Liu Q, Pong SS, Zeng Z, Zhang Q, Howard AD, et al. (1999) Identification of urotensin II as the endogenous ligand for the orphan G-protein-coupled receptor GPR14. *Biochem Biophys Res Commun* 266: 174–178.
30. Elshourbagy NA, Douglas SA, Shabon U, Harrison S, Duddy G, et al. (2002) Molecular and pharmacological characterization of genes encoding urotensin-II peptides and their cognate G-protein-coupled receptors from the mouse and monkey. *Br J Pharmacol* 136: 9–22.
31. Ziltener P, Mueller C, Haenig B, Scherz MW, Nayler O (2002) Urotensin II mediates ERK1/2 phosphorylation and proliferation in GPR14-transfected cell lines. *J Recept Signal Transduct Res* 22: 155–168.
32. Sauzeau V, Le Mellionec E, Bertoglio J, Scalbert E, Pacaud P, et al. (2001) Human urotensin II-induced contraction and arterial smooth muscle cell proliferation are mediated by RhoA and Rho-kinase. *Circ Res* 88: 1102–1104.
33. Gong H, Wang YX, Zhu YZ, Wang WW, Wang MJ, et al. (2004) Cellular distribution of GPR14 and the positive inotropic role of urotensin II in the myocardium in adult rat. *J Appl Physiol* 97: 2228–2235.
34. Johns DG, Ao Z, Naselsky D, Herold CL, Maniscalco K, et al. (2004) Urotensin-II-mediated cardiomyocyte hypertrophy: effect of receptor antagonism and role of inflammatory mediators. *Naunyn Schmiedeberg Arch Pharmacol* 370: 238–250.
35. Shi L, Ding W, Li D, Wang Z, Jiang H, et al. (2006) Proliferation and anti-apoptotic effects of human urotensin II on human endothelial cells. *Atherosclerosis* 188: 260–264.
36. Watanabe T, Pakala R, Katagiri T, Benedict CR (2001) Synergistic effect of urotensin II with mildly oxidized LDL on DNA synthesis in vascular smooth muscle cells. *Circulation* 104: 16–18.
37. Guidolin D, Albertin G, Oselladore B, Sorato E, Rebuffat P, et al. (2010) The pro-angiogenic activity of urotensin-II on human vascular endothelial cells involves ERK1/2 and PI3K signaling pathways. *Regul Pept* 162: 26–32.
38. Coulouarn Y, Lihmann I, Jegou S, Anouar Y, Tostivint H, et al. (1998) Cloning of the cDNA encoding the urotensin II precursor in frog and human reveals intense expression of the urotensin II gene in motoneurons of the spinal cord. *Proc Natl Acad Sci U S A* 95: 15803–15808.
39. Coulouarn Y, Jegou S, Tostivint H, Vaudry H, Lihmann I (1999) Cloning, sequence analysis and tissue distribution of the mouse and rat urotensin II precursors. *FEBS Lett* 457: 28–32.
40. Lin Y, Tsuchihashi T, Matsumura K, Fukuhara M, Ohya Y, et al. (2003) Central cardiovascular action of urotensin II in spontaneously hypertensive rats. *Hypertens Res* 26: 839–845.
41. Castel H, Diallo M, Chatenet D, Leprince J, Desrues L, et al. (2006) Biochemical and functional characterization of high-affinity urotensin II receptors in rat cortical astrocytes. *J Neurochem* 99: 582–595.
42. Jarry M, Diallo M, Lecointre C, Desrues L, Tokay T, et al. (2010) The vasoactive peptides urotensin II and urotensin II-related peptide regulate astrocyte activity through common and distinct mechanisms: involvement in cell proliferation. *Biochem J* 428: 113–124.
43. Desrues L, Lefebvre T, Diallo M, Gandolfo P, Leprince J, et al. (2008) Effect of GABA A receptor activation on UT-coupled signaling pathways in rat cortical astrocytes. *Peptides* 29: 727–734.
44. Sotelo C (2004) Cellular and genetic regulation of the development of the cerebellar system. *Prog Neurobiol* 72: 295–339.
45. Taft JR, Vertes RP, Perry GW (2005) Distribution of GFAP+ astrocytes in adult and neonatal rat brain. *Int J Neurosci* 115: 1333–1343.
46. Tretter V, Ehya N, Fuchs K, Sieghart W (1997) Stoichiometry and assembly of a recombinant GABAA receptor subtype. *J Neurosci* 17: 2728–2737.
47. Blair LA, Levitan ES, Marshall J, Dionne VE, Barnard EA (1988) Single subunits of the GABAA receptor form ion channels with properties of the native receptor. *Science* 242: 577–579.
48. Pritchett DB, Sontheimer H, Gorman CM, Kettenmann H, Seeburg PH, et al. (1988) Transient expression shows ligand gating and allosteric potentiation of GABAA receptor subunits. *Science* 242: 1306–1308.
49. Sigel E, Baur R, Trube G, Mohler H, Malherbe P (1990) The effect of subunit composition of rat brain GABAA receptors on channel function. *Neuron* 5: 703–711.
50. Krishek BJ, Xie X, Blackstone C, Huganir RL, Moss SJ, et al. (1994) Regulation of GABAA receptor function by protein kinase C phosphorylation. *Neuron* 12: 1081–1095.
51. Angelotti TP, Macdonald RL (1993) Assembly of GABAA receptor subunits: alpha 1 beta 1 and alpha 1 beta 1 gamma 2S subunits produce unique ion channels with dissimilar single-channel properties. *J Neurosci* 13: 1429–1440.
52. Diallo M, Jarry M, Desrues L, Castel H, Chatenet D, et al. (2008) [Orn5]URP acts as a pure antagonist of urotensinergic receptors in rat cortical astrocytes. *Peptides* 29: 813–819.
53. Chatenet D, Dubessy C, Leprince J, Boularan C, Carlier L, et al. (2004) Structure-activity relationships and structural conformation of a novel urotensin II-related peptide. *Peptides* 25: 1819–1830.
54. Clozel M, Binkert C, Birker-Robaczewska M, Boukhadra C, Ding SS, et al. (2004) Pharmacology of the urotensin-II receptor antagonist palosuran (ACT-058362; 1-[2-(4-benzyl-4-hydroxy-piperidin-1-yl)-ethyl]-3-(2-methyl-quinolin-4-yl)-urea sulfate salt): first demonstration of a pathophysiological role of the urotensin system. *J Pharmacol Exp Ther* 311: 204–212.
55. Behm DJ, McAtee JJ, Dodson JW, Neeb MJ, Fries HE, et al. (2008) Palosuran inhibits binding to primate UT receptors in cell membranes but demonstrates differential activity in intact cells and vascular tissues. *Br J Pharmacol* 155: 374–386.
56. Puia G, Vicini S, Seeburg PH, Costa E (1991) Influence of recombinant gamma-aminobutyric acid-A receptor subunit composition on the action of allosteric modulators of gamma-aminobutyric acid-gated Cl⁻ currents. *Mol Pharmacol* 39: 691–696.
57. Wafford KA, Whiting PJ, Kemp JA (1993) Differences in affinity and efficacy of benzodiazepine receptor ligands at recombinant gamma-aminobutyric acidA receptor subtypes. *Mol Pharmacol* 43: 240–244.
58. Bianchi MT, Haas KF, Macdonald RL (2001) Structural determinants of fast desensitization and desensitization-deactivation coupling in GABA_A receptors. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 21: 1127–1136.
59. Boileau AJ, Baur R, Sharkey LM, Sigel E, Czajkowski C (2002) The relative amount of cRNA coding for gamma2 subunits affects stimulation by benzodiazepines in GABA(A) receptors expressed in *Xenopus* oocytes. *Neuropharmacology* 43: 695–700.

60. Yoon KW (1994) Voltage-dependent modulation of GABAA receptor channel desensitization in rat hippocampal neurons. *Journal of neurophysiology* 71: 2151–2160.
61. Dominguez-Perrot C, Feltz P, Poulter MO (1996) Recombinant GABAA receptor desensitization: the role of the gamma 2 subunit and its physiological significance. *The Journal of physiology* 497(Pt 1): 145–159.
62. Moss SJ, Smart TG (1996) Modulation of amino acid-gated ion channels by protein phosphorylation. *Int Rev Neurobiol* 39: 1–52.
63. Brandon N, Jovanovic J, Moss S (2002) Multiple roles of protein kinases in the modulation of gamma-aminobutyric acid(A) receptor function and cell surface expression. *Pharmacol Ther* 94: 113–122.
64. Kittler JT, Moss SJ (2003) Modulation of GABAA receptor activity by phosphorylation and receptor trafficking: implications for the efficacy of synaptic inhibition. *Curr Opin Neurobiol* 13: 341–347.
65. Gartlon J, Parker F, Harrison DC, Douglas SA, Ashmeade TE, et al. (2001) Central effects of urotensin-II following ICV administration in rats. *Psychopharmacology (Berl)* 155: 426–433.
66. Jegou S, Cartier D, Dubessy C, Gonzalez BJ, Chatenet D, et al. (2006) Localization of the urotensin II receptor in the rat central nervous system. *J Comp Neurol* 495: 21–36.
67. Clark SD, Nothacker HP, Wang Z, Saito Y, Leslie FM, et al. (2001) The urotensin II receptor is expressed in the cholinergic mesopontine tegmentum of the rat. *Brain Res* 923: 120–127.
68. Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G (2000) GABA(A) receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* 101: 815–850.
69. Fritschy JM, Panzanelli P (2006) Molecular and synaptic organization of GABAA receptors in the cerebellum: Effects of targeted subunit gene deletions. *Cerebellum* 5: 275–285.
70. Laurie DJ, Wisden W, Seeburg PH (1992) The distribution of thirteen GABAA receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci* 12: 4151–4172.
71. Bovolin P, Santi MR, Puia G, Costa E, Grayson D (1992) Expression patterns of gamma-aminobutyric acid type A receptor subunit mRNAs in primary cultures of granule neurons and astrocytes from neonatal rat cerebella. *Proc Natl Acad Sci USA* 89: 9344–9348.
72. McKhann GM, D'Ambrosio R, Janigro D (1997) Heterogeneity of astrocyte resting membrane potentials and intercellular coupling revealed by whole-cell and gramicidin-perforated patch recordings from cultured neocortical and hippocampal slice astrocytes. *J Neurosci* 17: 6850–6863.
73. Haydar TF, Wang F, Schwartz ML, Rakic P (2000) Differential modulation of proliferation in the neocortical ventricular and subventricular zones. *J Neurosci* 20: 5764–5774.
74. LoTurco JJ, Owens DF, Heath MJ, Davis MB, Kriegstein AR (1995) GABA and glutamate depolarize cortical progenitor cells and inhibit DNA synthesis. *Neuron* 15: 1287–1298.
75. Nilsson M, Eriksson PS, Ronnback L, Hansson E (1993) GABA induces Ca2+ transients in astrocytes. *Neuroscience* 54: 605–614.
76. Ames RS, Sarau HM, Chambers JK, Willette RN, Aiyar NV, et al. (1999) Human urotensin-II is a potent vasoconstrictor and agonist for the orphan receptor GPR14. *Nature* 401: 282–286.
77. Mori M, Sugo T, Abe M, Shimomura Y, Kurihara M, et al. (1999) Urotensin II is the endogenous ligand of a G-protein-coupled orphan receptor, SENR (GPR14). *Biochem Biophys Res Commun* 265: 123–129.
78. Nothacker HP, Wang Z, McNeill AM, Saito Y, Merten S, et al. (1999) Identification of the natural ligand of an orphan G-protein-coupled receptor involved in the regulation of vasoconstriction. *Nat Cell Biol* 1: 383–385.
79. Qi J, Du J, Tang X, Li J, Wei B, et al. (2004) The upregulation of endothelial nitric oxide synthase and urotensin-II is associated with pulmonary hypertension and vascular diseases in rats produced by aorticaval shunting. *Heart Vessels* 19: 81–88.
80. Green SA, Spasoff AP, Coleman RA, Johnson M, Liggett SB (1996) Sustained activation of a G protein-coupled receptor via “anchored” agonist binding. Molecular localization of the salmeterol exosite within the 2-adrenergic receptor. *J Biol Chem* 271: 24029–24035.
81. Jakubik J, Tucek S, El-Fakahany EE (2002) Allosteric modulation by persistent binding of xanomeline of the interaction of competitive ligands with the M1 muscarinic acetylcholine receptor. *J Pharmacol Exp Ther* 301: 1033–1041.
82. Ono T, Kawaguchi Y, Kudo M, Kushikata T, Hashiba E, et al. (2008) Urotensin II evokes neurotransmitter release from rat cerebrocortical slices. *Neurosci Lett* 440: 275–279.
83. Pryor PR, Mullock BM, Bright NA, Gray SR, Luzio JP (2000) The role of intraorganellar Ca(2+) in late endosome-lysosome heterotypic fusion and in the reformation of lysosomes from hybrid organelles. *J Cell Biol* 149: 1053–1062.
84. Lin FT, Krueger KM, Kendall HE, Daaka Y, Fredericks ZL, et al. (1997) Clathrin-mediated endocytosis of the beta-adrenergic receptor is regulated by phosphorylation/dephosphorylation of beta-arrestin1. *J Biol Chem* 272: 31051–31057.
85. Herring D, Huang R, Singh M, Robinson LC, Dillon GH, et al. (2003) Constitutive GABAA receptor endocytosis is dynamin-mediated and dependent on a dileucine AP2 adaptin-binding motif within the beta 2 subunit of the receptor. *J Biol Chem* 278: 24046–24052.
86. Naga Prasad SV, Jayatilake A, Madamanchi A, Rockman HA (2005) Protein kinase activity of phosphoinositide 3-kinase regulates beta-adrenergic receptor endocytosis. *Nat Cell Biol* 7: 785–796.
87. Barnes EM, Jr. (2001) Assembly and intracellular trafficking of GABAA receptors. *Int Rev Neurobiol* 48: 1–29.
88. Proulx CD, Simaan M, Escher E, Laporte SA, Guillemette G, et al. (2005) Involvement of a cytoplasmic-tail serine cluster in urotensin II receptor internalization. *Biochem J* 385: 115–123.
89. Cao W, Luttrell LM, Medvedev AV, Pierce KL, Daniel KW, et al. (2000) Direct binding of activated c-Src to the beta 3-adrenergic receptor is required for MAP kinase activation. *J Biol Chem* 275: 38131–38134.
90. Gandolfo P, Patte C, Leprince J, Thoumas JL, Vaudry H, et al. (1997) The stimulatory effect of the octadecaneuropeptide (ODN) on cytosolic Ca2+ in rat astrocytes is not mediated through classical benzodiazepine receptors. *Eur J Pharmacol* 322: 275–281.
91. Lefebvre T, Gonzalez BJ, Vaudry D, Desrues L, Falluel-Morel A, et al. (2009) Paradoxical effect of ethanol on potassium channel currents and cell survival in cerebellar granule neurons. *J Neurochem* 110: 976–989.