1 Invasion by *Fallopia japonica* alters soil food webs through secondary metabolites

2

3 Abgrall, C.^{a,b}; Forey, E.^a; Mignot, L.^a; Chauvat, M.^a

4

5

6 ^a Ecodiv URA/EA 1293, IRSTEA, FR CNRS 3730 SCALE, UFR Sciences et Techniques,

7 Normandie université, Université de Rouen, 76821 Mont Saint Aignan Cedex, France

8 ^b Corresponding author ; corentin.abgrall1@univ-rouen.fr

9

10 Abstract

11

12 Biological invasions are a major threat to biodiversity with varying degrees of impact. There

13 is increasing evidence that allelopathy often plays an important role in explaining both

14 invasion success and impact on native taxa (e.g. novel weapons hypothesis). The effects of

15 these secondary metabolites on plant communities and microorganisms are well known.

16 However, their direct and indirect effects on soil fauna are unresolved, despite the

17 importance of the latter in ecosystem processes and, potentially, invasion mitigation.

18 Japanese knotweed (*Fallopia japonica*), an east-Asian species, which has proved to be

19 invasive in Europe, containing allelopathic secondary compounds inhibiting native plants

20 and microbial communities. The focal point of this study was the allelopathic effects of

21 knotweed on soil mesofauna (Nematoda, Collembola and Acari). During a one-month

22 laboratory experiment we added knotweed rhizome extract (KRE) at different

23 concentrations to soils collected in an invasion-prone area. The experiment consisted of

including or excluding secondary metabolites through the use of activated carbon filtration of KRE. This enabled us to separate effects caused by nutrient addition (i.e. trophic effects) and combined (trophic and allelopathic) effects. Relative effects of nutrient and secondary metabolites addition on abiotic and biotic soil variables were then quantified. We highlighted frequently contrasting trophic and allelopathic effects influenced in some cases by KRE concentration. Microbial assemblages, through fungal / microbial biomass ratio, did not show any congruent response to KRE secondary compounds but was more negatively impacted by nutrient addition. The use of a trophic-based path analysis led us to show that changes within the soil biota had repercussions on secondary consumers (e.g. bacterivorous nematodes and Collembola). Abundance within taxa at higher trophic levels such as predatory Acari (but not predatory nematodes) was also affected although to a lesser extent, likely in part due to the limited considered timeframe. Overall, we showed that, in controlled conditions, invasive allelopathic plants such as knotweeds can have effects on soil fauna at different trophic levels through addition of both nutrients and secondary metabolites to the soil. Considering the limited knowledge of allelopathic effects on the soil fauna and soil functions, this study provides new information on above- and belowground interactions.

Keywords

plant-soil interactions ; novel weapons hypothesis ; allelopathy ; trophic networks ; alien species

1. Introduction

48 Past and current introduction of invasive plant species and their spread in new ecosystems
49 is a major concern for conservation at a global level (Litt et al., 2014; Pyšek et al., 2012) due
50 to their severe impact on biodiversity (Murrell et al., 2011; Vilà et al., 2011) and ecosystem
51 processes (Bassett et al., 2011; Kohyt and Skubała, 2013). Only a small number of exotic
52 species become invasive in their introduced range (Reinhart and Callaway, 2006) through
53 distinctive characteristics (or traits) providing superior competitive ability when compared
54 to native species (Van Kleunen et al., 2010). These traits can be morphological in nature by
55 directly improving plant fitness (Van Kleunen et al., 2010) or physiological with the synthesis
56 of biochemical, secondary metabolites that influence the germination, growth, survival
57 and/or reproduction of other organisms (Inderjit et al., 2011b).

58 The novel weapon hypothesis (NVH) suggests that the success of many exotic invasive plant
59 species is due to the possession of allelopathic compounds unencountered by native
60 species, particularly native plant species (Callaway and Ridenour, 2004). Furthermore, it has
61 been shown that many invasive species have different allelopathic potential effects between
62 their native and introduced ranges (Inderjit et al., 2011a; Thorpe et al., 2009). These
63 biochemical compounds, exudated from plant roots (Callaway et al., 2008) or released from
64 degrading litter (Inderjit et al., 2011a) have powerful effects on ecosystem functioning by
65 impacting both organisms and ecological processes (Hättenschwiler et al., 2011;
66 Hättenschwiler and Vitousek, 2000; Reigosa et al., 2006; Wardle et al., 1998).

67 Under the soil, plants interact with a wide range of organisms including bacteria, fungi,
68 nematodes and various kinds of arthropods (Abgrall et al., 2017; Parepa et al., 2013). These
69 aboveground-belowground relationships can be antagonistic (e.g. herbivores, pathogens) or
70 mutualistic (e.g. mycorrhizous fungi, nitrogen-fixing bacteria) (Van der Putten et al., 2007).
71 Allelopathic biochemical that have a negative effect on plants can do so indirectly by
72 promoting or inhibiting particular soil biota (Callaway et al., 2008; Stinson et al., 2006).

73 Furthermore, the soil biota is known as having a structuring influence on plant community
74 composition, dynamics and phenology (Forey et al., 2015; Wardle, 2002), allelopathy
75 feedback from the soil biota could further increase invasion (Parepa et al., 2013).

76 Japanese knotweed (*Fallopia japonica* (Houtt.) Ronse Decr. 1988, Polygonaceae) was
77 introduced in Europe in the 19th century for its ornamental properties. It is now one of the
78 most destructive invasive species in Europe and North America (Lowe et al., 2000). *F.*
79 *japonica* spreads mostly by clonal rhizomatous growth with a single stem or rhizome node
80 being able to regenerate a full plant explaining the high-dispersion capacity of knotweed (De
81 Waal, 2001). Multiple species of the genus *Fallopia* such as *F. japonica*, *F. sachalinensis* as
82 well as their hybrid *F. ×bohemica* are known to contain and produce several secondary
83 metabolites (Murrell et al., 2011). Some of those compounds exhibit allelopathic properties
84 and can inhibit the germination or growth of other plant species (Aguilera et al., 2010;
85 Gerber et al., 2008) as well as bacteria (Hedenec et al., 2014) with mixed effects on fungi
86 (e.g. Daayf et al., 1995; Kumagai et al., 2005). A study by Vastano *et al.* (2000) revealed a
87 higher concentration of stilbenes in North American invasive *F. japonica* than in Chinese
88 native individuals of the same species tending to support the NVH in the case of knotweed.

89 One of these compounds, *trans*-resveratrol (3,4,5'-trihydroxystilbene), has been identified
90 as being produced by knotweed (Vastano et al., 2000). This molecule, which is also found in
91 grapevines, is known as having antifungal (Filip et al., 2003) and antibacterial properties
92 (Chan, 2002). Content analysis of resveratrol in knotweed tissues has been assessed by
93 Vaher & Koel (2003) who found that more than 80% of *trans*-resveratrol was located in the
94 roots and rhizomes, where the majority of plant-microorganisms interactions occur (Bais et
95 al., 2006).

96 Secondary metabolites present in knotweed rhizomes could have either a direct effect on
97 soil fauna either by repellence (Asplund et al., 2015), toxicity (Isman and Duffey, 1982) or an

indirect effect through changes in the soil biota (Ens et al., 2009). As evidence for direct toxicity of phenolic compounds is scarce, indirect effects through alterations of basal resources for secondary consumers appear more likely. In this paper, we studied the effect of knotweed rhizome aqueous extracts on the soil biota and fauna in order to provide additional information on the novel weapon hypothesis in this particular case. Indeed, while several studies have assessed knotweed allelopathic potential in invaded areas none, as far as we know, have considered the impact on the soil fauna in relation to this hypothesis. Therefore and based on the theory, we hypothesized that: (1) knotweed has a negative effect on microbial (and particularly bacterial) biomass through rhizome allelopathic secondary metabolites ; (2) this negative effect has repercussions on higher trophic levels through trophic cascades, and results in soil food web structure alteration ; (3) this negative effect is be slightly attenuated by a positive trophic effect of nutrient addition provided by knotweed rhizome extract ; (4) those effects, positive (i.e. trophic) or negative (i.e. allelopathic), are concentration-dependent.

2. Material & Methods

2.1. Material collection and experiment preparation

Belowground *F. japonica* biomass was harvested in early autumn 2016 within a spontaneously invaded plateau site in Normandy, France (49.455024° N; 1.062645° W). To the best of our knowledge, control measures have never been applied to this site. Samples were kept in an icebox for transportation to the laboratory. Rhizomes were water-cleaned and stored at 4°C prior to extraction. We used an electric grinder to break plant tissues and

facilitate osmosis. One hundred grams of ground plant material was mixed with 1000 ml of distilled water. This aqueous extract was kept at 19°C for 24h. Following Norsworthy (2003) the mixture was then passed through a series of sieves ranging from 1000 to 50 µm and then vacuum filtered through standard filter paper (> 20-25 µm). The extract was then further sterilized by filtering through 0.22 µm filter.

We collected soil from the upper 10 cm of a reaped grassland in a small valley. While the area was uninfested by *F. japonica* close-by sites (< 200 m) with similar topographical and edaphic conditions have been invaded for several years. Macrofauna as well as macroscopic plant materials were removed from the collected soil. Samples were gently, and unforcefully, sieved at 5 mm so as to preserve mesofauna and mixed. Ten 200 g samples were taken from the soil to sample initial Collembola, Acari and Nematoda communities. Soils were placed in 8 x 8 x 10 cm plastic pots. Filter paper (<10-20 µm) was placed at the bottom of the prevent leakage of the pot content. Sixty grams of fine grained sand was added above the filter paper forming a ~ 0.5 cm layer. The rest of the pot was filled with 310 ± 5 g of soil. Ten 400 g samples of mixed soil were also collected for analysis of initial physico-chemical conditions. A layer of 0.5 g of *Agropyron* sp. litter, which is the dominant species in the samples grassland and also present close to invaded sites, was added to provide physical habitat for the soil fauna. Pots were kept at 19°C in a phytotron with a 8h day / 16h night cycle for a week. In order to increase and homogenize abundance and compensate for possible losses during soil sieving each pot was then placed under 2 individual Berlese-Tullgren extractors, one containing topsoil (0 – 5 cm) and the other deeper soil (5 – 10 cm) from the same area.

2.2. Experimental design

To simulate varying natural conditions and test for concentration-dependence distilled water was used to provide different concentrations of the aforementioned aqueous KRE (0, 33, 66 & 100%). Half of the solution at each concentration, including distilled water, was filtered 3 times through activated carbon prior to watering. This filtration was conducted in order to remove potentially toxic organic compounds (Cheremisinoff and Ellerbusch, 1978) from the KRE. We also filtered distilled water in order to test for the effect of filtration itself. In total we obtained 8 different solution: filtered and unfiltered distilled water, filtered and unfiltered 33% KRE, filtered and unfiltered 66% KRE as well as filtered and unfiltered 100% KRE. Each solution was used to water 10 pots prepared as detailed above. The result was a balanced factorial design ($4 \times 2 \times 10 = 80$ pots with 10 replicates per modality). During the experiment, the pots were kept for four weeks (from early November to early December 2016) in a climate-controlled room ($21.0 \pm 1.9^\circ\text{C}$, 16h day / 8h night, $47.0 \pm 8.8\%$ humidity) and watered weekly with the corresponding solution.

2.3. Sampling, biochemical analysis and fauna identification

In order to verify the validity of our activated-carbon methodology we used HPLC to test for resveratrol concentration in filtered and unfiltered KRE. Resveratrol ($\text{C}_{14}\text{H}_{12}\text{O}_3$), a phenolic allelopathic compound, was measured using direct-injection high-performance liquid chromatography (ThermoFisher Scientific UltiMate 3000 UHPLC). We used a variable wavelength UV detector at 306 nm, equipped with a standard C18 column, a water-acetonitril (60:40) mobile phase and an isocratic flow of $1 \text{ ml} \cdot \text{min}^{-1}$ (Goldberg et al., 1994). We used commercially-available Resveratrol powder (CAS Number: 501-36-0) for calibration. At the end of the experiment several biotic and abiotic variables were assessed. Approximately 100g of fresh soil was used for springtail extraction in a Berlese-Tullgren funnel (Macfadyen, 1961). Samples were weighted and placed within sieves (stitch: 1 mm,

diameter: 80 mm, height: 50 mm) above a plastic funnel. Extraction, under a heat source, lasted for a week with individuals collected in 70% ethanol. This extraction method is dependent on the limited tolerance of these animals to desiccation and will therefore only extract active individuals. There is therefore no differentiation between individuals that were inactivated, killed or otherwise incapacitated. One hundred grams of fresh soil was used for nematode extraction in a Baermann funnel (McSorley and Walter, 1991). Dampened samples were placed in a porous paper (10-15 µm stitch) supported by a 2 mm sieve and placed above a water-filled and sealed funnel for 48h. This method has a limited efficiency in isolating slow moving and nematodes and will not isolate inactive individuals (Van Bezooijen, 2006) and thus is not exhaustive.

Mesofauna samples were separated into Acari and Collembola under a stereo binocular microscope. Collembola individuals were mounted in lactic acid on microscope slides for identification with a phase-contrast optical microscope. Collembola individuals were identified to the species level (Hopkin, 2007; Potapov, 2001; Thibaud, 2004). Acari were identified to the order or suborder level: Mesostigmata (Gamasida), Cryptostigmata (Oribatida) and Prostigmata (Actinedida) (Coineau and Cleva, 1997). The cohort Astigmatina (previously the suborder Astigmata) were included in the suborder Oribatida (Wang and Fan, 2010). After Baermann funnel extraction, nematodes were counted while active under a stereo binocular microscope. Following decantation nematodes were fixed using a 4% formaldehyde solution and mounted on microscope slides. Individuals were attributed to trophic groups (herbivores, bacterivores, fungivores and predators/omnivores) based on mouthpart examination under a compound optical microscope.

Four grams of fresh soil were used to measure soil ergosterol content using the method proposed by Gong, Guan & Witter (2001). Ergosterol is a sterol found within fungi and protozoa. Ergosterol concentration can be used as a proxy of soil fungal biomass. Two times

20 g of fresh soil was used to assess microbial biomass using chloroform fumigation and extraction (Brookes et al., 1985). Carbon extraction was performed in 100 ml of potassium sulphate 0.05M (K_2SO_4) for chloroform-fumigated or unfumigated samples.

Once springtail extraction was complete a 30 g dry soil aliquot used to assess soil abiotic variables. Soil pH was measured using 1:5 volumetric fraction in 1M potassium chloride (KCl) using a Mettler Toledo FiveEasy pH meter. Total carbon and nitrogen content was measured in a ThermoFisher Flash Analyzer 2000 after electric grinding of dry soil material.

2.4. Statistical analysis

We used ANOVA tests included in R Software 3.3.1 for statistical analysis. In order to assess for knotweed rhizome extracts (KRE) effect on the soil fauna and microbiology we calculated the relative differences between the considered modalities and our control by standardizing and normalizing our measured values in relation to control means. This was done by subtracting the average control value to each value for the considered modality. Based on our methodology and hypotheses we considered that several of the potential effects of KRE could be separated. We thus considered that subtracting the control mean to values obtained in modalities where KRE was unfiltered by activated carbon (AC) values gave the combined effect of KRE on the considered variable (Fig. 1). Based on the generally accepted hypothesis we considered that AC filtration retains knotweed secondary compounds. Therefore, subtracting control means to values measured in pots subjected to AC filtered KRE we calculated KRE effect not attributable to secondary metabolites (Fig. 1). We referred to this effect as a trophic effect that could directly affect soil microbiology by adding nutrients to the substrate with potential indirect repercussions on higher trophic levels.

Finally, the same methodology was used by subtracting for each concentration the mean AC filtered value to individual values measured in pots that received unfiltered KRE.

We used a null model approach in order to consider control stochastic variability. Null vectors were randomly generated for each variable based on the observed distribution parameters of the control. Significant differences between these null vectors and our data vectors was assessed using non-parametric Wilcoxon signed-rank test. An iterative procedure (one-thousand repetitions) and result aggregation enabled us to robustly, and conservatively, assess for statistical differences between control and treatments. These differences, when found, showed KRE effects on the considered variables.

We also assessed for differences in the structure of the soil food web with and without AC filtration, and thus with or without knotweed secondary metabolites. We used multigroup path analysis to model our empirically observed model and compare it to a “null” multigroup model. The “null” model had constrained intercepts and regression coefficients that was compared to the empirical model using ANOVA. This approach provides a means to assess for covarying responses of soil fauna compartments to potential allelopathy.

3. Results

3.1. Physico-chemistry & microbiology

Contrary to our hypothesis and the literature, we observed limited effect of knotweed rhizome extracts (KRE) on microbiological variables. There were no significant differences in ergosterol concentration, an indicator of fungal biomass, irrespective of concentration or filtration mainly due to high variability. Regarding microbial carbon, a proxy of overall

microbial biomass, nutrient addition seems to cause a decrease as concentration increases but insignificantly except at intermediate concentrations ($-10.74 \pm 4.04\%$; $p < 0.05$; Tab. 1) with repercussions on ergosterol / microbial ratio (17.25 ± 6.06 ; $p < 0.05$; Tab. 1). pH was also affected by KRE addition with a significant decrease in response to secondary metabolites addition at the lowest concentration ($-0.02 \pm 0\%$; $p < 0.05$; Tab. 1) which shifted to an increase at the highest concentration ($+1.38 \pm 0.5$; $p < 0.05$; Tab. 1). The C/N ratio remained largely unaffected by KRE input except at the highest concentration (combined effect: $-1.37 \pm 0.56\%$; $p < 0.05$; Tab. 1).

Resveratrol concentration for 100% v/v KRE was $2.27 \pm 0.23 \text{ mg.l}^{-1}$ while the literature suggests an IC_{50} (i.e. concentration for 50% mortality) of 9 mg.l^{-1} (Fan et al., 2010). We did not detect resveratrol in any detectable amount in KRE samples after activated carbon filtration, even at the highest concentrations. Several chromatograms detailing these analysis are provided in Supplementary Material A1.

3.2. Nematodes

Total nematode abundance showed a strong response to KRE input at all concentrations despite important differences in effect direction. For instance, total nematode abundance was reduced by half following addition of low-concentration KRE (Fig. 2; Tab. 2). Conversely, nematode abundance was almost doubled at the highest KRE concentration while no response was found at the intermediate concentration (Fig. 2; Tab. 2). This general trend (i.e. combined KRE effect) is the result of decreasing intensity and significance of responses to nutrient addition (from +86 % at minimum concentration to +21 % at maximum concentration; Fig. 2; Tab. 2) and highly contextual responses of nematodes to secondary

compounds addition (from -43 % at intermediate concentration to +48 % at maximum concentration; Fig. 2; Tab. 2).

Although significance and intensity differed, bacterivorous and fungivorous nematodes (41.07 ± 1.59 % and 22.35 ± 1.26 % of total nematodes abundance, respectively) response to KRE addition varied similarly to the general trend with a shift from a negative (-43.5 % / -79.8 %) to a positive (+58.3 % / +24.5 %) response with increasing KRE concentration (Tab. 2). Herbivorous nematodes abundance (28.0 ± 1.3 % of total abundance) varied somewhat differently with a significant, and positive, response to KRE addition only at the highest concentration (+177 % increase; Tab. 2). Nutrient addition appeared to elicit a generally positive response independently of concentration ($+64.8 \pm 17.6$ %) which was only significant at the highest concentration ($+59.2 \pm 21.3$ %; Tab. 2) while there was no significant response to secondary compounds addition. Predatory and omnivorous nematodes showed no significant response to KRE addition (Tab. 2).

3.3. Mesofauna

Total Acari relative abundance showed a strong positive response to KRE input at all concentrations without significant differences in intensity (from +138.1 to +223.8 % as concentration increased; Fig. 3) which appears to me mostly related to a response to secondary compounds addition. A similar pattern was observed for oribatid mites (Tab. 3). There were significant differences between responses at the lowest and highest concentration levels for both the combined and secondary compounds responses but not for response to nutrient addition (Tab. 3). Predators (i.e. mostly Gamasida) abundances responded only to the combined aspects of KRE addition, and only at the two lowest concentrations (Tab. 3).

Regarding Collembola abundance observed response patterns are positive although only
significant when considering combined KRE effects at intermediate and high concentration
(+93.9 and +66.2 %, respectively; Fig. 4). A positive, yet insignificant, effect of nutrient
addition seems to exist at the intermediate concentration ($+56.9 \pm 28.4$ %; $p < 0.10$; Tab. 3).
Taxonomic diversity (i.e. Shannon's diversity) responded positively to KRE addition (only
significant at the two lowest concentrations). A response (positive) to secondary compounds
was only found at the intermediate KRE concentration (Tab. 3). Collembola functional
richness and evenness, calculated using trait data from the COLTRAIT database (Salmon et
al., 2014; Salmon and Ponge, 2012), only responded at 66% v/v KRE, mainly linked to the
secondary allelopathic effect of KRE addition. Functional evenness decreased while
functional richness increased in both cases (Tab. 3).

3.4. Path analysis

Differences between the empirically observed and a "null" multigroup model with
constrained intercepts and regression coefficients was tested using ANOVA. It showed a
difference in model structure between the two groups (i.e. unfiltered and activated carbon
filtered KRE addition) ($n_{group1} = 40$, $n_{group2} = 40$, Chi^2 difference = 39.87, $p = 0.022$).
Allelopathic effect removal through activated carbon (AC) filtration increased the strength
of the relationship between KRE concentration and microbial carbon concentration (From -
0.005 $p = 0.816$ to -0.032 $p = 0.048$; Fig. 5) and the effect of the latter on Collembola
abundance (-6.998 $p = 0.384$. to -14.443 $p = 0.060$; Fig. 5), herbo-fungivorous Acari
abundance (4.217 $p = 0.530$. to -9.884 $p = 0.075$; Fig. 5) and bacterivorous nematodes
abundance (0.864 $p = 0.941$. to -25.575 $p = 0.027$; Fig. 5). The relationship between
bacterivorous and predatorous nematodes abundance remained unaffected by AC filtration

(0.126 $p = 0.000$. to 0.111 $p = 0.000$; Fig. 5). The relationship between herbo-fungivorous and predatorous Acari abundance was not significantly altered by AC filtration (0.030 $p = 0.593$. to 0.089 $p = 0.176$; Fig. 5).

Concerning the fungal pathway allelopathic effect removal through activated carbon (AC) filtration decreased the relationship, yet with still no significant relationship, between KRE concentration and fungal biomass (i.e. ergosterol concentration) (0.013 $p = 0.174$. to -0.007 $p = 0.571$; Fig. 5). Fungal biomass relationships with its consumers was also affected by allelopathic effect removal: with herbo-fungivorous Acari (18.681 $p = 0.166$. to -3.119 $p = 0.693$; Fig. 5), Collembola abundance (-1.464 $p = 0.928$. to 17.470 $p = 0.118$; Fig. 5) and, in a very limited way, fungivorous nematodes abundance (-1.973 $p = 0.836$. to 8.794 $p = 0.274$; Fig. 5). Strengths of relationships of these taxa with their predators also changed after allelopathic effect removal especially fungivorous nematodes (-0.213 $p = 0.011$ to -0.067 $p = 0.348$; Fig. 5) and Collembola (0.126 $p = 0.009$ to -0.050 $p = 0.282$; Fig. 5) with predatorous Acari. The relationship between fungivorous and predatorous nematodes abundance was mostly unaffected (-0.030 $p = 0.593$ to 0.089 $p = 0.176$; Fig. 5) as was the relationship between the two main predatorous groups (i.e. predatorous nematodes and Acari) (0.244 $p = 0.161$ to -0.119 $p = 0.496$; Fig. 5).

4. Discussion

The first hypothesis posited a negative effect of knotweed allelopathic secondary compounds (ASC) on microbial communities as generally observed *in situ* (Hedenec et al., 2014; Tamura and Tharayil, 2014). The effect of ASC addition was not directly tested on the soil fauna, due in part to the lack of a proper identification of all such potential compounds in knotweed (Fan et al., 2010). However, we were able to ascertain the removal of ASC from

a solution of knotweed rhizome extract (KRE) by using activated carbon which is known to suppress allelopathic effects (Ridenour & Callaway, 2001 but see Lau et al., 2008 for a critic of this methodology). We were able to test and demonstrate removal of one ASC, trans-resveratrol, from KRE through activated carbon filtration (see Supplementary Material A1). Differences in population responses between activated carbon filtered and unfiltered KRE addition to the soil was therefore considered to be mainly, but not only, due to removal of ASC. We then indirectly calculated a “secondary” effect of KRE. Contrary to our hypothesis, we generally did not find any significant antimicrobial effects on ergosterol concentration (a proxy of fungal biomass; Davis & Lamar, 1992), microbial carbon (a proxy of microbial biomass; Vance et al., 1987) or the ratio between the two (a rough indicator of microbial community structure; Djajakirana et al., 1996; Tab. 2). Ergosterol concentration itself remained unaffected by KRE at concentrations which is consistent with the contrasting, yet often positive, effects found in the literature on the effects on fungal biomass (Daayf et al., 1995; Lecerf et al., 2007; Tamura and Tharayil, 2014). The pro-microbial, albeit insignificant, effects are far more surprising and tend to refute our hypothesis and contrast with results found in the literature (Hedenec et al., 2014; Kumagai et al., 2005; Stefanowicz et al., 2016; Tamura and Tharayil, 2014). Most of these results were observed in the field, with multiple potential confounding factors, with only Daayf et al., 1995 and Kumagai et al., 2005 directly testing antimicrobial and antifungal properties of knotweeds secondary compounds in controlled conditions. For instance, a major source of knotweed allelopathic properties are linked to the slow degradation and release of phenolic compounds from leaf litter degradation (Lavoie, 2017) which we did not account for in this study. Overall while the antibacterial effect of knotweeds in general, and Japanese knotweed in particular, appear fairly conclusive in the literature our results tend to show that this cannot be attributed to, or only to, rhizome secondary metabolites.

The final hypothesis stipulated that knotweed KRE-addition effects, in particular ASC addition, were linearly concentration-dependent. This hypothesis cannot be properly segregated from the other and will be considered here to avoid repetition. This hypothesis was based on the literature which frequently mentions release of secondary metabolites in the environment by knotweeds as a major contributor to knotweed effects in their invasive range (Vastano et al., 2000). Other lab studies have shown concentration-dependent effects of some root-secreted phenolic compounds on microbial biomass (e.g. Zhang et al., 2015). These compounds; however, are not in the same family as resveratrol or catechin. We tested this hypothesis indirectly as the effect of secondary metabolites was calculated and not measured. Concerning the concentration-dependence of microbiological response to ASC the relationship remained insignificant in all cases for both fungal and microbial biomass (Tab. 1). Hence, while there may be concentration-dependence the effects themselves are insignificant, and we must therefore accept the alternative hypothesis in the case of ASC. Nutrient addition, however, significantly and negatively affected microbial biomass at 66% v/v concentration only, showing differences dependent on concentration. However, as this is for intermediate concentration there does not appear to be a linear relationship between concentration and response. When analyzing changes in soil food web structure we showed that allelopathic secondary compounds (ASC) removal increased the strength of the negative correlation between KRE concentration and microbial biomass, the corollary being that ASC addition would tend to decrease the strength of the relationship between the two (Fig. 5). Only after ASC removal was the relationship significant between the two variables. Therefore, while the combined, and ASC, effects on microbial biomass appear not to be concentration-dependent purely “trophic” effects are significantly so.

The second hypothesis stated that, if there were antimicrobial and antifungal effects of KRE, they would have repercussions on higher trophic levels through a trophic cascade with potential alterations of trophic structures. We showed previously that no antimicrobial or

antifungal effect were found in our experiment in response to KRE of allelopathic secondary compounds (ASC). However, the evidence revealed significant differences in abundance at higher trophic levels. Indeed, we did find significant effects of KRE on bacterivorous and fungivorous nematodes (both positive and negative) abundances as well as herbo-fungivorous Acari and Collembola abundances (mostly positive) from both allelopathic or combined effects (although with no consistent pattern at different concentrations; see Tab 2-3). We found no significant differences in predator nematodes abundances despite the changes in abundance of their prey. Predator Acari, on the other hand, had higher abundances following KRE addition. Accordingly, predators may have had a top-down effect on microbivores while being unaffected themselves by bottom-up regulation themselves in the considered timeframe. Unfortunately, we could not find any reports in the literature on nematodes abundances under knotweed and therefore cannot assess the representativity of our results. Skubala & Mierny (2009), on the other hand, found a significant negative effect of giant knotweed (*Fallopia sachalinensis*) on oribatid mites (mostly herbivores, fungivores or both) but no effect on Collembola abundance (mostly generalist fungivores and detritivores). They attributed observed effects in the field on liberation phenolic compounds from leaf litter degradation, not rhizome excretion. This is, to our knowledge, the only publication to date assessing knotweed effects on the soil mesofauna in spontaneously invaded sites. Hedenec et al. (2014) also considered Collembola and Acari, with mixed results and no differences with native species, but in an agricultural setting with giant knotweed used as a biofuel crop. Macroarthropod abundance, which we did not consider in this study, has generally been shown to be negatively affected by knotweed presence (Gerber et al., 2008; Kappes et al., 2007; Topp et al., 2008)

Collembola and Acari have a generation time of several weeks to months, depending on taxa and eco-morphological group (Choi et al., 2002; Ermilov and Lochynska, 2008; Joosse and Veltkamp, 1969; Park, 2007; Prinzing et al., 2002; Verhoef and Selm, 1983). Observed

differences are thus unlikely to be caused by a predator-prey intergenerational regulation but would be due to a more direct effect. Predatory nematodes and Acari have generation times ranging from 3 to 280 days depending on taxa (Abou-Awad et al., 2001; Khan et al., 2007; Ydergaard et al., 1997) as well as temperature. For some taxa within our study the experiment duration, 1 month, may have been insufficient to observe significant repercussions on higher trophic levels through intergenerational predator-prey relationships.

We also assessed how activated carbon filtration, and thus ASC removal, affected the structure of the soil trophic network by using multigroup path analysis (Fig. 5). The comparison of an empirical model to a constrained model, clearly showed that ASC removal significantly altered overall relationships between the various considered faunal groups. In this analysis, ASC removal also seemed to more readily alter relationships between taxa in the microbial food web (i.e. between microbial biomass and abundances of Collembola, oribatid mites and bacterivorous nematodes). This relationship was, however, negative. This would suggest that ASC removal from added KRE increases the interdependency of compartments/taxa within the microbial food web as well as the effect of KRE concentration on microbial biomass. The corollary, although we did not test it directly, would be that the allelopathic component of knotweed effect tends to limit between-group variability. In addition, it would appear that while nutrient input has concentration-dependent effects (at least on microbial biomass), nutrient and allelopathic effects are not dependent on concentration. Another striking feature is the change in the relationship between fungivorous nematodes and predatorous Acari, for which intensity was drastically reduced by ASC removal from KRE and most importantly switched from a negative to a positive correlation.

We also posited that part of knotweeds success as invaders was due to lack of adaptation by native species to the invader: the novel weapon hypothesis (Callaway and Ridenour, 2004). This framework most generally applies to other plant species with which there is more direct competition. While our methodology does not enable us to assess that effect, we expected negative effects (on abundance and diversity) of knotweed allelopathy on at least some taxa within the soil fauna which would help explain results found in several field studies (Gerber et al., 2008; Kappes et al., 2007; Skubala and Mierny, 2009; Topp et al., 2008). Such negative effects could have been direct through phytotoxicity or indirect through a trophic cascade or changes in habitat structure. This was not the case here with few negative responses of soil fauna to ASC addition. The only significant negative effects documented were for all types of nematodes in response to 66% KRE addition (Tab. 2).

The third hypothesis centered on the posited countering of negative allelopathic effects of KRE addition by nutrient addition. This appears to be the case, albeit insignificantly, for the ergosterol / microbial C ratio at the two highest concentrations which is negatively affected by the allelopathic component of KRE (i.e. favoring microbial biomass) and positively affected by the nutrient-addition component (i.e. favoring fungal biomass) with a combined positive effect. Field studies evaluating fungal:bacterial ratios have found conflicting results on that matter with both increased (Suseela et al., 2016; Tamura and Tharayil, 2014) and decreased (Stefanowicz et al., 2016) ratios under knotweed-invaded plots. The results tend to support the first case of decreased bacterial biomass, although fungal biomass remained unaffected (Tab. 1). These effects of knotweed are often attributed to increased litter biomass (e.g. Suseela et al., 2016) and changes in litter chemistry (higher litter C/N ratios: Dassonville, Guillaumaud, Piola, Meerts, & Poly, 2011; Mincheva et al., 2014; Urgenson, Reichard, & Halpern, 2009; higher litter lignin content: Aguilera et al., 2010) in addition to the already mentioned allelopathic effects. Our decomposition of effects between nutrient and secondary compounds addition tends to indicate that this positive effect on

471 fungal:microbial ratios is mainly attributed to a positive response of microbial biomass to
472 increased nutrient input to the soil. Due to lack of leaf litter this is not comparable to field
473 nutrient input, but nonetheless worth considering. Secondary compounds, which can be
474 twice as concentrated in knotweed-invaded plots (Suseela et al., 2016), appear to have a
475 negative effect on fungi:microbial biomass ratios in this case. Changes in microbial biomass
476 carbon have the most influence over shifts in this ratio in our case (i.e. negative effect of
477 nutrient-addition and positive effect of secondary compounds addition) which would to be
478 contrary to our hypothesis. The pH responded in a similar manner with a significant negative
479 nutrient addition effect, a significant positive secondary compounds effect but an
480 insignificant slightly positive combined effect. A decrease in soil pH has generally been
481 observed in field studies (Dassonville et al., 2011, 2008; Kappes et al., 2007) and is also
482 attributed to increased litter biomass in knotweed stands. This is, however, not always the
483 case (Stefanowicz et al., 2017; Stoll et al., 2012)

484 Finally the results regarding microarthropod abundances (i.e. Collembola and Acari) are
485 ambiguous with no evidence of the hypothesized pattern of attenuating effects. In fact, all
486 significant responses of both Acari and Collembola are positive. This would seem, contrary
487 to our hypothesis, to indicate a synergetic combined effect of KRE addition (mostly at
488 intermediate and high concentrations). In all cases of significant response to a combined
489 effect one or both components' response was neutral (nutrient addition especially).

490 Contrasting and inverse results from nutrient and secondary metabolites addition have also
491 been found for nematodes at intermediate KRE concentration with a negative ASC addition
492 effect, positive nutrient addition effect and neutral combined effect. Direct nematocidal
493 effects of plant secondary compounds have been documented in some cases, mostly in
494 laboratory studies (Chitwood, 2002). If there was such a direct effect of knotweed we would
495 expect it to also be present at the highest concentration, which is not the case in our study
496 (Fig. 2).

These results provide the basis for further research on knotweed such as more detailed characterization of knotweed ASC and their potential allelopathic effects as well as further field work. As assessment of indirect, and direct, allelopathic effects of phenolic compounds on soil fauna has rarely been done. Thus, this report should provide useful data for authors working on such a subject as information is currently scarce on the subject. Finally, we hope the results presented here will provide useful reference data for future biological invasions study and inform managers of invaded areas on knotweed potential impacts.

5. Conclusion

In conclusion, the results showed an effect of knotweed rhizome extract (KRE) on soil microbiology. Fungal biomass remained unaffected but microbial biomass as a whole responded negatively to KRE in some cases. Interestingly these negative responses, when they occurred, were mostly attributable to factors other than the allelopathic secondary compounds (ASC) within KRE, most notably nutrient addition. Calculated responses of microbial biomass to ASC addition were, albeit insignificantly so, positive. While KRE addition had an effect in most cases on taxa “higher” within the soil trophic networks, there were no evident and generalizable trophic cascades across trophic levels for a given KRE concentration. Path analysis did reveal important changes in soil food web structure (constructed based on hypothesized producer-consumer relationships) which appeared to be mostly within the bacterial pathway, and concentration-dependent. There was circumstantial, but not generalizable, evidence of compensating, or attenuating, effects of nutrient and ASC addition on various taxa. Rarely was ASC effect, when documented, concentration-dependent in the results.

6. Acknowledgements

We would like to thank the Région Normandie and the Grand Réseau de Recherche (GRR) Territoire, Environnement, Risques, Agronomie (TERA) for financial support for this experiment. We would also like to thank the Agence de l'Eau Seine-Normandie for their support. We also thank the two anonymous reviewers who provided exhaustive and constructive comments that significantly improved the paper. We would also like to thank Brad Schneider for proof-reading the manuscript as English is not our native language. Finally, we thank members of Ecodiv for their theoretical and practical help during this study.

7. References

- Abgrall, C., Chauvat, M., Langlois, E., Hedde, M., Mouillot, D., Salmon, S., Winck, B., Forey, E., 2017. Shifts and linkages of functional diversity between above- and below-ground compartments along a flooding gradient. *Functional Ecology* 31, 350–360.
- Abou-Awad, B.A., Korayem, A.M., Hassan, M.F., Abou-Elela, M.A., 2001. Life history of the predatory mite *Lasioseius athiasae* (Acari, Ascidae) on various kinds of food substances: a polypeptide analysis of prey consideration. *Journal of Applied Entomology* 125, 125–130.
- Aguilera, A.G., Alpert, P., Dukes, J.S., Harrington, R., 2010. Impacts of the invasive plant *Fallopia japonica* (Houtt.) on plant communities and ecosystem processes. *Biological Invasions* 12, 1243–1252.
- Asplund, J., Bokhorst, S., Kardol, P., Wardle, D.A., 2015. Removal of secondary compounds increases invertebrate abundance in lichens. *Fungal Ecology* 18, 18–25.

546 Bais, H.P., Weir, T.L., Perry, L.G., Gilroy, S., Vivanco, J.M., 2006. The role of root exudates in
547 rhizosphere interactions with plants and other organisms. *Annual Review of Plant*
548 *Biology* 57, 233–266.

549 Bassett, I., Paynter, Q., Beggs, J.R., 2011. Invasive *Alternanthera philoxeroides* (alligator
550 weed) associated with increased fungivore dominance in Coleoptera on decomposing
551 leaf litter. *Biological Invasions* 13, 1377–1385.

552 Brookes, P.C., Landman, A., Pruden, G., Jenkinson, D.S., 1985. Chloroform fumigation and
553 the release of soil nitrogen: A rapid direct extraction method to measure microbial
554 biomass nitrogen in soil. *Soil Biology and Biochemistry* 17, 837–842.

555 Callaway, R.M., Cipollini, D., Barto, K., Thelen, G.C., Hallett, S.G., Prati, D., Stinson, K.,
556 Klironomos, J., 2008. Novel weapons: Invasive plant suppresses fungal mutualists in
557 America but not in its native europe. *Ecology* 89, 1043–1055.

558 Callaway, R.M., Ridenour, W.M., 2004. Novel weapons: invasive success and the evolution of
559 increased competitive ability. *Frontiers in Ecology and the Environment* 2, 436–443.

560 Chan, M.M.-Y., 2002. Antimicrobial effect of resveratrol on dermatophytes and bacterial
561 pathogens of the skin. *Biochemical Pharmacology* 63, 99–104.

562 Cheremisinoff, P.N., Ellerbusch, F., 1978. Carbon adsorption handbook. Ann Arbor Science
563 Publishers.

564 Chitwood, D.J., 2002. Phytochemical based strategies for nematode control. *Annual Review*
565 *of Phytopathology* 40, 221–249.

566 Choi, W. Il, Ryoo, M. Il, Kim, J.-G., 2002. Biology of *Paronychiurus kimi* (Collembola:
567 Onychiuridae) under the influence of temperature, humidity and nutrition.
568 *Pedobiologia* 46, 548–557.

569 Coineau, Y., Cleva, R., 1997. Ces animaux minuscules qui nous entourent. Delachaux et
570 Niestlé.

571 Daayf, F., Schmitt, A., Bélanger, R.R., 1995. The effects of plant extracts of *Reynoutria*
572 *sachalinensis* of powdery mildew development and leaf physiology of long english
573 cucumber. *Plant Disease* 79, 577–580.

574 Dassonville, N., Guillaumaud, N., Piola, F., Meerts, P., Poly, F., 2011. Niche construction by
575 the invasive Asian knotweeds (species complex *Fallopia*): impact on activity,
576 abundance and community structure of denitrifiers and nitrifiers. *Biological Invasions*
577 13, 1115–1133.

578 Dassonville, N., Vanderhoeven, S., Vanparys, V., Hayez, M., Gruber, W., Meerts, P., 2008.
579 Impacts of alien invasive plants on soil nutrients are correlated with initial site
580 conditions in NW Europe. *Oecologia* 157, 131–140.

581 Davis, M.W., Lamar, R.T., 1992. Evaluation of methods to extract ergosterol for quantitation
582 of soil fungal biomass. *Soil Biology and Biochemistry* 24, 189–198.

583 De Waal, L.C., 2001. A viability study of *Fallopia japonica* stem tissue. *Weed Research* 41,
584 447–460.

585 Djajakirana, G., Joergensen, R.G., Meyer, B., 1996. Ergosterol and microbial biomass
586 relationship in soil. *Biology and Fertility of Soils* 22, 299–304.

587 Ens, E.J., French, K., Bremner, J.B., 2009. Evidence for allelopathy as a mechanism of
588 community composition change by an invasive exotic shrub, *Chrysanthemoides*
589 *monilifera* spp. *rotundata*. *Plant and Soil* 316, 125–137.

590 Ermilov, S.G., Lochynska, M., 2008. The influence of temperature on the development time
591 of three oribatid mite species (Acari, Oribatida). *North-Western Journal of Zoology* 4,

592 247–281.

593 Fan, P., Hostettmann, K., Lou, H., 2010. Allelochemicals of the invasive neophyte *Polygonum*
594 *cuspidatum* Sieb. & Zucc. (Polygonaceae). Chemoecology 20, 223–227.

595 Filip, V., Plocková, M., Šmidrkal, J., Špičková, Z., Melzoch, K., Schmidt, Š., 2003. Resveratrol
596 and its antioxidant and antimicrobial effectiveness. Food Chemistry 83, 585–593.

597 Forey, E., Coulibaly, S.F.M., Chauvat, M., 2015. Flowering phenology of a herbaceous species
598 (*Poa annua*) is regulated by soil Collembola. Soil Biology and Biochemistry 90, 30–33.

599 Gerber, E., Krebs, C., Murrell, C., Moretti, M., Rocklin, R., Schaffner, U., 2008. Exotic invasive
600 knotweeds (*Fallopia* spp.) negatively affect native plant and invertebrate assemblages
601 in European riparian habitats. Biological Conservation 141, 646–654.

602 Goldberg, D.M., Yan, J., Ng, E., Diamandis, E.P., Karumanchiri, A., Soleas, G., Waterhouse,
603 A.L., 1994. Direct Injection Gas Chromatographic Mass Spectrometric Assay for trans-
604 resveratrol. Analytical Chemistry 66, 3959–3963.

605 Gong, P., Guan, X., Witter, E., 2001. A rapid method to extract ergosterol from soil by
606 physical disruption. Applied Soil Ecology 17, 285–289.

607 Hättenschwiler, S., Coq, S., Barantal, S., Handa, I.T., 2011. Leaf traits and decomposition in
608 tropical rainforests: Revisiting some commonly held views and towards a new
609 hypothesis. New Phytologist 189, 950–965.

610 Hättenschwiler, S., Vitousek, P.M., 2000. The role of polyphenols in terrestrial ecosystem
611 nutrient cycling. Trends in Ecology and Evolution 15, 238–242.

612 Hedenec, P., Novotny, D., Ustak, S., Cajthaml, T., Slejska, A., Simackova, H., Honzik, R.,
613 Kovarova, M., Frouz, J., 2014. The effect of native and introduced biofuel crops on the
614 composition of soil biota communities. Biomass and Bioenergy 60, 137–146.

615 Hopkin, S.P., 2007. A key to the Collembola (springtails) of Britain and Ireland. FSC
 616 Publications.

617 Inderjit, Evans, H., Crocoll, C., Bajpai, D., Kaur, R., Feng, Y.L., Silva, C., Treviño Carreón, J.,
 618 Valiente-Banuet, A., Gershenzon, J., Callaway, R.M., 2011a. Volatile chemicals from leaf
 619 litter are associated with invasiveness of a neotropical weed in Asia. *Ecology* 92, 316–
 620 324.

621 Inderjit, Wardle, D.A., Karban, R., Callaway, R.M., 2011b. The ecosystem and evolutionary
 622 contexts of allelopathy. *Trends in Ecology and Evolution* 26, 655–662.

623 Isman, M.B., Duffey, S.S., 1982. Toxicity of tomato phenolic-compounds to the fruitworm,
 624 *Heliothis-Zea*. *Entomologia Experimentalis et Applicata* 31, 370–376.

625 Joosse, E.N.G., Veltkamp, E., 1969. Some aspects of growth, moulting and reproduction in
 626 five species of surface dwelling Collembola. *Netherlands Journal of Zoology* 20, 315–
 627 328.

628 Kappes, H., Lay, R., Topp, W., 2007. Changes in different trophic levels of litter-dwelling
 629 macrofauna associated with giant knotweed invasion. *Ecosystems* 10, 734–744.

630 Khan, Z., Kim, Y.H., Kim, S.G., Kim, H.W., 2007. Observations on the suppression of root-knot
 631 nematode (*Meloidogyne arenaria*) on tomato by incorporation of cyanobacterial
 632 powder (*Oscillatoria chlorina*) into potting field soil. *Bioresource Technology* 98, 69–73.

633 Kohyt, J., Skubała, P., 2013. Communities of mites (Acari) in litter and soil under the invasive
 634 red oak (*Quercus rubra* L.) and native pedunculate oak (*Q. robur* L.). *Biological Letters*
 635 50, 111–124.

636 Kumagai, H., Kawai, Y., Sawano, R., Kurihara, H., Yamazaki, K., Inoue, N., 2005. Antimicrobial
 637 substances from rhizomes of the giant knotweed *Polygonum sachalinense* against the

638 fish pathogen *Photobacterium damsela* subsp. *piscicida*. Zeitschrift Für
639 Naturforschung C 60, 39–44.

640 Lau, J. a., Puliafico, K.P., Kopshever, J. a., Steltzer, H., Jarvis, E.P., Schwarzländer, M., Strauss,
641 S.Y., Hufbauer, R.A., 2008. Inference of allelopathy is complicated by effects of
642 activated carbon on plant growth. New Phytologist 178, 412–423.

643 Lavoie, C., 2017. The impact of invasive knotweed species (*Reynoutria* spp.) on the
644 environment: review and research perspectives. Biological Invasions 19, 2319–2337.

645 Lecerf, A., Patfield, D., Boiché, A., Riipinen, M.P., Chauvet, E., Dobson, M., 2007. Stream
646 ecosystems respond to riparian invasion by Japanese knotweed (*Fallopia japonica*).
647 Canadian Journal of Fisheries and Aquatic Sciences 64, 1273–1283.

648 Litt, A.R., Cord, E.E., Fulbright, T.E., Schuster, G.L., 2014. Effects of Invasive Plants on
649 Arthropods. Conservation Biology 28, 1532–1549.

650 Lowe, S., Browne, M., Boudjelas, S., De Poorter, M., 2000. 100 of the world’s worst invasive
651 alien species: a selection from the global invasive species database. Invasive Species
652 Specialist Group Auckland, New Zealand.

653 Macfadyen, A., 1961. Improved funnel-type extractors for soil arthropods. The Journal of
654 Animal Ecology 30, 171–184.

655 McSorley, R., Walter, D.E., 1991. Comparison of soil extraction methods for nematodes and
656 microarthropods. Agriculture, Ecosystems and Environment 34, 201–207.

657 Mincheva, T., Barni, E., Varese, G.C., Brusa, G., Cerabolini, B., Siniscalco, C., 2014. Litter
658 quality, decomposition rates and saprotrophic mycoflora in *Fallopia japonica* (Houtt.)
659 Ronse Decraene and in adjacent native grassland vegetation. Acta Oecologica 54, 29–
660 35.

661 Murrell, C., Gerber, E., Krebs, C., Parepa, M., Schaffner, U., Bossdorf, O., 2011. Invasive
662 knotweed affects native plants through allelopathy. *American Journal of Botany* 98,
663 38–43.

664 Norsworthy, J.K., 2003. Allelopathic Potential of Wild Radish (*Raphanus raphanistrum*).
665 *Weed Science Society of America* 17, 307–313.

666 Parepa, M., Schaffner, U., Bossdorf, O., 2013. Help from under ground: soil biota facilitate
667 knotweed invasion. *Ecosphere* 4, art31.

668 Park, E.K., 2007. Effect of laboratory culture conditions on population growth of *Proisotoma*
669 *minuta* (Tullberg)(Collembola: Isotomidae). *Entomological Science* 10, 135–140.

670 Potapov, M., 2001. Synopses on Palaearctic Collembola: Isotomidae. *Abhandlungen Und*
671 *Berichte Des Naturkundemuseums Gorlitz* 73, 1–603.

672 Prinzing, A., Kretzler, S., Badejo, A., Beck, L., 2002. Traits of oribatid mite species that
673 tolerate habitat disturbance due to pesticide application. *Soil Biology and Biochemistry*
674 34, 1655–1661.

675 Pyšek, P., Jarošík, V., Hulme, P.E., Pergl, J., Hejda, M., Schaffner, U., Vilà, M., 2012. A global
676 assessment of invasive plant impacts on resident species, communities and
677 ecosystems: The interaction of impact measures, invading species' traits and
678 environment. *Global Change Biology* 18, 1725–1737.

679 Reigosa, M.J., Pedrol, N., Gonzalez, L., 2006. Allelopathy: A Physiological Process with
680 Ecological Implications. Springer Science & Business Media, Dordrecht, The
681 Netherlands.

682 Reinhart, K.O., Callaway, R.M., 2006. Soil biota and invasive plants. *New Phytologist* 170,
683 445–457.

684 Ridenour, W.M., Callaway, R.M., 2001. The relative importance of allelopathy in
685 interference: the effects of an invasive weed on a native bunchgrass. *Oecologia* 126,
686 444–450.

687 Salmon, S., Ponge, J.F., 2012. Species traits and habitats in springtail communities: A
688 regional scale study. *Pedobiologia* 55, 295–301.

689 Salmon, S., Ponge, J.F., Gachet, S., Deharveng, L., Lefebvre, N., Delabrosse, F., 2014. Linking
690 species, traits and habitat characteristics of Collembola at European scale. *Soil Biology*
691 *and Biochemistry* 75, 73–85.

692 Skubala, P., Mierny, A., 2009. Invasive *Reynoutria* taxa as a contaminant of soil. Does it
693 reduce abundance and diversity of microarthropods and damage soil habitat?
694 *Pesticides* 1–4, 57–62.

695 Stefanowicz, A.M., Stanek, M., Nobis, M., Zubek, S., 2017. Few effects of invasive plants
696 *Reynoutria japonica*, *Rudbeckia laciniata* and *Solidago gigantea* on soil physical and
697 chemical properties. *Science of the Total Environment* 574, 938–946.

698 Stefanowicz, A.M., Stanek, M., Nobis, M., Zubek, S., 2016. Species-specific effects of plant
699 invasions on activity, biomass, and composition of soil microbial communities. *Biology*
700 *and Fertility of Soils* 52, 841–852.

701 Stinson, K.A., Campbell, S.A., Powell, J.R., Wolfe, B.E., Callaway, R.M., Thelen, G.C., Hallett,
702 S.G., Prati, D., Klironomos, J.N., 2006. Invasive Plant Suppresses the Growth of Native
703 Tree Seedlings by Disrupting Belowground Mutualisms. *PLoS Biology* 4, e140.

704 Stoll, P., Gatzsch, K., Rusterholz, H.P., Baur, B., 2012. Response of plant and gastropod
705 species to knotweed invasion. *Basic and Applied Ecology* 13, 232–240.

706 Suseela, V., Alpert, P., Nakatsu, C.H., Armstrong, A., Tharayil, N., 2016. Plant–soil

707 interactions regulate the identity of soil carbon in invaded ecosystems: implication for
708 legacy effects. *Functional Ecology* 30, 1227–1238.

709 Tamura, M., Tharayil, N., 2014. Plant litter chemistry and microbial priming regulate the
710 accrual, composition and stability of soil carbon in invaded ecosystems. *New*
711 *Phytologist* 203, 110–124.

712 Thibaud, J.-M., 2004. Synopses on Palaearctic Collembola: Hypogastruridae. *Abhandlungen*
713 *Und Berichte Des Naturkundemuseums Gorlitz* 75, 1–287.

714 Thorpe, A.S., Thelen, G.C., Diaconu, A., Callaway, R.M., 2009. Root exudate is allelopathic in
715 invaded community but not in native community: Field evidence for the novel
716 weapons hypothesis. *Journal of Ecology* 97, 641–645.

717 Topp, W., Kappes, H., Rogers, F., 2008. Response of ground-dwelling beetle (Coleoptera)
718 assemblages to giant knotweed (*Reynoutria* spp.) invasion. *Biological Invasions* 10,
719 381–390.

720 Urgenson, L.S., Reichard, S.H., Halpern, C.B., 2009. Community and ecosystem
721 consequences of giant knotweed (*Polygonum sachalinense*) invasion into riparian
722 forests of western Washington, USA. *Biological Conservation* 142, 1536–1541.

723 Vaher, M., Koel, M., 2003. Separation of polyphenolic compounds extracted from plant
724 matrices using capillary electrophoresis. *Journal of Chromatography A* 990, 225–230.

725 Van der Putten, W.H., Klironomos, J.N., Wardle, D.A., 2007. Microbial ecology of biological
726 invasions. *Isme J* 1, 28–37.

727 Van Kleunen, M., Weber, E., Fischer, M., 2010. A meta-analysis of trait differences between
728 invasive and non-invasive plant species. *Ecology Letters* 13, 235–245.

729 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil

730 microbial biomass C. *Soil Biology and Biochemistry* 19, 703–707.

731 Vastano, B.C., Chen, Y., Zhu, N., Ho, C.T., Zhou, Z., Rosen, R.T., 2000. Isolation and
732 identification of stilbenes in two varieties of *Polygonum cuspidatum*. *Journal of*
733 *Agricultural and Food Chemistry* 48, 253–256.

734 Verhoef, H.A., Selm, A.J. van, 1983. Distribution and population dynamics of Collembola in
735 relation to soil moisture. *Ecography* 6, 387–388.

736 Vilà, M., Espinar, J.L., Hejda, M., Hulme, P.E., Jarošík, V., Maron, J.L., Pergl, J., Schaffner, U.,
737 Sun, Y., Pyšek, P., 2011. Ecological impacts of invasive alien plants: a meta-analysis of
738 their effects on species, communities and ecosystems. *Ecology Letters* 14, 702–708.

739 Wang, Z.Y., Fan, Q.H., 2010. Psoroptidia (Acari: Astigmatina) of China: a review of research
740 progress. *Zoosymposia* 4, 260–271.

741 Wardle, D.A., 2002. *Communities and ecosystems: linking the aboveground and*
742 *belowground components*. Princeton University Press.

743 Wardle, D.A., Nilsson, M.C., Gallet, C., Zackrisson, O., 1998. An ecosystem-level perspective
744 of allelopathy. *Biological Reviews* 73, 305–319.

745 Ydergaard, S., Enkegaard, A., Brødsgaard, H.F., 1997. The predatory mite *Hypoaspis miles*:
746 temperature dependent life table characteristics on a diet of sciarid larvae, *Bradysia*
747 *paupera* and *B. tritici*. *Entomologia Experimentalis et Applicata* 85, 177–187.

748 Zhang, Z., Qiao, M., Li, D., Zhao, C., Li, Y., Yin, H., Liu, Q., 2015. Effects of two root-secreted
749 phenolic compounds from a subalpine coniferous species on soil enzyme activity and
750 microbial biomass. *Chemistry and Ecology* 31, 636–649.

751

753 Figures

754

755 Figure 1 : Diagram of calculations involved in separating knotweed rhizome extract (KRE) effects. CT: control, AC: activated carbon filtration, KRE:

756 no AC filtration.

758 Figure 2 : Relative nematode total abundance (%) compared to control in relation to knotweed rhizome extract dilution levels and activated carbon
759 filtration with decomposition of effects. Symbols indicate levels of significativity of repeated statistical testing of differences between calculated
760 values and null generated controls. n.s.: $p > 0.10$, ~ : $p < 0.10$, * : $p < 0.05$, ** : $p < 0.01$, *** : $p < 0.001$, **** : $p < 0.0001$.

762 Figure 3 : Relative Acari total abundance compared to control in relation to knotweed rhizome extract dilution levels and activated carbon filtration
763 with decomposition of effects. Symbols indicate levels of significativity of repeated statistical testing of differences between calculated values and
764 null generated controls. n.s.: $p > 0.10$, ~ : $p < 0.10$, * : $p < 0.05$, ** : $p < 0.01$, *** : $p < 0.001$, **** : $p < 0.0001$.

766 Figure 4 : Relative Collembola total abundance compared to control in relation to knotweed rhizome extract dilution levels and activated carbon
767 filtration with decomposition of effects. Symbols indicate levels of significativity of repeated statistical testing of differences between calculated
768 values and null generated controls. n.s.: $p > 0.10$, ~ : $p < 0.10$, * : $p < 0.05$, ** : $p < 0.01$, *** : $p < 0.001$, **** : $p < 0.0001$.

769

770 Figure 5 : Multigroup path model of soil mesofaunal food webs after filtered or unfiltered knotweed rhizome addition. Differences between the
 771 observed multigroup model and a “null model” with fixed Intercepts and Regressions was assessed with an ANOVA. Green arrows indicate a

772 positive correlation while red arrows indicate a negative correlation. Arrow width is proportional to the strength of the relationship. KRE concent. =
 773 knotweed rhizome extract concentration level, Fungi = ergosterol concentration, Microbial Biomass = carbon amount in microbial biomass, Fungiv.
 774 nemat. = Fungivorous nematodes abundance, Bacter. nemat. = Bacterivorous nematodes abundance, Predat. nemato. = Predatorous nematodes
 775 abundance, Herb.-Fung. Acari = Herbo-fungivorous Acari, Predat. Acari = Predatorous Acari, Collemb.= Collembola.

776

777

778

779 Tables

780

781 Table 1: Relative differences in physico-chemical and microbiological variables compared to control values (%) between knotweed rhizome extract
 782 exposed pots and control pots. Values are means +/- SE. P-values are from repeated Wilcoxon rank-sum tests on absolute relative differences.

783

Percentage	33% v/v	66% v/v	100% v/v
differences			

from control

	Combined	Nutrient	Secondary	Combined	Nutrient	Secondary	Combined	Nutrient	Secondary
Ergosterol	8.05 ± 6.97	1.88 ± 4.69	0.06 ± 0.07	4.75 ± 4.97	8.99 ± 4.99	-3.89 ± 4.56	3.86 ± 4.26	0.11 ± 7.16	3.75 ± 4.25
content	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
Microbial	-4.2 ± 6.11	-4.29 ± 6.18	0 ± 0.06	-7.78 ± 5.91	-10.74 ± 4.04	3.32 ± 6.63	-5.31 ± 6.55	-11.38 ± 8.87	6.84 ± 7.39
carbon	n.s.	n.s.	n.s.	n.s.	p < 0.05	n.s.	n.s.	n.s.	n.s.
Ergosterol /	10.98 ± 11.32	3.98 ± 7.48	0.07 ± 0.11	10.81 ± 7.52	17.25 ± 6.06	-5.5 ± 6.42	9.17 ± 9.67	24.16 ± 22.81	-12.07 ± 7.79
Microbial C	n.s.	n.s.	n.s.	n.s.	p < 0.05	n.s.	n.s.	n.s.	n.s.
pH (KCl)	-1.05 ± 0.48	1.39 ± 0.4	-0.02 ± 0	0.93 ± 0.46	0.39 ± 0.43	0.53 ± 0.46	0.48 ± 0.49	-0.89 ± 0.29	1.38 ± 0.5
	p < 0.10	p < 0.01	p < 0.05	p < 0.10	n.s.	n.s.	n.s.	p < 0.05	p < 0.05
C/N ratio	-0.18 ± 0.55	-0.52 ± 0.49	0 ± 0.01	0.11 ± 0.67	0.64 ± 0.6	-0.52 ± 0.67	-1.37 ± 0.56	-0.07 ± 0.57	-1.3 ± 0.57
	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	p < 0.05	n.s.	p < 0.10

784

785

786

787

788 Table 2: Relative differences in nematode abundances between knotweed rhizome extract exposed pots and control pots. Values are means
 789 percentages of difference +/- SE. P-values are from repeated Wilcoxon rank-sum tests.

Percentage

differences

33% v/v

66% v/v

100% v/v

from control

Combined

Nutrient

Secondary

Combined

Nutrient

Secondary

Combined

Nutrient

Secondary

Bacterivores

-43.52 ±

-27.64 ±

20.78

59 ± 34.27

-0.64 ± 0.13

15.05

54.43 ± 25.03

-53.14 ± 9.75

58.3 ± 21.49

12.19 ± 36.48

41.11 ± 19.15

	p < 0.10	n.s.	n.s.	n.s.	p < 0.10	p < 0.001	p < 0.05	n.s.	p < 0.10
							177.12 ±		
Herbivores	16.12 ± 23.1	69.23 ± 32.53	-0.31 ± 0.14	43.93 ± 33.55	66.01 ± 38.26	-13.3 ± 20.21	60.86	59.19 ± 21.25	74.09 ± 38.23
	n.s.	p < 0.10	n.s.	n.s.	n.s.	n.s.	p < 0.05	p < 0.05	p < 0.10
		138.03 ±		-49.79 ±		-52.93 ±		-16.06 ±	
Fungivores	-79.83 ± 5.26	43.19	-0.92 ± 0.02	12.07	6.67 ± 18.76	11.32	24.52 ± 28.52	18.97	48.35 ± 33.98
	p < 0.0001	p < 0.05	n.s.	p < 0.01	n.s.	p < 0.01	n.s.	n.s.	n.s.
	-18.82 ±	-39.64 ±				-35.81 ±			
Predators/omnivores	19.73	24.42	0.34 ± 0.33	-8.71 ± 28.14	42.23 ± 33.72	19.78	47.14 ± 35.15	34.69 ± 52.25	9.24 ± 26.1
	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.

790

791

792

793

794

795 Table 3: Relative differences in mesofauna (Collembola and Acari) abundance, taxonomic and functional indices between knotweed rhizome extract

796 exposed pots and control pots. Values are means +/- SE. P-values are from repeated Wilcoxon rank-sum tests.

Percentage differences

from control

33% v/v

66% v/v

100% v/v

Combined

Nutrient

Secondary

Combined

Nutrient

Secondary

Combined

Nutrient

Secondary

Acari

73.58 ± 30.4

24.53 ±
16.92

0.39 ± 0.24

133.96 ±
30.71

13.21 ± 20.28

106.67 ±
27.13

35.85 ± 28.79

13.21 ±
31.82

20 ± 25.43

Gamasid mites

p < 0.05

n.s.

n.s.

p < 0.01

n.s.

p < 0.01

n.s.

n.s.

n.s.

Oribatid mites

203.85 ±
65.9969.23 ±
39.39

0.8 ± 0.39

219.23 ±
71.16142.31 ±
77.57

31.75 ± 29.37

415.38 ±
77.3938.46 ±
53.85272.22 ±
55.89

	p < 0.05	n.s.	n.s.	p < 0.05	p < 0.10	n.s.	p < 0.001	n.s.	p < 0.001
Collembola									
	n.s.	n.s.	n.s.	p < 0.01	p < 0.10	n.s.	p < 0.05	n.s.	p < 0.10
		43.05 ±		109.41 ±				79.26 ±	-33.16 ±
Shannon's diversity	57.54 ± 22.41	28.93	0.1 ± 0.16	20.98	51.9 ± 34.08	37.86 ± 13.81	19.82 ± 27.1	21.71	15.12
	p < 0.05	n.s.	n.s.	p < 0.001	n.s.	p < 0.05	n.s.	p < 0.01	p < 0.10
		15.61 ±							-47.14 ±
Functional richness	24.93 ± 34.62	22.45	0.08 ± 0.3	70.38 ± 27.46	5 ± 32.7	62.26 ± 26.15	-40 ± 27.42	13.5 ± 29.19	24.15
	n.s.	n.s.	n.s.	p < 0.05	n.s.	p < 0.05	n.s.	n.s.	p < 0.10
			-0.04 ±						
Functional evenness	-5.3 ± 4.13	-0.85 ± 4.7	0.04	-8.88 ± 3.46	1.26 ± 3.98	-10.02 ± 3.42	-1.31 ± 5.2	-1.61 ± 4.33	0.31 ± 5.28
	n.s.	n.s.	n.s.	p < 0.05	n.s.	p < 0.05	n.s.	n.s.	n.s.