



Forward-angle electron spectroscopy in heavy-ion atom collisions studied at the ESR

P-M Hillenbrand, S. Hagmann, D Banaś, K-H Blumenhagen, C Brandau, W. Chen, E. de Filippo, A. Gumberidze, D. Guo, D. Jakubassa-Amundsen, et al.

► To cite this version:

P-M Hillenbrand, S. Hagmann, D Banaś, K-H Blumenhagen, C Brandau, et al.. Forward-angle electron spectroscopy in heavy-ion atom collisions studied at the ESR. Journal of Physics: Conference Series, 2015, 635 (2), pp.022005. 10.1088/1742-6596/635/2/022005 . hal-02341306

HAL Id: hal-02341306

<https://normandie-univ.hal.science/hal-02341306>

Submitted on 1 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

PAPER • OPEN ACCESS

Forward-angle electron spectroscopy in heavy-ion atom collisions studied at the ESR

To cite this article: P-M Hillenbrand *et al* 2015 *J. Phys.: Conf. Ser.* **635** 022005

View the [article online](#) for updates and enhancements.

You may also like

- [A study of radiative double electron capture in bare chromium ions at the ESR](#)
N Winters, A Warczak, J A Tanis et al.

- [Developments towards a transverse free-electron target for the storage ring CRYRING@ESR](#)
C. Brandau, A. Borovik, B. M. Döhring et al.

- [Search for High-energy Neutrinos from Binary Neutron Star Merger GW170817 with ANTARES, IceCube, and the Pierre Auger Observatory](#)
A. Albert, M. André, M. Anghinolfi et al.



ECS Membership = Connection

ECS membership connects you to the electrochemical community:

- Facilitate your research and discovery through ECS meetings which convene scientists from around the world;
- Access professional support through your lifetime career;
- Open up mentorship opportunities across the stages of your career;
- Build relationships that nurture partnership, teamwork—and success!

[Join ECS!](#)

[Visit electrochem.org/join](#)



Forward-angle electron spectroscopy in heavy-ion atom collisions studied at the ESR

P.-M. Hillenbrand^{1,2}¹, S. Hagmann^{1,3}², D. Banas⁴, K.-H. Blumenhagen⁵, C. Brandau^{1,2}, W. Chen¹, E. De Filippo⁶, A. Gumberidze¹, D. L. Guo⁷, D. H. Jakubassa-Amundsen⁸, M. Lestinsky¹, Yu. A. Litvinov¹, J. M. Monti⁹, A. Müller², H. Rothard¹⁰, S. Schippers², M. S. Schöffler³, U. Spillmann¹, A. Surzhykov⁵, S. Trotsenko^{1,5}, A. B. Voitkiv¹¹, X. L. Zhu⁸, and Th. Stöhlker^{1,5}

¹GSI Darmstadt, Germany, ²Universität Giessen, Germany, ³Universität Frankfurt, Germany, ⁴JKU Kielce, Poland, ⁵Helmholtz Institut Jena, Germany, ⁶INFN Catalania, Italy, ⁷IMP Lanzhou, China, ⁸LMU München, Germany, ⁹CONICET Rosario, Argentina, ¹⁰CIMAP-CIRIL-Ganil Caen, France, ¹¹Universität Düsseldorf, Germany

Synopsis For the collisions system $\text{U}^{88+} + \text{N}_2$ @ 90 MeV/u the emitted cusp electrons have been studied, and the three processes contributing to the electron cusp were compared to individual theoretical calculations.

In collisions of heavy highly-charged projectile ions with atomic targets, the energy distribution of the emitted electrons is a characteristic observable for the underlying elementary processes. At the experimental storage ring ESR of the heavy-ion accelerator facility GSI, a dedicated magnetic electron spectrometer was built downstream from the gas-jet target, which enabled the measurement of high-energetic electrons emitted in ion-atom collisions within a small cone in the forward direction. Using this electron spectrometer in combination with detectors for emitted x rays and charge-exchanged projectiles, the study of the collision system $\text{U}^{88+} + \text{N}_2$ @ 90 MeV/u revealed three processes resulting in the emission of electrons with a velocity similar to the projectile velocity, i.e., cusp electrons:

(a) The process of **electron loss to continuum** (ELC) corresponds to the ionization of an electron from the projectile into the projectile continuum during the collision with the target,




For the ELC, the measured spectrum was compared to first-order perturbation theory using fully-relativistic Dirac wavefunctions [1].

(b) The process of **electron capture to continuum** (ECC) corresponds to the capture of a target electron into the projectile continuum, while the excess energy is carried away by the recoil of the generated target ion:



For the ECC, the measured spectrum was compared to calculations in the impulse approximation using semi-relativistic Sommerfeld-Maue wavefunctions, and to calculations in the continuum-distorted-wave approach [2].

(c) The process of **radiative electron capture to continuum** (RECC) corresponds to the capture of a target electron into the projectile continuum, while the excess energy is carried away by a photon:



This latter process can be seen as the high-energy endpoint of bremsstrahlung studied in inverse kinematics. For the RECC, the measured spectra were compared to calculations in the impulse approximation using semi-relativistic Sommerfeld-Maue wavefunctions, and to calculations using fully-relativistic Dirac wavefunctions [3].

Within this study it was shown that each of the three processes is characterized by a unique electron cusp shape, being a clear messenger for the underlying charge-exchange mechanism.

References

- [1] P.-M. Hillenbrand *et al.* 2014 *Phys. Rev. A* **90** 042713
- [2] P.-M. Hillenbrand *et al.* 2015 *Phys. Rev. A* **91** 022705
- [3] P.-M. Hillenbrand *et al.* 2014 *Phys. Rev. A* **90** 022707

¹E-mail: P.M.Hillenbrand@gsi.de

²E-mail: S.Hagmann@gsi.de

