



**HAL**  
open science

# $\beta$ -Lactam Exposure Triggers Reactive Oxygen Species Formation in *Enterococcus faecalis* via the Respiratory Chain Component DMK

Loïc Léger, Aurélie Budin-Verneuil, Margherita Cacaci, Abdellah Benachour, Axel Hartke, Nicolas Verneuil

► **To cite this version:**

Loïc Léger, Aurélie Budin-Verneuil, Margherita Cacaci, Abdellah Benachour, Axel Hartke, et al.  $\beta$ -Lactam Exposure Triggers Reactive Oxygen Species Formation in *Enterococcus faecalis* via the Respiratory Chain Component DMK. *Cell Reports*, 2019, 29 (8), pp.2184-2191.e3. 10.1016/j.celrep.2019.10.080 . hal-02371675

**HAL Id: hal-02371675**

<https://hal-normandie-univ.archives-ouvertes.fr/hal-02371675>

Submitted on 21 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution-NonCommercial 4.0 International License

## **$\beta$ -Lactam Exposure Triggers Reactive Oxygen Species**

### **Formation in *Enterococcus faecalis* via the**

### **Respiratory Chain Component DMK**

Loïc Léger,<sup>1</sup> Aurélie Budin-Verneuil,<sup>1</sup> Margherita Cacaci,<sup>2</sup> Abdellah Benachour,<sup>1</sup> Axel

5 Hartke,<sup>1,3\*</sup> and Nicolas Verneuil<sup>1</sup>

<sup>1</sup>UR Risques Microbiens, Normandie Univ, UNICAEN, U2RM, 14000 Caen, France

<sup>2</sup>Università Cattolica del Sacro Cuore, Istituto di Microbiologia, Roma, Italia

<sup>3</sup>Lead Contact

\*Correspondence: [axel.hartke@unicaen.fr](mailto:axel.hartke@unicaen.fr)

## 10 **SUMMARY**

Whereas the primary actions of  $\beta$ -lactams are well characterized, their downstream effects are less well understood. Although their targets are extracellular,  $\beta$ -lactams stimulate respiration in *Escherichia coli* leading to increased intracellular accumulation of reactive oxygen species (ROS). Here we show that  $\beta$ -lactams over a large concentration range trigger a strong increase

15 in ROS production in *Enterococcus faecalis* under aerobic but not anaerobic conditions. Both amoxicillin, of which the bacterium is susceptible, and cefotaxime, of which *E. faecalis* is resistant, triggers this response. This stimulation of ROS formation depends mainly on demethylmenaquinone (DMK), a component of the *E. faecalis* respiratory chain, but in contrast to *E. coli* is observed only in the absence of respiration. Our results suggest that in *E.*

20 *faecalis*,  $\beta$ -lactams increase electron flux through the respiratory chain, thereby stimulating the auto-oxidation of reduced DMK in the absence of respiration, which triggers increased extracellular ROS production.

## INTRODUCTION

Beta-lactam antibiotics are bactericidal drugs that block the activity of extracellular enzymes, known as penicillin-binding proteins (PBPs) which catalyze cross-linking of peptidoglycan peptide side chains thereby inhibiting cell wall synthesis (Tipper and Strominger, 1965; Wise and Park, 1965). It is generally believed that killing by  $\beta$ -lactams is due to their bacteriolytic activity. However, bacteriolysis cannot fully explain the lethality of these antibiotics. For example, penicillin has been shown to kill pathogens like *Streptococcus pyogenes* and *Enterococcus hirae* efficiently without inducing a significant lysis of the cells (Daneo-Moore et al., 1988; Gutmann and Tomasz, 1982; McDowell and Lemanski, 1988). A more recent study concluded that the killing of *Enterococcus faecalis* induced by amoxicillin was due to two distinct mechanisms, one involving autolysins and a second action of the drug independent of lysis (Dubée et al., 2011).

In several works, the secondary actions of  $\beta$ -lactams were analyzed. A recent study showed that  $\beta$ -lactams induce a deleterious futile cycle of cell wall synthesis and degradation thereby depleting cellular resources contributing to their killing activity in *Escherichia coli* (Cho et al. 2014). Other studies showed that  $\beta$ -lactams and other bactericidal antibiotics induce complex toxic metabolic perturbations leading to cell damage (Kohanski et al., 2007; Belenky et al 2015; Dwyer et al., 2009). The authors proposed that these drugs induce an intracellular oxidative stress by increasing respiration and with it the release of reactive oxygen species (ROS) from the respiratory chain overwhelming the cellular antioxidant defenses, leading to DNA damage and finally to cell death (Kohanski et al., 2007; Dwyer et al. 2009). These studies were mainly conducted with *E. coli* but numerous follow up studies on different Gram-negative and Gram-positive bacteria were in accordance with the proposed ROS-based model [see Dwyer et al., (2015) for a recent review]. Concerning  $\beta$ -lactams, major results supporting the induction of oxidative stress by these drugs are: (i) amoxicillin strongly

induced the expression of the *recA* gene encoding the major recombinase involved in DNA repair, caused DNA damage and, in the presence of ferric iron, induced the expression of anti-oxidant activities in *Pseudomonas* (Yeom et al., .2010); (ii) compared to the wild-type parent strain, *Acinetobacter baumannii* and *E. coli recA* mutants were more susceptible to  $\beta$ -lactams (Kohanski et al., 2007; Aranda et al., 2011); (iii) a combination of an iron chelator and a radical scavenger as well as the natural antioxidant glutathione provided protection against oxacillin-mediated killing of *Staphylococcus aureus* (Liu et al., 2012); (iv) reduction of the Fenton reaction by iron chelation by the ferritin-like protein Fri is suggested to be at the basis of  $\beta$ -lactam tolerance of *Listeria monocytogenes* (Krawczyk-Balska et al., 2012). Despite these substantial data, the ROS-based killing model was later challenged by other studies persuasively demonstrating that killing by the bactericidal antibiotics used in the former studies was independent from oxidative stress (Ezraty et al., 2013; Keren et al., 2013; Liu and Imlay, 2013).

These controversial viewpoints reflect that our mechanistic understanding of the action of bactericidal antibiotics on bacteria remains incomplete. In this report, we use the health care-associated pathogen *E. faecalis* as a model bacterium to test the proposal that bactericidal antibiotics induce ROS formation. This gram-positive facultative aerobic/anaerobic bacterium is a usual member of the intestinal microbiome of humans and has interesting properties to address this important question of downstream actions of antibiotics. Compared to other gram-positive cocci, enterococci exhibit decreased susceptibility to penicillins and high-level resistance to cephalosporins. We showed previously that this  $\beta$ -lactam tolerance as well as tolerance to vancomycin is linked to the superoxide dismutase SodA in *E. faecalis* and *E. faecium* (Bizzini et al., 2009; Ladjouzi et al., 2013). Unlike *E. coli*, enterococci have no tricarboxylic acid cycle, a pathway central in the Kohanski et al. (2007) model. Furthermore, respiration, responsible for ROS formation in this model, is facultative and only functional in

the presence of haematin or fumarate, although respiration by extracellular electron transfer in iron-enriched biofilms has been reported recently for this species (Keogh et al., 2018). *E. faecalis* can produce extracellular superoxide ( $O_2^-$ ) through autoxidation of membrane-associated demethylmenaquinone (DMK) (Huycke et al., 2001). It has been demonstrated that  
5 this radical is also produced in the mammalian intestinal tract (Huycke and Moore, 2002; Moore et al., 2004) and responsible for chromosomal instability (Wang and Huycke, 2007; Wang et al., 2008), inflammation and colorectal cancer (CRC) (Balish and Warner, 2002; Kim et al., 2005). The objective of the present work was to analyze if  $\beta$ -lactam antibiotics increase ROS formation in *E. faecalis* wild-type cells using drug concentrations potentially  
10 encountered by these pathogens during antimicrobial treatment. We show that this is indeed the case and provide a mechanistic explanation for this secondary action of these cell-wall targeting drugs.

## RESULTS

### Superoxide dismutase-deficient mutant is sensitive to amoxicillin

Amoxicillin is still one of the treatments of choice for enterococcal infections that lack mechanisms for high-level resistance (Mercuro et al., 2018). Therefore, we first analyzed whether this bactericidal drug induces an oxidative stress in *E. faecalis*. Since enterococci exhibit a high resistance to oxidants (Flahaut et al., 1998), we reasoned that the most direct way to explore the proposed role of ROS in amoxicillin mortality would be to use mutants defective either in  $O_2^-$  or in peroxide defenses as indicators of oxidative stress. *E. faecalis* harbors a manganese-cofactored superoxide dismutase (MnSod or SodA) and three peroxidases, the NADH peroxidase (Npr), the alkyl hydroperoxide reductase (AhpCF) and a thiol peroxidase (Tpx). In this work, the  $\Delta sodA$  and  $\Delta npr\Delta tpx\Delta ahpCF$  mutants known to be hypersensitive to ROS were used (La Carbona et al., 2007; Verneuil et al., 2006). Since under our experimental conditions the minimal inhibition concentration (MIC) of amoxicillin was 0.5  $\mu\text{g}/\text{mL}$  for all strains, we tested two different concentrations (0.25 and 0.5  $\mu\text{g}/\text{mL}$ ). As shown in Figure 1A, in the absence of amoxicillin, growth of the triple peroxidase mutant was comparable to that of the parental strain whereas survival of the  $\Delta sodA$  mutant decreased slightly between 16 hours and 24 hours of incubation under the used aeration condition. This is likely due to the generation and accumulation of ROS since no decrease in survival of the  $\Delta sodA$  mutant was observed with cultures grown in anaerobic conditions (Figure S1A). Rapid killing of all strains was observed with 0.5  $\mu\text{g}/\text{mL}$  of amoxicillin (Figure 1B). No significant differences in survival between the wild-type and the  $\Delta sodA$  mutant strain were observed under this condition whereas the triple peroxidase mutant was somewhat more resistant to killing by the antibiotic than the two former strains (Figure 1B). Rapid killing also occurred under anaerobic conditions at this amoxicillin concentration (Figure S1B). We concluded that at 0.5  $\mu\text{g}/\text{mL}$  of the antibiotic, lethality of the bacteria was independent of an oxidative stress.

At 0.25  $\mu\text{g}/\text{mL}$  of amoxicillin (half MIC), the survival of all strains was comparable during the first 16 hours of incubation (Fig. 1C). During this period, killing of the strains was minimal and increased slightly in the case of the wild-type and triple peroxidase mutant after 24 hours of exposition to the antibiotic. In contrast, the killing of the  $\Delta\text{sodA}$  mutant decreased by  $\sim 5.7 \log_{10}$  between 16 and 24 hours (Figure 1C). This impressive mortality of the  $\Delta\text{sodA}$  mutant was not observed under anaerobic conditions (Figure S1C). These results are consistent with a ROS-based killing mode of the MnSod deficient *E. faecalis* mutant at half MIC amoxicillin. Decrease of survival starts after 16 hours of aerobic amoxicillin treatment suggesting that ROS accumulates until creating some sort of damage that became lethal for the  $\Delta\text{sodA}$  mutant.

### **Amoxicillin promotes $\text{H}_2\text{O}_2$ generation**

$\text{O}_2^-$  is readily dismutated into  $\text{H}_2\text{O}_2$  ( $2 \text{O}_2^- \leftrightarrow \text{H}_2\text{O}_2 + \text{O}_2$ ) (Bielski and Richter, 1977) and therefore we wondered if catalase which catalyses the dismutation of  $\text{H}_2\text{O}_2$  to molecular oxygen and water would decrease killing by amoxicillin. As shown in Figure 2A, the survival of the  $\Delta\text{sodA}$  mutant was nearly restored to the wild-type level in the presence of active catalase (but not with heat-inactivated catalase) strongly suggesting that besides  $\text{O}_2^-$ ,  $\text{H}_2\text{O}_2$  is also generated during amoxicillin treatment and causatively contributes to antibiotic-mediated killing of the  $\Delta\text{sodA}$  mutant. In the next step, we verified this hypothesis by directly quantifying  $\text{H}_2\text{O}_2$  accumulation by *E. faecalis* cultivated with and without amoxicillin. In the absence of the antibiotic, the  $\Delta\text{sodA}$  mutant generated  $\text{H}_2\text{O}_2$  reaching a maximum concentration of around 700  $\mu\text{M}$  after 16 hours of incubation and was slightly lower at 24 hours (Figure 2B). No  $\text{H}_2\text{O}_2$  was detected in cultures treated with 0.5  $\mu\text{g}/\text{mL}$  of amoxicillin, which is likely due to the rapid ROS-independent killing of the cells at this concentration. However, in the presence of half MIC amoxicillin, cells produced  $\text{H}_2\text{O}_2$  continuously over the whole incubation time to reach  $>2000 \mu\text{M}$  after 24 hours; that is around 4-times more than in

the absence of the drug (Figure 2B). We wondered then if amoxicillin treatments also induce an increased H<sub>2</sub>O<sub>2</sub> production in wild-type cells which are not killed by 0.25 µg/mL of amoxicillin. Relative to the  $\Delta sodA$  mutant, the H<sub>2</sub>O<sub>2</sub> concentrations measured were globally lower but we found that even in the parental strain the amoxicillin treatment triggered an increased continuous accumulation of H<sub>2</sub>O<sub>2</sub> over the 24 hours incubation period (Figure 2C). At the end, the H<sub>2</sub>O<sub>2</sub> concentration accumulated was >600 µM in the presence of 0.25 µg/mL of the drug which correspond to 8.4-fold more of this ROS than in the absence of the antibiotic (Figure 2C). We then extended these analyses by testing H<sub>2</sub>O<sub>2</sub> production in the presence of a wider range of amoxicillin concentrations. As expected, no H<sub>2</sub>O<sub>2</sub> accumulation was observed with lethal concentrations ( $\geq 0.5$  µg/mL) of the drug (Figure 2D). However, concentrations of amoxicillin below 0.25 µg/mL seems to trigger more accumulation of H<sub>2</sub>O<sub>2</sub> although the differences were not statistically significant when compared to 0.25 µg/mL (Figure 2D). Of note, H<sub>2</sub>O<sub>2</sub> levels accumulated by the  $\Delta npr\Delta tpx\Delta ahpCF$  strain were of the same order of magnitude as those determined with the  $\Delta sodA$  mutant (Figure 2B). We concluded from the combined results that amoxicillin stimulates H<sub>2</sub>O<sub>2</sub> formation in *E. faecalis*.

### **A *recA* mutant is as resistant as the wild-type strain to amoxicillin**

We wondered if the amoxicillin treatment would also increase intracellular ROS production. In *E. coli*, ampicillin induces the SOS response and DNA damage, which has been attributed to the increased formation of hydroxyl radicals by the drug (Kohanski et al. 2007; Miller et al. 2004). *E. coli* and *A. baumannii* strains deficient in RecA, a protein promoting the central steps in recombination and recombinational DNA repair, are more sensitive to ampicillin, supporting the increased intracellular ROS production and DNA damage in cells exposed to this antibiotic (Aranda et al., 2011; Kohanski et al., 2007). We used a well defined  $\Delta recA$  mutant (Boumghar-Bourtchaj et al., 2009) to analyse if this scenario also hold true in



*E. faecalis* cells treated with amoxicillin. This mutant was highly more susceptible to UV radiation (Figure S2A) but was as resistant as wild-type strain to a treatment with 0.25 µg/mL of amoxicillin (Figure S2B). We concluded that intracellular concentration of oxidants should be low since no RecA-dependent DNA damages are caused by the drug in *E. faecalis*.

## 5 **Amoxicillin-induced H<sub>2</sub>O<sub>2</sub> accumulation appears to depend mainly on DMK**

Next, we addressed the question about the molecular mechanism(s) of drug-induced ROS formation. The cell death pathway model described in *E. coli* proposed the involvement of the respiratory chain as the primary source of increased intracellular ROS formation by antibiotics (Kohanski et al., 2007). *E. faecalis* is capable of aerobic respiration and the presence of cytochromes and DMK has been identified in its membrane (Huycke et al., 2001). However, since enterococci lack the ability to synthesize heme, their cytochromes are only functional in the presence of a source for this prosthetic group (Pritchard and Wimpenny, 1978; Ritchey and Seeley, 1974). In the absence of heme, aerobic respiration is blocked, and it has been demonstrated that this leads to an increase in extracellular O<sub>2</sub><sup>-</sup> generation through adventitious autooxidation of membrane-associated reduced DMK (DMKH<sub>2</sub>) (Huycke et al., 2001). Since a known H<sub>2</sub>O<sub>2</sub>-producing pathway in *E. faecalis* is the spontaneous disproportionation of extracellular O<sub>2</sub><sup>-</sup> (Huycke et al., 2002), we wondered whether the amoxicillin-induced ROS formation could be a result of the stimulation of autooxidation of DMK. To test this possibility, we measured survival of a  $\Delta sodA$  *menB::tet* double mutant in the presence of amoxicillin. The *menB* gene encodes 1,4-dihydroxy-2-naphthoic acid synthase catalyzing an essential step in the DMK biosynthesis pathway (Huycke et al., 2001). Relative to the  $\Delta sodA$  single mutant, survival of the  $\Delta sodA$  *menB::tet* double mutant increased by >4 log<sub>10</sub> (Figure 3A), likely due to reduced ROS production. This was confirmed by measuring H<sub>2</sub>O<sub>2</sub> concentrations in the culture medium of the *menB::tet* single and the  $\Delta sodA$  *menB::tet* double mutants. H<sub>2</sub>O<sub>2</sub> concentration in the former strain was under the detection limit and the

latter strain accumulated 6-times and 3-times less H<sub>2</sub>O<sub>2</sub> than the  $\Delta sodA$  single mutant at 16 hours and 24 hours, respectively (Figure 3B). This strongly suggested DMK as the major source of H<sub>2</sub>O<sub>2</sub> accumulation.

If right, depletion of electrons from the DMKH<sub>2</sub> pool should also decrease killing by amoxicillin. This was tested by addition of haematin which induces synthesis of cytochrome *bd* oxidase allowing aerobic respiration as well as by addition of fumarate which allows anaerobic respiration via a fumarate reductase (Huycke et al., 2001). In the presence of haematin, killing of the  $\Delta sodA$  mutant (Figure 3C) and accumulation of H<sub>2</sub>O<sub>2</sub> were significantly reduced (Figure 3D). However, it was recently reported that in addition to its respiratory role, cytochrome *bd* from *E. coli* displays catalase activity (Borisov et al., 2013). If true for the cytochrome *bd* of *E. faecalis*, it may reduce amoxicillin-induced H<sub>2</sub>O<sub>2</sub> accumulation. Moreover, *E. faecalis* also harbors a gene encoding a heme-dependent catalase (Frankenberg et al., 2002) and it could not be excluded that it may also contribute to H<sub>2</sub>O<sub>2</sub> degradation in the presence of haematin. Therefore, we performed experiments in presence of amoxicillin and fumarate. *E. faecalis* expresses fumarate reductase, a membrane-associated enzyme which catalyzes the reduction of fumarate to succinate. Electrons are provided by DMKH<sub>2</sub> and this process allows non-oxidative respiration in the absence of haematin (Huycke, 2002). The addition of fumarate improved the survival (Figure 3E) and reduced H<sub>2</sub>O<sub>2</sub> accumulation (Figure 3F) of the  $\Delta sodA$  mutant. Collectively, these results strongly suggest that amoxicillin-induced H<sub>2</sub>O<sub>2</sub> accumulation depends mainly on autooxidation reactions involving DMK.

### **Amoxicillin appears to cause an increased electron flow through DMK**

Stimulation of the autooxidation reactions involving DMK by amoxicillin suggests an increase in electron flow through the respiratory chain of *E. faecalis*. We used fumarate

respiration to verify this hypothesis. As *E. faecalis* does not express succinate dehydrogenase activity (Aue and Deiel, 1967), an increase in succinate formation in the presence of exogenous fumarate and amoxicillin would reflect an increased electron flux through the respiratory chain. We therefore analyzed succinate levels in the supernatants of wild-type  
5 cultures challenged with the antibiotic and found that, relative to untreated control, amoxicillin induced a statistically significant 2-fold increase in succinate production rate (Figure 4A), supporting the hypothesis that amoxicillin causes an increase in electron flow through DMK and fumarate reductase.

In the absence of fumarate and haematin, an increased electron flux through DMK  
10 cycle would similarly elevate the rate of  $O_2^-$  formation, by increasing autooxidation reactions. Thus, we evaluated the extracellular  $O_2^-$  production rate of amoxicillin-treated and untreated cells and found that the antibiotic induced a >2-fold increase in the formation of  $O_2^-$  per minutes for  $10^9$  CFU (Figure 4B). However, the difference between untreated and treated cells seems not statistically significant. Furthermore, it could not be excluded that part of the  
15 cells in the culture have been damaged by the antibiotic treatment and consequently are not able to form colonies. These cells might be still metabolically active and produce  $O_2^-$  which might overestimate the production rate of this radical. Nevertheless, as we observed a similar statistically significant increase in succinate production rate (Figure 4A), we concluded that these results support the hypothesis that amoxicillin causes an increased electron flow through  
20 DMK which in absence of fumarate and haematin, promotes the formation of  $O_2^-$ .

### **Cefotaxime also increases ROS production in *E. faecalis***

*E. faecalis* is intrinsically highly resistant to cephalosporins. Therefore, we wondered whether this  $\beta$ -lactam would also stimulate the generation of ROS. As shown in Figure S3, this is indeed the case. The  $\Delta sodA$  mutant was  $\sim 6.1 \log_{10}$  more susceptible than the parental strain to

treatment with 1  $\mu\text{g/mL}$  cefotaxime, a third-generation cephalosporin (Figure S3A). The antibiotic promoted  $\text{H}_2\text{O}_2$  production in both strains (Figures S3B and S3C) and this stimulation was observed over a wide concentration range of the antibiotic in the wild-type strain (Figure S3D). Cefotaxime treatment also increased, relative to untreated cells, both  
5 extracellular  $\text{O}_2^-$  production by ~2-fold (Figure S3E) as well as fumarate respiration (Figure S3F).

## DISCUSSION

The present study contributes to an ongoing discussion concerning the secondary mode of actions of antibiotics (Liu and Imlay, 2013). A better understanding of these less well understood downstream effects might pave the way for the development of new treatment strategies (Brynildsen et al., 2013). Here, we clearly demonstrate that  $\beta$ -lactams increase the production of ROS in *E. faecalis*, whether in the wild-type strain or the  $\Delta sodA$  and  $\Delta npr\Delta tpx\Delta ahpCF$  mutants. Unexpectedly and unrelated to antibiotic treatment, the  $\Delta sodA$  mutant, for hitherto unknown reasons, accumulated more  $H_2O_2$  than the wild-type strain (Figures 2B and 2C). Since the mutant deficient of all peroxidases of *E. faecalis* accumulated comparable high concentrations of  $H_2O_2$  (Figure 2B), it might be that peroxidases are inactive in the  $\Delta sodA$  background.

In *E. coli*, the respiratory chain is suspected to be at the basis of ROS generation under antibiotic treatment (Kohanski et al., 2007). Our results show that the ROS production appears to depend mainly on DMK, which is consistent with this proposal. However, in contrast to the Kohanski et al. (2007) study, significant ROS production is only observed in our study if electrons are stalled in the DMK pool, which is the case in environments lacking haematin or fumarate. In the presence of  $\beta$ -lactams, the pool of  $DMKH_2$  might increase which consequently will stimulate  $O_2^-$  formation. In the presence of haematin or fumarate, electrons will not accumulate in the DMK pool strongly attenuating the adventitious reaction of molecular oxygen with  $DMKH_2$ . However, further experimentations are now needed to decipher the detailed molecular changes induced by  $\beta$ -lactam antibiotics leading to the increased generation of ROS in *E. faecalis*.

It is worth mentioning that in a previous study we analyzed tolerance of *E. faecalis* to high antibiotic concentration (20xMIC) (Bizzini et al. 2009). We focused in this former study

mainly on vancomycin, a glycopeptide inhibiting cell wall synthesis by targeting the pentapeptide precursors. Whereas the wild-type strain was tolerant to 20xMIC of this drug, survival of the  $\Delta sodA$  mutant decreased by 4log<sub>10</sub>. In contrast to the present study, this decrease in survival was independent of DMK since the  $\Delta sodA menB::tet$  double mutant was  
5 as sensitive to vancomycin as the  $\Delta sodA$  single mutant. This difference to the present work is likely due to the different antibiotics and/or concentrations used in the two studies.

Albesa et al. (2004) showed that *S. aureus*, *E. coli*, and *E. faecalis* sensitive to ciprofloxacin exhibited oxidative stress when they were incubated with the antibiotic, whereas resistant strains did not. Surprisingly, our results demonstrate that cefotaxime also increase  
10 ROS generation in *E. faecalis* although these bacteria are highly resistant to the drug (MIC  $\geq 256$   $\mu\text{g/mL}$ ). Cephalosporin resistance of enterococci is not fully understood but one well characterized component is a specialized low-affinity PBP (Pbp5) that remains active in the presence of these drugs. Pbp5 allows therefore to perform cross-linking of peptidoglycan and growth of the bacteria in the presence of the antibiotic (Arbeloa et al., 2004; Signoretto et al.,  
15 1994). Another key component necessary for intrinsic cephalosporin resistance in *E. faecalis* is the CroR-CroS two-component signal transduction system (Djorić and Kristich, 2015). The loss of intrinsic resistance to cephalosporins of CroRS mutants remains obscure but it has been reported that oxidative stress enhances intrinsic cephalosporin resistance in *E. faecalis* and that in a CroR-CroS-dependent manner (Djorić and Kristich, 2015). The authors showed  
20 that mutants accumulating more H<sub>2</sub>O<sub>2</sub> than the parental strain or wild-type cultures incubated with extracellular added H<sub>2</sub>O<sub>2</sub> demonstrated increased resistance to ceftriaxone. This increased resistance is specific for cephalosporins since no increase in MICs has been observed with other cell wall targeting drugs including the  $\beta$ -lactam ampicillin. In view of these results it might be suggested that the increased synthesis of O<sub>2</sub><sup>-</sup> and H<sub>2</sub>O<sub>2</sub> during  $\beta$ -

lactam treatment evidenced in our study corresponds to an active adaptive stress response with the aim to better survive the antibiotic challenge.

*E. faecalis* is suspected to play a role in colorectal carcinogenesis. This pro-carcinogenic potential seems to be linked with its capacity to generate  $O_2^-$  by the autooxidation of DMK since tumor formation was observed with wild-type bacteria but not with *menB* mutants (Wang et al., 2012). The results presented in this communication might therefore reveal an important concern on the careful use of  $\beta$ -lactam antibiotics since amoxicillin increases significantly the generation of ROS in *E. faecalis* which could increase the risk of CRC development. Interestingly, several recent nested case-control studies analyzing large population-based databases demonstrated indeed a higher risk for CRC with increased number of antibiotic courses and exposure intensities, especially with the most frequently prescribed penicillins (Cao et al., 2018; Dik et al., 2016). Our results might indicate that *E. faecalis* could be a prime candidate to provide a mechanistic explanation for the increased incidence of CRC after therapy with  $\beta$ -lactams. This risk might be even amplified in patients treated with broad-spectrum cephalosporins, since enterococci exhibit intrinsic resistance to these antibiotics, enabling them to proliferate and achieve abnormally high densities in the gastrointestinal tract under these conditions (Donskey et al., 2000).

In conclusion, our study demonstrates that  $\beta$ -lactam antibiotics, beside of inhibiting their extracellular targets, have clearly off-target intracellular effects in *E. faecalis*. They seem to modify metabolism leading to an increased production of ROS mainly *via* DMK autooxidation. On the long run, this could be harmful to patients frequently treated with these drugs. Mechanisms by which  $\beta$ -lactams elicit these metabolic alterations are still poorly understood on the molecular level and their understanding might allow the development of new drugs devoid of these undesired downstream effects. Next steps will be to identify the

signaling cascade activated by the drugs action outside the cell and the rearrangements of intracellular metabolism triggered by these signals.



## **ACKNOWLEDGMENTS**

We are especially grateful to J.A. Imlay (University of Illinois, USA) and B. Ezraty (Université Aix-Marseille, France) for critically reading of the manuscript and helpful comments on this topic. Also, we thank J.C. Giard (Université Caen Normandie, France) for the generous gift of antibiotics. The expert technical assistance of I. Rincé and A.C. Appourchaux (Université Caen Normandie, France) was greatly appreciated. This work as well as the doctoral fellowship of Loïc Léger was funded by the French Ministry of Education and Science.

## **AUTHOR CONTRIBUTIONS**

10 Conceptualization: L.L., A.B.V., A.B., A.H., and N.V.; Methodology: L.L., A.B.V., and M.C.; Validation: L.L., A.B.V., A.B., A.H., and N.V.; Formal Analysis: L.L. and A.B.V.; Investigation: L.L., A.B.V., and M.C.; Resources: L.L., A.B.V., A.B., A.H., and N.V.; Writing – Original Draft Preparation: L.L., A.B.V., A.B., A.H., and N.V.; Writing – Review & Editing: L.L., A.B.V., A.B., A.H., and N.V.; Visualization: L.L., A.B.V., A.B., A.H., and  
15 N.V.; Supervision: A.B., A.H., and N.V.; Project Administration: A.B.V. and N.V.; Funding Acquisition: A.B., A.H., and N.V.

## **DECLARATION OF INTERESTS**

The authors declare no competing interests.

## MAIN FIGURE TITLES AND LEGENDS

### Figure 1. The $\Delta sodA$ mutant is sensitive to amoxicillin at 0.25 $\mu\text{g}/\text{mL}$

(A–C) Survival of *E. faecalis* JH2-2 wild-type strain (squares),  $\Delta sodA$  (diamonds) and  $\Delta npr\Delta tpx\Delta ahpCF$  (triangles) mutants growing under aerobic conditions in the absence of  
5 antibiotic (A), in the presence of 0.5  $\mu\text{g}/\text{mL}$  (B) or 0.25  $\mu\text{g}/\text{mL}$  amoxicillin (C).

Data are represented as means of 2 to 5 experiments, and error bars depict standard deviations. Asterisks indicate a statistically significant difference between mutants and the wild-type strain at 24 hours. \* $p < 0.05$ , \*\*\* $p < 0.001$  (Student's t test). ncd: no colonies detected.

10 See also Figures S1 and S3.

### Figure 2. Amoxicillin promotes $\text{H}_2\text{O}_2$ generation

(A) Survival of the  $\Delta sodA$  mutant exposed to 0.25  $\mu\text{g}/\text{mL}$  amoxicillin under aerobic conditions in the presence of ~500 U/mL bovine catalase (Cat) or heat-inactivated catalase (H-I Cat). For comparisons, survival curve of the wild-type strain exposed to 0.25  $\mu\text{g}/\text{mL}$   
15 amoxicillin under aerobic conditions is also shown.

(B and C)  $\text{H}_2\text{O}_2$  concentration of supernatants of  $\Delta sodA$  and  $\Delta npr\Delta tpx\Delta ahpCF$  mutants (B) and wild-type strain (C) cultures grown in the absence of antibiotic (open bars) or after treatment with 0.25  $\mu\text{g}/\text{mL}$  amoxicillin (closed bars) under aerobic conditions.

(D)  $\text{H}_2\text{O}_2$  in culture supernatants of wild-type strain cultivated under various concentrations of  
20 amoxicillin for 24 hours under aerobic conditions.

Data are represented as means of 2 to 4 experiments, and error bars depict standard deviations. For  $\text{H}_2\text{O}_2$  assays, asterisks indicate a statistically significant difference between

treated and untreated cells. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  (Student's t test). ncd: no colonies detected.

See also Figure S3.

**Figure 3. Amoxicillin-induced H<sub>2</sub>O<sub>2</sub> accumulation appears to depend mainly on DMK**

5 (A–F) Cultures are realized with amoxicillin at 0.25  $\mu\text{g/mL}$  under aerobic conditions.

(A) Survival of  $\Delta\text{sodA}$  (diamonds) and  $\Delta\text{sodA menB}::\text{tet}$  (circles) mutants.

(B) H<sub>2</sub>O<sub>2</sub> concentration of supernatants of  $\Delta\text{sodA}$  (open squares) and  $\Delta\text{sodA menB}::\text{tet}$  (closed squares) cultures.

(C) Survival of the  $\Delta\text{sodA}$  mutant in the presence of 8  $\mu\text{M}$  haematin/0.56  $\mu\text{M}$  NaOH (closed  
10 diamonds) or 0.56  $\mu\text{M}$  NaOH (open diamonds).

(D) H<sub>2</sub>O<sub>2</sub> concentration of supernatants of  $\Delta\text{sodA}$  mutant cultures supplemented with 8  $\mu\text{M}$  haematin/0.56  $\mu\text{M}$  NaOH (closed bars) or 0.56  $\mu\text{M}$  NaOH (open bars).

(E) Survival of the  $\Delta\text{sodA}$  mutant in the presence (closed diamonds) or absence (open diamonds) of 8 mM fumarate.

15 (F) H<sub>2</sub>O<sub>2</sub> concentration of supernatants of  $\Delta\text{sodA}$  mutant cultures supplemented with (closed bars) or without (open bars) 8 mM fumarate.

Data are represented as means of 2 to 4 experiments, and error bars depict standard deviations. For H<sub>2</sub>O<sub>2</sub> assays, asterisks indicate a statistically significant difference between 2 data sets. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  (Student's t test). ncd: no colonies detected.

20 See also Figure S2.

**Figure 4. Amoxicillin appears to cause an increased electron flow through DMK**

(A) Succinate production in the wild-type strain after 4 hours of growth in the presence of 20 mM fumarate in the absence of antibiotic (open bars) or after treatment with 0.25  $\mu\text{g/mL}$  amoxicillin (closed bars) under aerobic conditions.

5 (B) Extracellular  $\text{O}_2^-$  production of wild-type strain after 24 hours of growth in the absence of antibiotic (open bar) or after treatment with 0.25  $\mu\text{g/mL}$  amoxicillin (closed bar) under aerobic conditions.

Data are represented as means of 2 to 4 experiments, and error bars depict standard deviations. Asterisks indicate a statistically significant difference between mutants and the wild-type strain at 24 hours. \* $p < 0.05$  (Student's t test).

10 See also Figure S3.

## STAR★METHODS

### LEAD CONTACT AND MATERIALS AVAILABILITY

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Aurélie Budin-Verneuil (aurelie.verneuil@unicaen.fr).

- 5 All unique/stable reagents generated in this study are available from the Lead Contact without restriction.

### EXPERIMENTAL MODEL AND SUBJECT DETAILS

#### Bacterial Strains

The bacterial strains used in this study are listed in Table S1. Strains of *E. faecalis* were  
10 grown in M17 medium (Terzaghi and Sandine, 1975) supplemented with 0.3% glucose (GM17) at 37°C (Table S2). *E. coli* VE14188 was cultivated at 37°C in Luria-Bertani (LB) broth (Miller, 1972) with agitation or on LB agar. When appropriate, tetracycline (10 µg/mL for *E. faecalis*, 12.5 µg/mL for *E. coli*) was added to the medium.

### METHOD DETAILS

#### 15 Survival

For overnight cultures, a single colony was grown at 37°C in 10 mL of GM17, with tetracycline if necessary, for about 16 hours. For survival in aerobic conditions, the culture was diluted in 100 mL of GM17 in a 1000 mL Erlenmeyer flask to obtain a final OD<sub>600nm</sub> of 0.01. The bacterial suspension was incubated at 37°C with moderate shaking (60 rpm) for  
20 about 3 hours until reaching OD<sub>600nm</sub> of 0.5 (exponential growth phase). Cells were then collected by centrifugation (10 minutes at 3,000 g) and resuspended in fresh GM17 prewarmed at 37°C. Eight milliliters of culture were then distributed into 100 mL Erlenmeyer flasks and 2 mL of antibiotics [diluted in GM17 to obtain the defined final concentration of

amoxicillin or cefotaxime (Panpharma, France), respectively] were added. In the case of the controls without antibiotic, 2 mL of GM17 were added to these flasks. Samples were collected immediately ( $T_0$ ) and after incubation for 8 h, 16 h and 24 h at 37°C with shaking at 60 rpm. All samples were then serially diluted in 0.9% saline, spread onto GM17 plates, and  
5 incubated at 37°C for about 48 hours. Survival was determined by counting colonies on plates harboring between 30 and 300 colonies.

For survival in anaerobic conditions, an overnight culture (~16 hours) was diluted in 100 mL of GM17 in a 100 mL sealed bottle to obtain a final  $OD_{600nm}$  of 0.01 and incubated at 37°C without agitation for about 3 hours until  $OD_{600nm}$  of 0.5. The survival experiments were  
10 carried out as previously described except that it was performed in 10 mL sealed tubes containing 10 mL of the above culture and incubated without agitation.

When used, catalase, fumarate, or haematin (Sigma-Aldrich, MO) were diluted to a final concentration of 500 U/mL, 8 mM and 8  $\mu$ M, respectively. Haematin was prepared as a 20 mM stock solution in 1.4 mM NaOH. Catalase was inactivated by boiling for 5 min.

## 15 **H<sub>2</sub>O<sub>2</sub> Assay**

The concentrations of H<sub>2</sub>O<sub>2</sub> were measured in the supernatants using Amplex® Red Hydrogen Peroxide/Peroxidase Assay Kit (Invitrogen, OR) as described previously (Bizzini et al., 2009). Briefly, supernatants of cultures treated with antibiotics as described above were collected at the indicated time points by centrifugation (10 min at 3,000 g) and diluted 100-  
20 fold in 50 mM sodium phosphate buffer, pH 7.4. The H<sub>2</sub>O<sub>2</sub> concentrations were determined using 50  $\mu$ L of these dilutions and 50  $\mu$ L of a working solution of 100  $\mu$ M Amplex® Red reagent and 0.2 U/mL HRP. In parallel, H<sub>2</sub>O<sub>2</sub> standard curves (0 to 5  $\mu$ M) were prepared using GM17 diluted 100-fold in 1X Reaction Buffer. Reactions were incubated at room temperature for 30 minutes. Fluorescence was then measured with a FlexStation 3 reader

(Molecular Devices, CA) using excitation at 530 nm and fluorescence detection at 590 nm. Background fluorescence, determined from no-H<sub>2</sub>O<sub>2</sub> control reactions, has been subtracted for each value.

### **Construction of the *menB::tet* Mutant**

5 To construct the *menB::tet* mutant, we used the recombinant pVE14218 (a pORI derivative plasmid) harboring an internal fragment of *menB* gene (*menB::pVE14218*) previously used to construct the  $\Delta$ *sodA menB::tet* double mutant (Bizzini et al., 2009). The *menB::tet* mutant was obtained following a single crossing over insertion mutagenesis based on a two-vector system using the plasmids pGhost3 (Maguin et al., 1992), which provides functional RepA<sup>Ts</sup>,  
10 and the conditionally replicating pVE14218 (Tet<sup>R</sup>) (Rigottier-Gois et al., 2011).

### **HPLC Analysis of Succinate Production**

Succinate concentrations were determined by HPLC (Waters 600 Controller). As described in “Survival” section, cells were treated with antibiotics in presence of fumarate (20 mM) for 4 hours and collected by centrifugation. The filtrates of supernatants as well as internal  
15 standards were separated on an Aminex HPX-87H column (Biorad) maintained at 65°C and eluted for 30 min using a 5 mM H<sub>2</sub>SO<sub>4</sub> mobile phase with a flow rate of 0.6  $\mu$ l/min. Peak areas of substrates and derivative products were determined with the Borwin chromatography software (JMBS Developments).

### **Extracellular O<sub>2</sub><sup>-</sup> Assay**

20 Cells treated with antibiotics as described above were collected after 24 hours by centrifugation (10 min at 3,000 g), washed twice with Washing Buffer (7.5 mM ammonium sulfate, 6 mM sodium chloride, 33 mM potassium dihydrogenphosphate, 60 mM potassium hydrogenphosphate, and 1 mM magnesium chloride, pH 7.3), and resuspended in Washing Buffer supplemented with 0.33% glucose to an approximate OD<sub>550nm</sub> of 0.2. The extracellular

O<sub>2</sub><sup>-</sup> production was measured as described by Huycke et al. (1996) with slight modifications. Briefly, after being warmed to room temperature, ferricytochrome *c* was added to cell suspension in 1 cm-cuvettes to obtain a final concentration of 20 μM, and the reduction of ferricytochrome *c* was monitored over 10 min at 550 nm in the presence or absence of SOD (25 μg/mL; Sigma-Aldrich, MO) in a SmartSpecTMPlus spectrophotometer (Biorad, CA). The rate of ferricytochrome *c* reduction was calculated using linear regression. The rate of reduction in the absence of SOD was corrected from the rate measured in its absence. O<sub>2</sub><sup>-</sup> production was calculated as described by Huycke et al. (1996), using an extinction coefficient of 21.5 mM<sup>-1</sup> cm<sup>-1</sup> for reduced cytochrome *c*. In contrast to Huycke et al. (1996), viable counts were determined to normalize the rate of O<sub>2</sub><sup>-</sup> production (nmol/min) to 10<sup>9</sup> CFU.

## QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical parameters and significances are reported in the figure legends and figures. Means are considered significantly different when  $p < 0.05$ . In figures, asterisks denote statistical significances as calculated by Student's *t* test (\*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ ). Statistical analyses were performed in GraphPad Prism 5. No statistical calculations could be done for 24 hour survivals lacking CFU counts (refer to Figure 1, Figure 2, Figure 3 and Figure S3). These results are highlighted in the corresponding figures by "ncd" meaning "no colonies detected".

## DATA AND CODE AVAILABILITY

The published article includes all datasets generated or analyzed during this study.



## REFERENCES




- Albesa, I., Becerra, M.C., Battán, P.C., and Páez, P.L. (2004). Oxidative stress involved in the antibacterial action of different antibiotics. *Biochem. Biophys. Res. Commun.* *317*, 605–609.
- Aranda, J., Bardina, C., Beceiro, A., Rumbo, S., Cabral, M.P., Barbé, J., and Bou, G. (2011). Acinetobacter baumannii RecA protein in repair of DNA damage, antimicrobial resistance, general stress response, and virulence. *J. Bacteriol.* *193*, 3740–3747.
- Arbeloa, A., Segal, H., Hugonnet, J.E., Josseaume, N., Dubost, L., Brouard, J.P., Gutmann, L., Mengin-Lecreulx, D., and Arthur, M. (2004). Role of class A penicillin-binding proteins in PBP5-mediated beta-lactam resistance in Enterococcus faecalis. *J. Bacteriol.* *186*, 1221–1228.
- Aue, B.J., and Deiel, R.H. (1967). Fumarate reductase activity of Streptococcus faecalis. *J. Bacteriol.* *93*, 1770–1776.
- Balish, E., and Warner, T. (2002). Enterococcus faecalis induces inflammatory bowel disease in interleukin-10 knockout mice. *Am. J. Pathol.* *160*, 2253–2257.
- Belenky, P., Ye, J.D., Porter, C.B.M., Cohen, N.R., Lobritz, M.A., Ferrante, T., Jain, S., Korry, B.J., Schwarz, E.G., Walker, G.C., et al. (2015). Bactericidal Antibiotics Induce Toxic Metabolic Perturbations that Lead to Cellular Damage. *Cell Rep.* *13*, 968–980.
- Bielski, B.H.J., and Richter, H.W. (1977). A study of the superoxide radical chemistry by stopped-flow radiolysis and radiation induced oxygen consumption. *J. Am. Chem. Soc.* *99*, 3019–3023.
- Bizzini, A., Zhao, C., Auffray, Y., and Hartke, A. (2009). The Enterococcus faecalis superoxide dismutase is essential for its tolerance to vancomycin and penicillin. *J. Antimicrob. Chemother.* *64*, 1196–1202.
- Borisov, V.B., Forte, E., Davletshin, A., Mastronicola, D., Sarti, P., and Giuffrè, A. (2013). Cytochrome bd oxidase from Escherichia coli displays high catalase activity: an additional defense against oxidative stress. *FEBS Lett.* *587*, 2214–2218.
- Boumghar-Bourtchai, L., Dhalluin, A., Malbruny, B., Galopin, S., and Leclercq, R. (2009). Influence of recombination on development of mutational resistance to linezolid in Enterococcus faecalis JH2-2. *Antimicrob. Agents Chemother.* *53*, 4007–4009.
- Brynildsen, M.P., Winkler, J.A., Spina, C.S., MacDonald, I.C., and Collins, J.J. (2013). Potentiating antibacterial activity by predictably enhancing endogenous microbial ROS production. *Nat. Biotechnol.* *31*, 160–165.
- Cao, Y., Wu, K., Mehta, R., Drew, D.A., Song, M., Lochhead, P., Nguyen, L.H., Izard, J., Fuchs, C.S., Garrett, W.S., et al. (2018). Long-term use of antibiotics and risk of colorectal adenoma. *Gut* *67*, 672–678.
- Cho, H., Uehara, T., and Bernhardt, T. G. (2014). Beta-lactam antibiotics induce a lethal malfunctioning of the bacterial cell wall synthesis machinery. *Cell* *159*, 1300–1311.





- Daneo-Moore, L., Fletcher, S.H., Massida, O., and Pittaluga, F. (1988). Penicillin tolerance in *Enterococcus hirae* ATCC 9790. In Antibiotic inhibition of bacterial cell surface assembly and function, P. Actor, L. Daneo-Moore, M.L. Higgins, M.R.J. Salton, and G.D. Shockmann, eds. (Washington, D.C.: American Society for Microbiology Press), pp. 628–635.
- 5 Dik, V.K., van Oijen, M.G., Smeets, H.M., and Siersema, P.D. (2016). Frequent Use of Antibiotics Is Associated with Colorectal Cancer Risk: Results of a Nested Case-Control Study. *Dig. Dis. Sci.* *61*, 255–264.
- Djorić, D., and Kristich, C.J. (2015). Oxidative stress enhances cephalosporin resistance of *Enterococcus faecalis* through activation of a two-component signaling system. *Antimicrob. Agents Chemother.* *59*, 159–169.
- 10 Donskey, C.J., Chowdhry, T.K., Hecker, M.T., Hoyen, C.K., Hanrahan, J.A., Hujer, A.M., Hutton-Thomas, R.A., Whalen, C.C., Bonomo, R.A., and Rice, L.B. (2000). Effect of antibiotic therapy on the density of vancomycin-resistant enterococci in the stool of colonized patients. *N. Engl. J. Med.* *343*, 1925–1932.
- 15 Dubée, V., Chau, F., Arthur, M., Garry, L., Benadda, S., Mesnage, S., Lefort, A., and Fantin, B. (2011). The in vitro contribution of autolysins to bacterial killing elicited by amoxicillin increases with inoculum size in *Enterococcus faecalis*. *Antimicrob. Agents Chemother.* *55*, 910–912.
- Dwyer, D. J., Kohanski, M. A., and Collins, J. J. (2009). Role of reactive oxygen species in antibiotic action and resistance. *Curr. Opin. Microbiol.* *12*, 482–489.
- 20 Dwyer, D.J., Collins, J.J., and Walker, G.C. (2015). Unraveling the physiological complexities of antibiotic lethality. *Annu. Rev. Pharmacol. Toxicol.* *55*, 313–332.
- Ezraty, B., Vergnes, A., Banzhaf, M., Duverger, Y., Huguenot, A., Brochado, A.R., Su, S.Y., Espinosa, L., Loiseau, L., Py, B., et al. (2013). Fe-S cluster biosynthesis controls uptake of aminoglycosides in a ROS-less death pathway. *Science* *340*, 1583–1587.
- 25 Flahaut, S., Laplace, J.M., Frère, J., and Auffray, Y. (1998). The oxidative stress response in *Enterococcus faecalis*: relationship between H<sub>2</sub>O<sub>2</sub> tolerance and H<sub>2</sub>O<sub>2</sub> stress proteins. *Lett. Appl. Microbiol.* *26*, 259–264.
- Frankenberg, L., Brugna, M., and Hederstedt, L. (2002). *Enterococcus faecalis* heme-dependent catalase. *J. Bacteriol.* *184*, 6351–6356.
- 30 Gutmann, L., and Tomasz, A. (1982). Penicillin-resistant and penicillin-tolerant mutants of group A Streptococci. *Antimicrob. Agents Chemother.* *22*, 128–136.
- Huycke, M.M., Joyce, W., and Wack, M.F. (1996). Augmented production of extracellular superoxide by blood isolates of *Enterococcus faecalis*. *J. Infect. Dis.* *173*, 743–746.
- 35 Huycke, M.M., Moore, D., Joyce, W., Wise, P., Shepard, L., Kotake, Y., and Gilmore, M.S. (2001). Extracellular superoxide production by *Enterococcus faecalis* requires demethylmenaquinone and is attenuated by functional terminal quinol oxidases. *Mol. Microbiol.* *42*, 729–740.







- Huycke, M.M. (2002). Physiology of Enterococci. In *The Enterococci: Pathogenesis, Molecular Biology, and Antibiotic Resistance*, M.S. Gilmore, D.B. Clewell, P. Courvalin, G.M. Dunny, B.E. Murray, and L.B. Rice, eds. (Washington, D.C.: American Society for Microbiology), pp. 133–176.
- 5 Huycke, M.M., Abrams, V., and Moore, D.R. (2002). Enterococcus faecalis produces extracellular superoxide and hydrogen peroxide that damages colonic epithelial cell DNA. *Carcinogenesis* 23, 529–536.
- Huycke, M.M., and Moore, D.R. (2002). In vivo production of hydroxyl radical by Enterococcus faecalis colonizing the intestinal tract using aromatic hydroxylation. *Free Radic. Biol. Med.* 33, 818–826.
- 10
- Jacob, A.E., and Hobbs, S.J. (1974). Conjugal Transfer of Plasmid-Borne Multiple Antibiotic Resistance in Streptococcus faecalis var. zymogenes. *J. Bacteriol.* 117, 360–372.
- Keogh, D., Lam, L.N., Doyle, L.E., Matysik, A., Pavagadhi, S., Umashankar, S., Low, P.M., Dale, J.L., Song, Y., Ng, S.P., et al. (2018). Extracellular Electron Transfer Powers Enterococcus faecalis Biofilm Metabolism. *MBio* 9.
- 15
- Keren, I., Wu, Y., Inocencio, J., Mulcahy, L.R., and Lewis, K. (2013). Killing by bactericidal antibiotics does not depend on reactive oxygen species. *Science* 339, 1213–1216.
- Kim, S.C., Tonkonogy, S.L., Albright, C.A., Tsang, J., Balish, E.J., Braun, J., Huycke, M.M., and Sartor, R.B. (2005). Variable phenotypes of enterocolitis in interleukin 10-deficient mice monoassociated with two different commensal bacteria. *Gastroenterology* 128, 891–906.
- 20
- Kohanski, M.A., Dwyer, D.J., Hayete, B., Lawrence, C.A., and Collins, J.J. (2007). A common mechanism of cellular death induced by bactericidal antibiotics. *Cell* 130, 797–810.
- Krawczyk-Balska, A., Marchlewicz, J., Dudek, D., Wasiak, K., and Samluk, A. (2012). Identification of a ferritin-like protein of Listeria monocytogenes as a mediator of  $\beta$ -lactam tolerance and innate resistance to cephalosporins. *BMC Microbiol.* 12, 278.
- 25
- La Carbona, S., Sauvageot, N., Giard, J.C., Benachour, A., Posteraro, B., Auffray, Y., Sanguinetti, M., and Hartke, A. (2007). Comparative study of the physiological roles of three peroxidases (NADH peroxidase, Alkyl hydroperoxide reductase and Thiol peroxidase) in oxidative stress response, survival inside macrophages and virulence of Enterococcus faecalis. *Mol. Microbiol.* 66, 1148–1163.
- 30
- Ladjouzi, R., Bizzini, A., Lebreton, F., Sauvageot, N., Rincé, A., Benachour, A., and Hartke, A. (2013). Analysis of the tolerance of pathogenic enterococci and Staphylococcus aureus to cell wall active antibiotics. *J. Antimicrob. Chemother.* 68, 2083–2091.
- Liu, Y., Liu, X., Qu, Y., Wang, X., Li, L., and Zhao, X. (2012). Inhibitors of Reactive Oxygen Species Accumulation Delay and/or Reduce the Lethality of Several Antistaphylococcal Agents. *Antimicrob. Agents Chemother.* 56, 6048–6050.
- 35
- Liu, Y., and Imlay, J.A. (2013). Cell death from antibiotics without the involvement of reactive oxygen species. *Science* 339, 1210–1213.

- Maguin, E., Duwat, P., Hege, T., Ehrlich, D., and Gruss, A. (1992). New thermosensitive plasmid for gram-positive bacteria. *J. Bacteriol.* *174*, 5633–5638.
- McDowell, T.D., and Lemanski, C.L. (1988). Absence of autolytic activity (peptidoglycan nicking) in penicillin-induced nonlytic death in a group A streptococcus. *J. Bacteriol.* *170*, 1783–1788.
- 5
- Mercuro, N.J., Davis, S.L., Zervos, M.J., and Herc, E.S. (2018). Combatting resistant enterococcal infections: a pharmacotherapy review. *Expert Opin. Pharmacother.* *19*, 979–992.
- Miller, J.H. (1972). *Experiments in Molecular Genetics* (Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press).
- 10
- Miller, C., Thomsen, L.E., Gaggero, C., Mosseri, R., Ingmer, H., and Cohen, S.N. (2004). SOS response induction by beta-lactams and bacterial defense against antibiotic lethality. *Science* *305*, 1629–1631.
- Moore, D.R., Kotake, Y., and Huycke, M.M. (2004). Effects of iron and phytic acid on production of extracellular radicals by *Enterococcus faecalis*. *Exp. Biol. Med.* (Maywood) *229*, 1186–1195.
- 15
- Pritchard, G.G., and Wimpenny, J.W. (1978). Cytochrome formation, oxygen-induced proton extrusion and respiratory activity in *Streptococcus faecalis* var. *zymogenes* grown in the presence of haematin. *J. Gen. Microbiol.* *104*, 15–22.
- Rigottier-Gois, L., Alberti, A., Houel, A., Taly, J.F., Palcy, P., Manson, J., Pinto, D., Matos, R.C., Carrilero, L., Montero, N., et al. (2011). Large-Scale Screening of a Targeted *Enterococcus faecalis* Mutant Library Identifies Envelope Fitness Factors. *PLoS ONE* *6*.
- 20
- Ritchey, T.W., and Seeley, H.W. (1974). Cytochromes in *Streptococcus faecalis* var. *zymogenes* grown in a haematin-containing medium. *J. Gen. Microbiol.* *85*, 220–228.
- Signoretto, C., Boaretti, M., and Canepari, P. (1994). Cloning, sequencing and expression in *Escherichia coli* of the low-affinity penicillin binding protein of *Enterococcus faecalis*. *FEMS Microbiol. Lett.* *123*, 99–106.
- 25
- Terzaghi, B.E., and Sandine, W.E. (1975). Improved medium for lactic streptococci and their bacteriophages. *Appl. Microbiol.* *29*, 807–813.
- Tipper, D.J., and Strominger, J.L. (1965). Mechanism of action of penicillins: a proposal based on their structural similarity to acyl-D-alanyl-D-alanine. *Proc. Natl. Acad. Sci. U. S. A.* *54*, 1133–1141.
- 30
- Verneuil, N., Mazé, A., Sanguinetti, M., Laplace, J.M., Benachour, A., Auffray, Y., Giard, J.C., and Hartke, A. (2006). Implication of (Mn)superoxide dismutase of *Enterococcus faecalis* in oxidative stress responses and survival inside macrophages. *Microbiology* *152*, 2579–2589.
- 35
- Wang, X., and Huycke, M.M. (2007). Extracellular superoxide production by *Enterococcus faecalis* promotes chromosomal instability in mammalian cells. *Gastroenterology* *132*, 551–561.



- Wang, X., Allen, T.D., May, R.J., Lightfoot, S., Houchen, C.W., and Huycke, M.M. (2008). *Enterococcus faecalis* induces aneuploidy and tetraploidy in colonic epithelial cells through a bystander effect. *Cancer Res.* 68, 9909–9917.
- 5 Wang, X., Yang, Y., Moore, D.R., Nimmo, S.L., Lightfoot, S.A., and Huycke, M.M. (2012). 4-hydroxy-2-nonenal mediates genotoxicity and bystander effects caused by *Enterococcus faecalis*-infected macrophages. *Gastroenterology* 142, 543–551.e7.
- Wise, E.M. Jr, and Park, J.T. (1965). Penicillin: its basic site of action as an inhibitor of a peptide cross-linking reaction in cell wall mucopeptide synthesis. *Proc. Natl. Acad. Sci. U. S. A.* 54, 75–81.
- 10 Yeom, J., Imlay, J.A., and Park, W. (2010). Iron Homeostasis Affects Antibiotic-mediated Cell Death in *Pseudomonas* Species. *J. Biol. Chem.* 285, 22689–22695.











**A****B**

# No $\beta$ -lactam



*E. faecalis*

# $\beta$ -lactams



*E. faecalis*