



HAL
open science

Conic nonholonomic constraints on surfaces and control systems

Timothée Schmoderer, Witold Respondek

► **To cite this version:**

Timothée Schmoderer, Witold Respondek. Conic nonholonomic constraints on surfaces and control systems. 2022. hal-03634034

HAL Id: hal-03634034

<https://hal-normandie-univ.archives-ouvertes.fr/hal-03634034>

Preprint submitted on 2 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Conic nonholonomic constraints on surfaces and control systems

Timothée Schmoderer · Witold Respondek.

Received: date / Accepted: date

Abstract This paper addresses the equivalence problem of conic submanifolds in the tangent bundle of a smooth 2-dimensional manifold. Those are given by a quadratic relation between the velocities and are treated as nonholonomic constraints whose admissible curves are trajectories of the corresponding control systems, called quadratic systems. We deal with the problem of characterising and classifying conic submanifolds under the prism of feedback equivalence of control systems, both control-affine and fully nonlinear. The first main result of this work is a complete description of non-degenerate conic submanifolds via a characterisation under feedback transformations of the novel class of quadratic control-affine systems. This characterisation can explicitly be tested on structure functions defined for any control-affine system and gives a normal form of quadratizable systems and of conic submanifolds. Then, we consider the classification problem of regular conic submanifolds (ellipses, hyperbolas, and parabolas), which is treated via feedback classification of quadratic control-nonlinear systems. Our classification includes several normal forms of quadratic systems (in particular, normal forms not containing functional parameters as well as those containing neither functional nor real parameters), and, as a consequence, gives a classification of regular conic submanifolds.

Keywords nonlinear control system · feedback equivalence · conic submanifolds · nonholonomic constraint · normal forms · pseudo-Riemannian geometry

Mathematics Subject Classification (2020) 93B52 · 37N35 · 93A10 · 93B27 · 53B20 · 53B30

Contents

1	Introduction	1
2	Preliminaries	4
3	Quadratizable control-affine systems	9
3.1	Characterisation of quadratizable control-affine systems	12
3.2	Normal form of quadratizable control-affine system	13
4	Classification of quadratic systems	16
4.1	Classification of elliptic and hyperbolic systems	18
4.2	Classification of parabolic systems	23
5	Conclusions	26
A	Simultaneous rectification of two rank 1 distributions in \mathbb{R}^2	28
B	Resolution of equation (6)	28
C	Detailed computation of the proof of Theorem 3.2	28
C.1	Resolution of equation (10)	28
C.2	Smooth form of h'	29
D	Details of the computations of Lemma 4.2	29
E	Gaussian curvature for metric given in terms of vector fields	29

1 Introduction

Let \mathcal{X} be a smooth connected manifold of dimension $n = 2$ (a surface), equipped with local coordinates x . In the tangent bundle $T\mathcal{X}$ of \mathcal{X} , we consider a smooth 3-dimensional submanifold \mathcal{S} , a hypersurface, given

by

$$\mathcal{S} = \{(x, \dot{x}) \in T\mathcal{X}, S(x, \dot{x}) = 0\},$$

where $S : T\mathcal{X} \rightarrow \mathbb{R}$ is a smooth scalar function satisfying $\text{rk} \frac{\partial S}{\partial \dot{x}}(x, \dot{x}) = 1$ for all $(x, \dot{x}) \in \mathcal{S}$. Two submanifolds $\mathcal{S} = \{S(x, \dot{x}) = 0\} \subset T\mathcal{X}$ and $\tilde{\mathcal{S}} = \{\tilde{S}(\tilde{x}, \dot{\tilde{x}}) = 0\} \subset T\tilde{\mathcal{X}}$ are said to be equivalent if there exists a diffeomorphism $\phi : \mathcal{X} \rightarrow \tilde{\mathcal{X}}$ and a smooth nonvanishing function $\delta : T\mathcal{X} \rightarrow \mathbb{R}$ such that

$$\tilde{S}(\phi(x), D\phi(x)\dot{x}) = \delta(x, \dot{x}) S(x, \dot{x}),$$

where $D\phi$ is the derivative (tangent map) of ϕ . This definition implies that for all $(x, \dot{x}) \in \mathcal{S}$ we have $(\tilde{x}, \dot{\tilde{x}}) = (\phi(x), D\phi(x)\dot{x}) \in \tilde{\mathcal{S}}$, hence the application $(\phi, D\phi)$ maps the graph of $S^{-1}(0)$ into that of $\tilde{S}^{-1}(0)$. Equivalence of submanifolds \mathcal{S} and $\tilde{\mathcal{S}}$ means simply that implicit underdetermined ordinary differential equations $S(x, \dot{x}) = 0$ and $\tilde{S}(\tilde{x}, \dot{\tilde{x}}) = 0$ are equivalent; see e.g. [2, Definition 2].

It is natural to ask how to characterise and classify submanifolds \mathcal{S} of certain particular classes, for instance the class of linear submanifolds given by $S_{lin}(x, \dot{x}) = \omega(x)\dot{x} = 0$ or the class of affine submanifolds given by $S_{aff}(x, \dot{x}) = \omega(x)\dot{x} + h(x) = 0$, where ω is a smooth differential 1-form on \mathcal{X} and h is a smooth function on \mathcal{X} . Those questions have been widely studied under the prism of Pfaffian equations (linear and affine) and go back to Pfaff, Darboux, Cartan [15, 5, 6]. Although the problem of classification of Pfaffian equations is still open in its full generality, many important results have been obtained for various classes of linear Pfaffian equations (contact and quasi-contact case, Martinet case, singularities, [24, 25, 13, 10, 8]) and of affine Pfaffian equations (dimension two [9], three [17, 16], and arbitrary [26, 8]).

We will call \mathcal{S}_q a quadratic, or a conic, submanifold if it is given by

$$\mathcal{S}_q(x, \dot{x}) = \dot{x}^t \mathbf{g}(x) \dot{x} + 2\omega(x)\dot{x} + h(x) = 0,$$

with all involved objects being smooth, i.e. for each point $x \in \mathcal{X}$, the set $\mathcal{S}_q(x)$ forms a conic curve in $T_x\mathcal{X}$. The image of a quadratic submanifold $\mathcal{S}_q = \{S_q = 0\}$ under a diffeomorphism $\tilde{x} = \phi(x)$ is a quadratic submanifold $\tilde{\mathcal{S}}_q = \{\tilde{S}_q = 0\}$, where $\tilde{S}_q(\tilde{x}, \dot{\tilde{x}}) = \dot{\tilde{x}}^t \tilde{\mathbf{g}} \dot{\tilde{x}} + \tilde{\omega} \dot{\tilde{x}} + \tilde{h}$, with $\mathbf{g} = \phi^* \tilde{\mathbf{g}}$, $\omega = \phi^* \tilde{\omega}$, and $h = \phi^* \tilde{h}$. On the other hand, a submanifold $\tilde{\mathcal{S}}$ equivalent to a quadratic \mathcal{S}_q is given by $\phi^* \tilde{S} = \delta \cdot S_q = 0$ and $\tilde{\mathcal{S}}$ need not be quadratic because of the presence of the function δ . The first purpose of this work is to provide a local characterisation of submanifolds \mathcal{S} that are equivalent to quadratic submanifolds \mathcal{S}_q . In particular, we will identify submanifolds equivalent to elliptic, hyperbolic, and parabolic conics, given respectively by

$$\begin{aligned} \mathcal{S}_E = \left\{ a^2(z - c_0)^2 + b^2(y - c_1)^2 = 1 \right\}, \quad \mathcal{S}_H = \left\{ a^2(z - c_0)^2 - b^2(y - c_1)^2 = 1 \right\}, \\ \text{and } \mathcal{S}_P = \left\{ ay^2 - z + by + c = 0 \right\}, \end{aligned}$$

where a, b, c_0, c_1 , and c , are smooth functions of $x = (z, y)$, we choose the order (z, y) to be consistent with some normal forms existing in the literature, satisfying $a(\cdot) \neq 0$ and $b(\cdot) \neq 0$ in the elliptic and hyperbolic cases and $a(\cdot) \neq 0$ in the parabolic case. We will also discuss the case of passing smoothly from one type to another. To state and discuss general facts on elliptic, hyperbolic, and parabolic submanifolds we set $\mathcal{S}_Q = \{\mathcal{S}_E, \mathcal{S}_H, \mathcal{S}_P\}$.

The second goal of this work is to provide a classification of elliptic \mathcal{S}_E , hyperbolic \mathcal{S}_H , and parabolic \mathcal{S}_P conics. In the elliptic and hyperbolic cases, we will first describe the submanifolds with the functions $a(x) = b(x)$ and call them *conformally-flat*; second, if, additionally, $a = b = 1$, then we call them *flat* elliptic or hyperbolic submanifold. Finally, we will describe the forms for which we also have $c_0, c_1 \in \mathbb{R}$ (in the same coordinate system as the one for which we have $a = b = 1$), called a *constant-form* elliptic and hyperbolic submanifolds; the particular case of $c_0 = c_1 = 0$ is called a *null-form* to emphasis the absence of any (functional, continuous, or discrete) parameter. In the parabolic case, we will describe the submanifolds with $a = 1$, called *weakly-flat*, and characterise those with, additionally, $b = 0$ (called *strongly-flat*) and, moreover, with $c \in \mathbb{R}$ (called *constant-form*); in the particular case where $c = 0$ we call the parabolic submanifold a *null-form*. Our classification is summarised by Table 1 in Section 4, where a characterisation of all above subclasses is given.

Our analysis of the equivalence problem of submanifolds will be based on attaching to a submanifold $\mathcal{S} = \{S(x, \dot{x}) = 0\} \subset T\mathcal{X}$ two control systems. First,

$$\Xi_{\mathcal{S}} : \dot{x} = F(x, w), \quad x \in \mathcal{X}, \quad w \in \mathcal{W} \subset \mathbb{R},$$

where $\dot{x} - F(x, w) = 0$ is a regular parametric representation of \mathcal{S} , that is for all values of the parameter $w \in \mathcal{W}$ (interpreted as a scalar control) we have $S(x, F(x, w)) = 0$ and $\text{rk} \frac{\partial F}{\partial w}(x, w) = 1$, and second,

$$\Sigma_{\mathcal{S}} : \begin{cases} \dot{x} = F(x, w) \\ \dot{w} = u \end{cases}, \quad (x, w) \in \mathcal{X} \times \mathcal{W}, \quad u \in \mathbb{R},$$

called, respectively, a first and second extension of \mathcal{S} . Notice that for $\Xi_{\mathcal{S}}$ the control w enters in a nonlinear way, whereas for $\Sigma_{\mathcal{S}}$ the control u enters in an affine way, but for the price of augmentation of the dimension of the state space. Observe that, since $S(x, \dot{x}) = 0$, defining \mathcal{S} , relates the positions x with the velocities \dot{x} , it describes a nonholonomic constraint. We say that a smooth curve $x(t) \in \mathcal{X}$ satisfies the nonholonomic constraint given by \mathcal{S} if we have $(x(t), \dot{x}(t)) \in \mathcal{S}$. Clearly, $x(t)$ satisfies the nonholonomic constraint described by \mathcal{S} (equivalently, satisfies the implicit differential equation $S(x(t), \dot{x}(t)) = 0$) if and only if $x(t)$ is a trajectory of $\Xi_{\mathcal{S}}$ for a certain smooth control $w(t)$ or, equivalently, $(x(t), w(t))$ is a trajectory of $\Sigma_{\mathcal{S}}$ for a smooth control $u(t)$. A crucial observation that links studying submanifolds $\mathcal{S} \subset T\mathcal{X}$ and their extensions $\Xi_{\mathcal{S}}$ and $\Sigma_{\mathcal{S}}$ is that the equivalence of submanifolds corresponds to the equivalence of control systems $\Xi_{\mathcal{S}}$ and $\Sigma_{\mathcal{S}}$ via feedback transformations, general for $\Xi_{\mathcal{S}}$, and control-affine for $\Sigma_{\mathcal{S}}$, as assured by [Proposition 2.1](#).

Organisation of the paper. In the next section, we will recall some definitions of control theory and we will show that the problem of characterising and classifying submanifolds of $T\mathcal{X}$ can be replaced by that of characterising and classifying their first and second extensions $\Xi_{\mathcal{S}}$ and $\Sigma_{\mathcal{S}}$ under feedback transformations (see [Proposition 2.1](#)). Moreover, we will give a first rough classification of non-degenerate conic submanifolds, introducing elliptic \mathcal{S}_E , hyperbolic \mathcal{S}_H , and parabolic \mathcal{S}_P subclasses (see [Lemma 2.1](#)). In [Section 3](#), we will define a general second extension of a conic submanifold \mathcal{S}_q , called a quadratic system Σ_q . Just as we do for conic submanifolds, we will identify elliptic Σ_E , hyperbolic Σ_H , and parabolic Σ_P systems as particular cases of Σ_q , see [Definition 3.1](#) and [Proposition 3.1](#). In [Theorem 3.1](#), we will fully characterise the class of quadratic systems by means of a checkable relation between well-defined structure functions attached to any control-affine system. The conditions obtained in that theorem allow to give a normal form for all quadratizable control-affine systems (i.e. systems Σ that are feedback equivalent to Σ_q), see [Theorem 3.2](#), which in turn leads to a normal form for all non-degenerate conic submanifolds \mathcal{S}_q . We will also show how our characterisation and the normal form apply to the classes of elliptic, hyperbolic, and parabolic control-affine systems, which gives us a deeper insight into our conditions and in our normal form (see [Corollary 3.1](#) and [Corollary 3.2](#)). Finally, in [Section 4](#), we will be interested in the classification of elliptic, hyperbolic, and parabolic submanifolds as presented in [Table 1](#) given there. This problem is dealt with using the classification of their first extensions (treated as control-nonlinear systems) under feedback transformations. We first show that the classification of elliptic, hyperbolic, and parabolic submanifolds presented in [Table 1](#) in [Section 4](#) is reflected in properties of a triple of vector fields attached to their parametrisations; see [Lemma 4.1](#) statements *(i)* to *(iv)* for the elliptic and hyperbolic cases and statements *(v)* to *(viii)* for the parabolic case. To every quadratic control-nonlinear system Ξ_E , Ξ_H , or Ξ_P (first extensions of elliptic, hyperbolic, and parabolic submanifolds, respectively), we will attach a frame of the tangent bundle (see the paragraph before [Proposition 4.1](#)) and we give conditions for that frame to be commutative: it turns out that in the elliptic and hyperbolic cases this requires that a certain pseudo-Riemannian metric is flat (see [Proposition 4.3](#)), whereas in the parabolic case this problem can be solved without any extra assumptions (see [Proposition 4.5](#)). Then we show how we can additionally normalize the systems while preserving the commutativity of that frame. Our classifications include several normal and canonical forms, given by [Proposition 4.4](#) for elliptic and hyperbolic systems and by [Theorem 4.1](#) for parabolic systems. We summarise the structure of the paper in [figure 1](#).

Related works. A classification of quadratic control systems was initiated by Bonnard in [\[3\]](#). His work differs from our as he considered homogeneous systems of degree 2 with respect to all state variables. Hence, his class of quadratic control systems is a subclass of our parabolic systems (where we require that only one variable enters quadratically) but he considers the general dimension n while our results concern 3-dimensional systems only. In [\[12\]](#), Krener and Kang studied the problem of equivalence, via feedback, to polynomial systems of degree 2 modulo higher order terms. This work was continued in [\[11\]](#) and [\[21\]](#) for any degree but all those results are given for formal classification only. Examples of control systems subject to conic nonholonomic constraints appear in various domains of physics and engineering applications. In the next section, we will discuss Dubin's car [\[7\]](#) which is a simple model of a vehicle, as well as its hyperbolic counterpart [\[14\]](#). We also mention [\[27\]](#), where the planar tilting manoeuvre problem is considered under small angle assumption, the studied control system is elliptic with respect to the states.

2 Preliminaries

Main notations.




Fig. 1 Walk-through the paper with separation between characterisation and classification results

$\mathcal{X}, T\mathcal{X}, x = (z, y)$	A smooth 2d manifold, its tangent bundle, and its local coordinates.
$\mathcal{M}, \xi = (x, w) = (z, y, w)$	A smooth 3d manifold and its local coordinates.
ϕ, ϕ_*	A diffeomorphism and its tangent map.
\mathcal{S}	Smooth submanifold of $T\mathcal{X}$ given by an equation of the form $S(x, \dot{x}) = 0$, with $\text{rk} \frac{\partial S}{\partial \dot{x}}(x, \dot{x}) = 1$.
S_q	Quadratic, conic, submanifold described by $S_q(x, \dot{x}) = \dot{x}^t \mathbf{g}(x) \dot{x} + 2\omega(x)\dot{x} + h(x) = 0$.
$S_Q = \{S_E, S_H, S_P\}$	Set of elliptic, hyperbolic, and parabolic submanifolds.
Ξ_S	Regular parametrisation of a submanifold \mathcal{S} , called a first extension, seen as a control-nonlinear system.
Σ_S	Extension of Ξ_S , called a second extension of a submanifold \mathcal{S} , seen as a control-affine system on a three dimensional manifold \mathcal{M} .

In this section, we introduce all tools and concepts that we will need in this paper. First, we recall some notions on the equivalence of submanifolds (of the tangent bundle) and of control systems. Second, most importantly, we show that the problem of equivalence of submanifolds can be replaced by that of equivalence of their first and second extensions, see [Proposition 2.1](#). Finally, we present quadratic submanifolds and we introduce the subclasses of elliptic, hyperbolic, and parabolic submanifolds. In the paper the word *smooth* will always mean C^∞ -smooth and throughout all systems, functions, manifolds and submanifolds are assumed to be smooth.

Equivalence notions. Recall that we have defined two submanifolds $\mathcal{S} = \{S(x, \dot{x}) = 0\} \subset T\mathcal{X}$ and $\tilde{\mathcal{S}} = \{\tilde{S}(\tilde{x}, \tilde{\dot{x}}) = 0\} \subset T\tilde{\mathcal{X}}$ to be equivalent if there exists a diffeomorphism $\phi : \mathcal{X} \rightarrow \tilde{\mathcal{X}}$ and a smooth nonvanishing function $\delta : T\mathcal{X} \rightarrow \mathbb{R}$ such that

$$\tilde{S}(\phi(x), D\phi(x)\dot{x}) = \delta(x, \dot{x}) S(x, \dot{x}).$$

If ϕ is defined in a neighbourhood \mathcal{X}_0 of x_0 and δ is defined in a neighbourhood of $(x_0, \dot{x}_0) \in \mathcal{S}$ only, then we say that \mathcal{S} and $\tilde{\mathcal{S}}$ are locally equivalent at (x_0, \dot{x}_0) and $(\tilde{x}_0, \tilde{\dot{x}}_0) = (\phi(x_0), D\phi(x_0)\dot{x}_0)$, respectively.

Example 1. The submanifolds \mathcal{S} and $\tilde{\mathcal{S}}$ given by $S(x, \dot{x}) = \dot{z} - (-1 + \sqrt{1 + \dot{y}})^2 = 0$, around $(x_0, \dot{x}_0) = (0, 0)$, and $\tilde{S}(\tilde{x}, \tilde{\dot{x}}) = \tilde{\dot{z}} - \left(\frac{\tilde{\dot{y}}}{2}\right)^2 = 0$, around $(\tilde{x}_0, \tilde{\dot{x}}_0) = (0, 0)$, respectively, are equivalent via

$$(\tilde{z}, \tilde{y}) = \phi(x) = (z, y - z) \quad \text{and} \quad \delta(x, \dot{x}) = -\frac{1}{4} \left(\dot{z} - \dot{y} - 2 - 2\sqrt{1 + \dot{y}} \right).$$

Example 2. Since on a 2-dimensional manifold all metrics are locally conformally flat (see [\[20, Addendum 1 of Chapter 9\]](#) for the Riemannian case and [\[19, Theorem 7.2\]](#) for the Lorentzian one), it follows that the submanifold given by $S(x, \dot{x}) = a(x)\dot{z}^2 + 2b(x)\dot{z}\dot{y} + c(x)\dot{y}^2 - 1 = 0$ (where $ac - b^2 \neq 0$) is locally equivalent to that given by $\tilde{S}(\tilde{x}, \tilde{\dot{x}}) = \left(\tilde{\dot{z}}^2 \pm \tilde{\dot{y}}^2\right) - r(\tilde{x})^2 = 0$.

Example 3. Since $\text{rk} \frac{\partial S}{\partial \dot{x}}(x, \dot{x}) = 1$, assume that $\frac{\partial S}{\partial \dot{x}}(x_0, \dot{x}_0) \neq 0$; recall that $x = (z, y)$. By the implicit function theorem, we can locally write $S(x, \dot{x}) = \delta(x, \dot{x})(\dot{z} - s(x, \dot{y}))$, where $\delta(x_0, \dot{x}_0) \neq 0$. So, a submanifold \mathcal{S} is always locally equivalent to the one given by $\dot{z} - s(x, \dot{y}) = 0$.

Two general control systems $\Xi : \dot{x} = F(x, w)$ and $\tilde{\Xi} : \dot{\tilde{x}} = \tilde{F}(\tilde{x}, \tilde{w})$, not necessarily coming from parametrisations of submanifolds, are called *feedback equivalent* if there exists a diffeomorphism $\Phi : \mathcal{X} \times \mathcal{W} \rightarrow \tilde{\mathcal{X}} \times \tilde{\mathcal{W}}$ of the form

$$(\tilde{x}, \tilde{w}) = \Phi(x, w) = (\phi(x), \psi(x, w)),$$

which transforms the first system into the second, i.e.

$$D\phi(x)F(x, w) = \tilde{F}(\phi(x), \psi(x, w)).$$

The map $\phi(x)$ plays the role of a coordinates change in the state space \mathcal{X} , and the map ψ is called feedback as it changes the parametrisation by control w in a way that depends on the state x . If the diffeomorphism Φ is defined in a neighbourhood of (x_0, w_0) only, and $\Phi(x_0, w_0) = (\tilde{x}_0, \tilde{w}_0)$, then we say that the two systems are *locally feedback equivalent* at (x_0, w_0) and $(\tilde{x}_0, \tilde{w}_0)$, respectively. If the diffeomorphism Φ is defined on the product $\mathcal{X}_0 \times \mathcal{W}$, that is, it is global with respect to w , where \mathcal{X}_0 is a neighbourhood of a state x_0 , and $\phi(x_0) = \tilde{x}_0$, then we say that Ξ and $\tilde{\Xi}$ are locally feedback equivalent at x_0 and \tilde{x}_0 , respectively. The latter local feedback equivalence will be especially useful for the class of control-affine systems

$$\Sigma : \dot{\xi} = f(\xi) + g(\xi)u, \quad u \in \mathbb{R},$$

where f and g are smooth vector fields on \mathcal{M} (equipped with local coordinates ξ); below the state space \mathcal{M} will be of dimension 3 so \mathcal{M} is not \mathcal{X} , which is of dimension 2. For control-affine systems we will restrict the feedback transformations to the control-affine transformations

$$\psi(\xi, u) = \alpha(\xi) + \beta(\xi)u,$$

where $\alpha(\xi)$ and $\beta(\xi)$ are smooth functions satisfying $\beta(\cdot) \neq 0$. In that case, we denote the feedback transformation by the triple (ϕ, α, β) and if $\phi = \text{id}$, then this action is called a *pure feedback* transformation and is denoted (α, β) . Observe that a general control system $\Xi : \dot{x} = F(x, w)$ can be extended to a control-affine system Ξ^e by augmenting the state space with the control w and introducing the new control $u = \dot{w}$, which gives

$$\Xi^e : \begin{cases} \dot{x} = F(x, w) \\ \dot{w} = u \end{cases}, \quad u \in \mathbb{R}.$$

Notice that Ξ^e lives on the manifold $\mathcal{M} = \mathcal{X} \times \mathcal{W}$ of dimension $n = 3$.

In introduction, we already attached to a submanifold $\mathcal{S} \subset T\mathcal{X}$ a regular parametrisation given by a control-nonlinear system $\Xi_{\mathcal{S}} : \dot{x} = F(x, w)$ satisfying $S(x, F(x, w)) = 0$ and $\text{rk} \frac{\partial F}{\partial w}(x, w) = 1$. Since constructing $\Xi_{\mathcal{S}}$ requires to introduce an extra variable w (a control), we will also call it a *first extension* of \mathcal{S} . Then we can attach to $\Xi_{\mathcal{S}}$ its extension $\Xi_{\mathcal{S}}^e$, denoted $\Sigma_{\mathcal{S}}$ and given by

$$\Sigma_{\mathcal{S}} : \begin{cases} \dot{x} = F(x, w) \\ \dot{w} = u \end{cases}, \quad u \in \mathbb{R},$$

and called a *second extension* of \mathcal{S} . To distinguish different control systems attached to the submanifolds \mathcal{S} and $\tilde{\mathcal{S}}$, we will denote $\Xi_{\mathcal{S}}$ (resp. $\Sigma_{\mathcal{S}}$) by $\tilde{\Xi}_{\tilde{\mathcal{S}}}$ (resp. by $\tilde{\Sigma}_{\tilde{\mathcal{S}}}$). The following proposition shows that the problem of equivalence of submanifolds corresponds to the equivalence under feedback transformations of their corresponding first and second extensions.

Proposition 2.1 (Equivalence of equivalence relations) *Consider submanifolds $\mathcal{S} = \{S(x, \dot{x}) = 0\}$ of $T\mathcal{X}$ and $\tilde{\mathcal{S}} = \{\tilde{S}(\tilde{x}, \dot{\tilde{x}}) = 0\}$ of $T\tilde{\mathcal{X}}$. The following statements are locally equivalent:*

- (i) *The submanifolds \mathcal{S} and $\tilde{\mathcal{S}}$ are equivalent via $\phi(x)$ and $\delta(x, \dot{x})$.*
- (ii) *Their regular parametrisations (first extensions) $\Xi_{\mathcal{S}}$ and $\tilde{\Xi}_{\tilde{\mathcal{S}}}$ are feedback equivalent via $\phi(x)$ and $\psi(x, w)$.*
- (iii) *Their second extensions $\Sigma_{\mathcal{S}}$ and $\tilde{\Sigma}_{\tilde{\mathcal{S}}}$ are feedback equivalent via $\varphi(x, w) = (\phi(x), \psi(x, w))$ and (α, β) .*

That is, the following diagram is commutative:

$$\begin{array}{ccccc} \mathcal{S} & \xrightarrow{\text{parametrisation}} & \Xi_{\mathcal{S}} & \xrightarrow{\text{extension}} & \Sigma_{\mathcal{S}} \\ (\phi, \delta) \downarrow & & (\phi, \psi) \downarrow & & \downarrow (\varphi, \alpha, \beta) \\ \tilde{\mathcal{S}} & \xrightarrow{\text{parametrisation}} & \tilde{\Xi}_{\tilde{\mathcal{S}}} & \xrightarrow{\text{extension}} & \tilde{\Sigma}_{\tilde{\mathcal{S}}} \end{array}$$

Proof (i) ⇒ (ii). To show that the equivalence of submanifolds implies the equivalence under feedback of their first extensions, we will first use a diffeomorphism (of the state space) to analyse both first extensions in the same coordinate system. Then, we will show that the parameters (controls) of those two first extensions are related by a pure feedback transformation. To obtain invertibility of the feedback transformation, our argument will strongly rely on the regularity of the submanifold, $\text{rk} \frac{\partial S}{\partial \dot{x}} = 1$, and of their first extensions, $\text{rk} \frac{\partial F}{\partial w} = 1$. Assume that two submanifolds \mathcal{S} and $\tilde{\mathcal{S}}$ given, respectively, by $S(x, \dot{x}) = 0$ and $\tilde{S}(\tilde{x}, \dot{\tilde{x}}) = 0$ are equivalent via a diffeomorphism $\tilde{x} = \phi(x)$ and a nonvanishing function $\delta(x, \dot{x})$, that is $\tilde{S}(\phi(x), D\phi(x)\dot{x}) = \delta(x, \dot{x})S(x, \dot{x})$. Consider $\Xi_{\mathcal{S}} : \dot{x} = F(x, w)$ and $\tilde{\Xi}_{\tilde{\mathcal{S}}} : \dot{\tilde{x}} = \tilde{F}(\tilde{x}, \tilde{w})$, two regular parametrisations of those submanifolds. Then, using $0 = \tilde{S}(\tilde{x}, \dot{\tilde{x}}) = \tilde{S}(\tilde{x}, \tilde{F}(\tilde{x}, \tilde{w}))$, we have

$$\tilde{S}(\phi(x), \tilde{F}(\phi(x), \tilde{w})) = \delta(x, (D\phi(x))^{-1} \tilde{F}(\phi(x), \tilde{w}))S(x, (D\phi(x))^{-1} \tilde{F}(\phi(x), \tilde{w})), \quad \forall \tilde{w} \in \tilde{W},$$

implying $S(x, \hat{F}(x, \tilde{w})) = 0$, where $\hat{F}(x, \tilde{w}) = (D\phi(x))^{-1} \tilde{F}(\phi(x), \tilde{w})$. Therefore, $\dot{x} = F(x, w)$ and $\dot{x} = \hat{F}(x, \tilde{w})$ are two regular parametrisations of the same submanifold \mathcal{S} . We will prove that $F(x, w)$ and $\hat{F}(x, \tilde{w})$ are related by an invertible (pure) feedback transformation of the form $\tilde{w} = \psi(x, w)$.

Since $\text{rk} \frac{\partial S}{\partial \dot{x}}(x_0, \dot{x}_0) = 1$, we may assume that $\frac{\partial S}{\partial \dot{z}}(x_0, \dot{x}_0) \neq 0$, where $x = (z, y)$ and, by the implicit function theorem, we have $S(x, \dot{x}) = \delta(x, \dot{x})(\dot{z} - s(x, \dot{y})) = 0$. Let $\dot{z} = F_1(x, w)$, $\dot{y} = F_2(x, w)$ be a regular parametrisation of \mathcal{S} , it follows that $\frac{\partial F_2}{\partial w}(x_0, w_0) \neq 0$. Indeed, if $\frac{\partial F_2}{\partial w}(x_0, w_0) = 0$, then $\dot{z} - s(x, \dot{y}) = 0$ (recall that $\delta \neq 0$), hence $F_1 - s(x, F_2) = 0$ and thus $\frac{\partial F_1}{\partial w} - \frac{\partial s}{\partial y} \frac{\partial F_2}{\partial w} = 0$ implying that $\frac{\partial F_1}{\partial w}(x_0, w_0) = 0$; contradiction.

Hence, the two regular parametrisations of $\mathcal{S} = \{S(x, \dot{x}) = 0\}$ given by

$$\Xi_{\mathcal{S}} : \begin{cases} \dot{z} = F_1(x, w) \\ \dot{y} = F_2(x, w) \end{cases} \quad \text{and} \quad \hat{\Xi}_{\mathcal{S}} : \begin{cases} \dot{z} = \hat{F}_1(x, \tilde{w}) \\ \dot{y} = \hat{F}_2(x, \tilde{w}) \end{cases}$$

satisfy $\frac{\partial F_2}{\partial w}(x_0, w_0) \neq 0$ and $\frac{\partial \hat{F}_2}{\partial \tilde{w}}(x_0, \tilde{w}_0) \neq 0$. Therefore, $\tilde{w} = \hat{F}_2^{-1}(x, \dot{y}) = \hat{F}_2^{-1}(x, F_2(x, w))$, where \hat{F}_2^{-1} is the inverse with respect to the second argument. And using $\dot{z} = s(x, \dot{y})$ we obtain $\dot{z} = F_1(x, w) = s(x, \dot{y}) = \hat{F}_1(x, \tilde{w})$. Thus, $\Xi_{\mathcal{S}}$ and $\hat{\Xi}_{\mathcal{S}}$ are feedback equivalent via $\tilde{w} = \psi(x, w)$ with $\psi(x, w) = \hat{F}_2^{-1}(x, F_2(x, w))$ and the systems $\Xi_{\mathcal{S}}$ and $\tilde{\Xi}_{\tilde{\mathcal{S}}}$ are feedback equivalent since $\tilde{\Xi}_{\tilde{\mathcal{S}}}$ is the system $\hat{\Xi}_{\mathcal{S}}$ mapped via the diffeomorphism $\tilde{x} = \phi(x)$.

(ii) ⇒ (i). Assume that the two regular parametrisations $\Xi_{\mathcal{S}} : \dot{x} = F(x, w)$ and $\tilde{\Xi}_{\tilde{\mathcal{S}}} : \dot{\tilde{x}} = \tilde{F}(\tilde{x}, \tilde{w})$ of \mathcal{S} and $\tilde{\mathcal{S}}$, respectively, are feedback equivalent via $\tilde{x} = \phi(x)$ and $\tilde{w} = \psi(x, w)$. Denote $\Phi = \phi^{-1}$, and apply the diffeomorphism $x = \Phi(\tilde{x})$ to $\tilde{\Xi}_{\tilde{\mathcal{S}}}$ to obtain a new vector field (parametrized by \tilde{w}) $\hat{F}(x, \tilde{w}) = D\Phi(\phi(x))\tilde{F}(\phi(x), \tilde{w})$ related with $F(x, w)$ by the pure feedback transformation $\tilde{w} = \psi(x, w)$. Denote $F = (F_1, F_2)^t$ and $\hat{F} = (\hat{F}_1, \hat{F}_2)^t$ in the $x = (z, y)$ coordinates. Without loss of generality we can assume that $\frac{\partial F_2}{\partial w}(x_0, w_0) \neq 0$ and $\frac{\partial \hat{F}_2}{\partial \tilde{w}}(x_0, \tilde{w}_0) \neq 0$.

Now, apply to $\tilde{\mathcal{S}}$ the same diffeomorphism $x = \Phi(\tilde{x})$, whose inverse we denoted by $\tilde{x} = \phi(x)$, and set $\hat{S}(x, \dot{x}) = \tilde{S}(\phi(x), D\phi(x)\dot{x})$. Since by definition of $\tilde{\Xi}_{\tilde{\mathcal{S}}}$ we have $\tilde{S}(\tilde{x}, \tilde{F}(\tilde{x}, \tilde{w})) = 0$, we conclude that $\hat{S}(x, \hat{F}(x, \tilde{w})) = 0$. Then we claim that $\frac{\partial \hat{S}}{\partial \dot{z}}(x_0, \dot{x}_0) \neq 0$; indeed, if $\frac{\partial \hat{S}}{\partial \dot{z}}(x_0, \dot{x}_0) = 0$, then $\frac{\partial S}{\partial w}S(x, F(x, w)) = 0$ yields $\frac{\partial S}{\partial z} \frac{\partial F_1}{\partial w} + \frac{\partial S}{\partial y} \frac{\partial F_2}{\partial w} = 0$ and thus $\frac{\partial S}{\partial y}(x_0, \dot{x}_0) = 0$ giving a contradiction. The same observation holds for $\frac{\partial \hat{S}}{\partial \dot{z}}(x_0, \dot{x}_0)$. Thus by the implicit function theorem we have $S(x, \dot{x}) = \delta(x, \dot{x})(\dot{z} - s(x, \dot{y}))$ and $\hat{S}(x, \dot{x}) = \hat{\delta}(x, \dot{x})(\dot{z} - \hat{s}(x, \dot{y}))$, with $\delta \neq 0$ and $\hat{\delta} \neq 0$.

Using $\dot{z} = s(x, \dot{y})$ and $\dot{z} = \hat{s}(x, \dot{y})$, we obtain $s(x, \dot{y}) = F_1(x, w) = \hat{F}_1(x, \tilde{w}) = \hat{s}(x, \dot{y})$, hence we have

$$S(x, \dot{x}) = \delta(x, \dot{x})(\dot{z} - s(x, \dot{y})) = \delta(x, \dot{x})(\dot{z} - \hat{s}(x, \dot{y})) = \frac{\delta}{\hat{\delta}} \tilde{S}(\phi(x), D\phi(x)\dot{x})$$

establishing the equivalence between \mathcal{S} and $\tilde{\mathcal{S}}$.

(ii) ⇒ (iii). If $\Xi_{\mathcal{S}}$ and $\tilde{\Xi}_{\tilde{\mathcal{S}}}$ are feedback equivalent, then

$$D\phi(x)F(x, w) = \tilde{F}(\phi(x), \psi(x, w)).$$

Thus the diffeomorphism $(\phi(x), \psi(x, w))$, of the augmented state space (x, w) , together with the control-affine feedback

$$\tilde{u} = \frac{\partial \psi}{\partial x} F(x, w) + \frac{\partial \psi}{\partial w} u$$

transform $\Sigma_{\mathcal{S}}$ into $\tilde{\Sigma}_{\tilde{\mathcal{S}}}$.

(iii) \Rightarrow (ii). Assume that $\Sigma_{\mathcal{S}}$ into $\tilde{\Sigma}_{\tilde{\mathcal{S}}}$ are feedback equivalent via $(\tilde{x}, \tilde{w}) = \Phi(x, w)$ and $\tilde{u} = \alpha(x, w) + \beta(x, w)u$. Since the distribution $\text{span}\{\frac{\partial}{\partial w}\}$ is sent by ϕ_* into $\text{span}\{\frac{\partial}{\partial \tilde{w}}\}$, it follows that Φ has the triangular form $(\phi(x), \psi(x, w))$. Therefore, feedback equivalence of the systems $\Xi_{\mathcal{S}}$ and $\tilde{\Xi}_{\tilde{\mathcal{S}}}$ is established via the diffeomorphism $\tilde{x} = \phi(x)$ and the reparametrisation $\tilde{w} = \psi(x, w)$. \square

Remark 2.1 The use of extensions of control-nonlinear systems was introduced in [22] and used to study controllability and observability of nonlinear systems, and then to analyse linearizability and decoupling [23]. Moreover, notice that the same proof as that of (ii) \Leftrightarrow (iii) shows that any two control-nonlinear systems Ξ and $\tilde{\Xi}$ (which need not be regular parametrisations of submanifolds) are feedback equivalent if and only if their extensions Ξ^e and $\tilde{\Xi}^e$ are equivalent via control-affine feedback; see [9].

Remark 2.2 The equivalence (i) \Leftrightarrow (ii) does not hold, in general, if the parametrisation $\Xi_{\mathcal{S}} : \dot{x} = F(x, w)$ does not satisfy the regularity condition $\frac{\partial F}{\partial w}(x, w) \neq 0$ that we assume. To see that, consider the submanifold \mathcal{S} given by $\dot{z} - \dot{y}^2 = 0$. The parametrisations of \mathcal{S}

$$\Xi_{\mathcal{S}} : \begin{cases} \dot{z} = w^2 \\ \dot{y} = w \end{cases} \quad \text{and} \quad \tilde{\Xi}_{\mathcal{S}} : \begin{cases} \dot{z} = \tilde{w}^6 \\ \dot{y} = \tilde{w}^3 \end{cases}$$

are not feedback equivalent around $w_0 = 0$, and the reason is that $\tilde{\Xi}_{\mathcal{S}}$ fails to satisfy $\frac{\partial \tilde{F}}{\partial \tilde{w}}(\tilde{w}_0) \neq 0$ at $\tilde{w}_0 = 0$.

According to the previous proposition, in order to deal with the problem of equivalence of submanifolds of the tangent bundle $T\mathcal{X}$ it is interesting, first, to study the problem of equivalence of general control-nonlinear systems Ξ and, second, the problem of equivalence of general control-affine systems Σ .

Conic submanifolds. Recall that a conic, or quadratic, submanifold is given (in a, thus in any, local coordinate system) by $S_q(x, \dot{x}) = 0$, where S_q is a smooth polynomial of degree two in \dot{x} , i.e.

$$S_q(x, \dot{x}) = \dot{x}^t \mathbf{g}(x) \dot{x} + 2\omega(x) \dot{x} + h(x).$$

The map S_q can be represented by the triple $S_q = (\mathbf{g}, \omega, h)$, where \mathbf{g} is a symmetric $(0, 2)$ -tensor (possibly degenerated), ω is a one-form and h is a function. Clearly, two conics S_q of $T\mathcal{X}$ and \tilde{S}_q of $T\tilde{\mathcal{X}}$, given by (\mathbf{g}, ω, h) and $(\tilde{\mathbf{g}}, \tilde{\omega}, \tilde{h})$, respectively, are equivalent if and only if there exists a diffeomorphism $\tilde{x} = \phi(x)$ and a non-vanishing function $\delta = \delta(x)$ on \mathcal{X} such that $\delta \mathbf{g} = \phi^* \tilde{\mathbf{g}}$, $\delta \omega = \phi^* \tilde{\omega}$, and $\delta h = \phi^* \tilde{h}$. In particular, observe that the tensors \mathbf{g} and $\tilde{\mathbf{g}}$, which are (pseudo-)Riemannian metrics (possibly degenerated), are conformally equivalent.

It is well-known in affine geometry that such conic equations can be classified by the signature of the matrix $M_q(x) = \begin{pmatrix} \mathbf{g}(x) & \omega(x)^t \\ \omega(x) & h(x) \end{pmatrix}$ and that of $\mathbf{g}(x)$. We will use the following two determinants

$$\Delta_1(x) = \det(M_q(x)) \quad \text{and} \quad \Delta_2(x) = \det(\mathbf{g}(x)). \quad (1)$$

Of course, Δ_1 and Δ_2 depend on the choice of coordinates, however, since a diffeomorphism $\tilde{x} = \phi(x)$ transforms them as $\Delta_i = \theta^2 \phi^*(\tilde{\Delta}_i)$, for $i = 1, 2$, where $\theta(x) = \det D\phi(x)$, their zero-level sets $\{\Delta_i(x) = 0\}$ are invariantly related to the submanifold \mathcal{S}_q . In this work we will characterise non-degenerate conics, that is, non-empty and satisfying $\Delta_1 \neq 0$; notice that non-empty conics at points of degeneration $\Delta_1(x) = 0$ form in fact (pairs of) linear subspaces of $T_x\mathcal{X}$. Excluding empty \mathcal{S}_q is needed when considering elliptic submanifolds (see lemma below) and it implies that M_q is indefinite. The non-degeneracy assumption $\Delta_1(x) \neq 0$ implies $\frac{\partial S_q}{\partial \dot{x}}(x, \dot{x}) \neq 0$ (but the converse does not hold in general) and that, locally around x_0 , $\text{rk } \mathbf{g}(x) \geq 1$.

If $\Delta_2(x_0) \neq 0$ or if $\Delta_2 \equiv 0$ in a neighbourhood of x_0 , then we can describe three particular types of conics given by the classification lemma below. Notice, however, that this lemma does not describe conics for which we pass smoothly from one type to another (see remark below the proof of [Theorem 3.2](#) for that case).

Lemma 2.1 (Classification of non-degenerate conics) *Consider a nonempty conic \mathcal{S}_q , given by (\mathbf{g}, ω, h) , and assume $\Delta_1(x_0) \neq 0$. Then, locally around x_0 , we have*

- (i) If $\Delta_2(x_0) > 0$, then \mathcal{S}_q is equivalent to $\mathcal{S}_E = \{a^2(\dot{z} - c_0)^2 + b^2(\dot{y} - c_1)^2 = 1\}$,
(ii) If $\Delta_2(x_0) < 0$, then \mathcal{S}_q is equivalent to $\mathcal{S}_H = \{a^2(\dot{z} - c_0)^2 - b^2(\dot{y} - c_1)^2 = 1\}$,
(iii) If $\Delta_2 \equiv 0$, then \mathcal{S}_q is equivalent to $\mathcal{S}_P = \{a\dot{y}^2 - \dot{z} + b\dot{y} + c = 0\}$,

where a, b, c_0, c_1 , and c are smooth functions satisfying $a \neq 0$ and $b \neq 0$ in the elliptic and hyperbolic cases, and $a \neq 0$ in the parabolic case.

We call \mathcal{S}_E , resp. \mathcal{S}_H , resp. \mathcal{S}_P , an elliptic, resp. a hyperbolic, resp. a parabolic, submanifold and we will use the notation \mathcal{S}_Q to denote the set $\{\mathcal{S}_E, \mathcal{S}_H, \mathcal{S}_P\}$ of those three particular forms. Observe that for the parabolic form \mathcal{S}_P , the non-degeneracy assumption $\Delta_1 \neq 0$ implies $\text{rk } \mathbf{g} \equiv 1$ and the existence of a nonvanishing one-form $\omega = -dz + bdy$ satisfying $\omega \notin \text{ann}(\ker \mathbf{g})$, whereas in the elliptic and hyperbolic cases this one-form, given by $\omega = -(c_0 dz \pm c_1 dy)$, can vanish at some points. Those three classes of submanifolds are related to the signature of the metric \mathbf{g} ; indeed if $\text{sgn}(\mathbf{g})$ is constant in a neighbourhood of x_0 , then \mathcal{S}_E , resp. \mathcal{S}_H , resp. \mathcal{S}_P , corresponds to $\text{sgn}(\mathbf{g}) = (+, +)$, resp. $\text{sgn}(\mathbf{g}) = (+, -)$, resp. $\text{sgn}(\mathbf{g}) = (+, 0)$; notice that we can always assume that there is at least one positive eigenvalue, otherwise take the equivalent submanifold given by $\tilde{S} = -S$.

Proof Consider a submanifold \mathcal{S}_q given in local coordinates by $S_q(x, \dot{x}) = 0$, with

$$S_q(x, \dot{x}) = \dot{x}^t \begin{pmatrix} \mathfrak{g}_{11} & \mathfrak{g}_{12} \\ \mathfrak{g}_{12} & \mathfrak{g}_{22} \end{pmatrix} \dot{x} + 2(\omega_1, \omega_2) \dot{x} + h,$$

where all functions \mathfrak{g}_{ij} , ω_1 , ω_2 , and h depend smoothly on $x \in \mathcal{X}$.

- (i)-(ii) We deal with $\Delta_2(x_0) \neq 0$, that is, the elliptic and hyperbolic cases together. In those cases, \mathbf{g} is a non-degenerate symmetric $(0, 2)$ -tensor, therefore it can be interpreted as a pseudo-Riemannian metric. Since on 2-dimensional manifolds all metrics are smoothly diagonalisable (actually, all metrics are conformally flat; see [20, Addendum 1 of Chapter 9] for the Riemannian case and [19, Theorem 7.2] for the Lorentzian one), introduce coordinates $\tilde{x} = \phi(x) = (z, y)$ such that in those coordinates, \mathcal{S}_q can be written (we drop the tildes for more readability),

$$S_q = \lambda_1(x)^2 \dot{z}^2 \pm \lambda_2(x)^2 \dot{y}^2 + 2\omega(x)\dot{x} + h(x)$$

or, equivalently (since $\lambda_1(\cdot) \neq 0$ and $\lambda_2(\cdot) \neq 0$), as

$$S_q = \lambda_1^2 \left(\dot{z} + \frac{\omega_1}{\lambda_1^2} \right)^2 \pm \lambda_2^2 \left(\dot{y} \pm \frac{\omega_2}{\lambda_2^2} \right)^2 + h - \lambda_1^2 \left(\frac{\omega_1}{\lambda_1^2} \right)^2 \mp \lambda_2^2 \left(\frac{\omega_2}{\lambda_2^2} \right)^2$$

Notice that for this form we have $\Delta_1 = \pm \lambda_1^2 \lambda_2^2 \left(h - \frac{(\omega_1)^2}{\lambda_1^2} \mp \frac{(\omega_2)^2}{\lambda_2^2} \right)$ which, by our assumption, does not vanish. Denote $c_0 = -\frac{\omega_1}{\lambda_1^2}$, $c_1 = \mp \frac{\omega_2}{\lambda_2^2}$, and divide by $\tilde{h} = -h + \lambda_1^2 \left(\frac{\omega_1}{\lambda_1^2} \right)^2 \pm \lambda_2^2 \left(\frac{\omega_2}{\lambda_2^2} \right)^2$, observe that $\tilde{h} \neq 0$ as $\tilde{h} = \mp \frac{1}{\lambda_1^2 \lambda_2^2} \Delta_1$, to obtain

$$S_q = \frac{\lambda_1^4 \lambda_2^2}{\mp \Delta_1} (\dot{z} - c_0)^2 \pm \frac{\lambda_1^2 \lambda_2^4}{\mp \Delta_1} (\dot{y} - c_1)^2 - 1,$$

where the upper, resp. lower, sign corresponds to the elliptic, resp. hyperbolic, case. In the elliptic case, if $\Delta_1 > 0$ the conic is empty which is excluded by assumption, therefore $\Delta_1 < 0$ and set $a^2 = \frac{\lambda_1^4 \lambda_2^2}{-\Delta_1}$ and $b^2 = \frac{\lambda_1^2 \lambda_2^4}{-\Delta_1}$ to obtain \mathcal{S}_E . In the hyperbolic case, if $\Delta_1 < 0$, then permute the variables (z, y) and we can thus always obtain a conic defined by S_q in the above form with $\Delta_1 > 0$. Then, set $a^2 = \frac{\lambda_1^4 \lambda_2^2}{\Delta_1}$ and $b^2 = \frac{\lambda_1^2 \lambda_2^4}{\Delta_1}$ to obtain \mathcal{S}_H .

- (iii) Assume $\Delta_2 \equiv 0$. Since $\Delta_1 \neq 0$, we have $\text{rk } \mathbf{g}(x) = 1$ in a neighbourhood of x_0 , which implies that $\mathfrak{g}_{11}(x_0)\mathfrak{g}_{22}(x_0) \neq 0$ and thus, without loss of generality, we can assume that $\mathfrak{g}_{22}(x_0) \neq 0$. Then, by $\text{rk } \mathbf{g}(x) = 1$, the distribution $\ker \mathbf{g} = \text{span} \left\{ \mathfrak{g}_{22}(x) \frac{\partial}{\partial z} - \mathfrak{g}_{12}(x) \frac{\partial}{\partial y} \right\}$ is locally of constant rank and thus we introduce coordinates $\tilde{x} = (\tilde{z}, \tilde{y})$ such that $\ker \mathbf{g} = \text{span} \left\{ \frac{\partial}{\partial \tilde{z}} \right\}$ in which we have

$$\tilde{S}_q = \tilde{a}(\tilde{x}) \dot{\tilde{y}}^2 + 2\tilde{\omega}(\tilde{x}) \dot{\tilde{x}} + \tilde{h}(\tilde{x}),$$

whose determinant $\Delta_1 = -\tilde{a}(\tilde{\omega}_1)^2 \neq 0$ implies that $\tilde{\omega}_1 \neq 0$. Dividing \tilde{S}_q by $-2\tilde{\omega}_1$ we obtain the desired form \mathcal{S}_P with $a = \frac{\tilde{a}}{-2\tilde{\omega}_1}$, $b = \frac{\tilde{\omega}_2}{-\tilde{\omega}_1}$, and $c = \frac{\tilde{h}}{-2\tilde{\omega}_1}$. \square

Example 4. There is a well known example of a control-nonlinear system subject to an elliptic constraint \mathcal{S}_E , namely Dubin's car [7]. The state of the system is the centre of mass of the vehicle $(z, y) \in \mathbb{R}^2$, and we control the orientation of the vehicle (with respect to the z -axis) via $w \in \mathbb{S}^1$. Then, the dynamics of Dubin's car reads

$$\begin{cases} \dot{z} = r \cos(w) \\ \dot{y} = r \sin(w) \end{cases}, \quad r \in \mathbb{R}^*,$$

which clearly is a first extension, regular parametrisation, of the elliptic submanifold $\dot{z}^2 + \dot{y}^2 = r^2$.

We have established all notions necessary for our characterisation and classification of conic submanifolds \mathcal{S}_q by studying the feedback equivalence of their first and second extensions $\Xi_{\mathcal{S}_q}$ and $\Sigma_{\mathcal{S}_q}$.

3 Quadratic control-affine systems

Notations.

$\Sigma = (f, g)$	A control-affine system on a 3-dimensional manifold and with scalar control.
(ϕ, α, β)	Feedback transformations acting on control-affine systems.
$[g, f], \text{ad}_g^k f, L_g()$	Lie bracket, iterated Lie bracket, Lie derivative.
\mathcal{G}	Distribution spanned by the vector field g .
$\Sigma_q = (f_q, g_q)$	Quadratic control-affine system given, in coordinates (x, w) , by the vector fields $f_q = f_q \frac{\partial}{\partial x}$ and $g_q = \frac{\partial}{\partial w}$, which satisfy Definition 3.1 .
(ρ, τ)	Structure functions attached to a control-affine system $\Sigma = (f, g)$, see condition (C2) of Theorem 3.1 .
$\Sigma_Q = \{\Sigma_E, \Sigma_H, \Sigma_P\}$	Set of elliptic, hyperbolic, parabolic subclasses of quadratic systems Σ_q .
Σ_h	Second extension of a submanifold $\mathcal{S} = \{\dot{z} = h(x, \dot{y})\}$.

In this section, we introduce the novel class of quadratic control-affine systems Σ_q that describes second extensions of quadratic submanifolds \mathcal{S}_q given by

$$S_q(x, \dot{x}) = \dot{x}^t g(x) \dot{x} + 2\omega(x) \dot{x} + h(x) = 0$$

and satisfying $\Delta_1 \neq 0$. Next, we address the equivalence problem of a control-affine system Σ to a quadratic control-affine system Σ_q , and in that way, due to [Proposition 2.1](#), we provide a characterisation of quadratic submanifolds \mathcal{S}_q . As a corollary, we will give a characterisation of the elliptic, hyperbolic, and parabolic submanifolds via a characterisation of corresponding subclasses of quadratic control-affine systems. Moreover, by studying our conditions, we will give a normal form of control-affine systems that are feedback equivalent to a quadratic one, and, as a consequence, this will give us a normal form of quadratic submanifolds that smoothly passes from the elliptic to the hyperbolic classes.

On a 3-dimensional manifold \mathcal{M} , equipped with local coordinates ξ , we consider the control-affine system

$$\Sigma : \dot{\xi} = f(\xi) + g(\xi)u,$$

with a scalar control $u \in \mathbb{R}$ and smooth vector fields f and g . A control-affine system Σ is denoted by the pair $\Sigma = (f, g)$, and we set $\mathcal{G} = \text{span}\{g\}$ the distribution spanned by the vector field g . Moreover, we will use the following notations: given two vector fields g and f on \mathcal{M} , by $[g, f]$ we denote the Lie bracket of g and f , in coordinates we have $[g, f] = \frac{\partial f}{\partial \xi} g - \frac{\partial g}{\partial \xi} f$, and $\text{ad}_g^k f = [g, \text{ad}_g^{k-1} f]$ stands for the iterated Lie bracket, with the convention $\text{ad}_g^0 f = f$.

Definition 3.1 (Quadratic and quadratic systems) We say that a control-affine system $\Sigma = (f, g)$ is *quadratic* if it is feedback equivalent to

$$\Sigma_q : \begin{cases} \dot{x} = f_q(x, w) \\ \dot{w} = u \end{cases},$$

where $x \in \mathcal{X}$, a 2-dimensional manifold, $w \in \mathcal{W} \subset \mathbb{R}$, and $f_q(x, w)$ satisfies $\left(\frac{\partial^2 f_q}{\partial w^2} \wedge \frac{\partial f_q}{\partial w}\right)(x, w) \neq 0$, and

$$\frac{\partial^3 f_q}{\partial w^3} = \tau_q(x) \frac{\partial f_q}{\partial w},$$

with τ_q a smooth function of the indicated variable $x \in \mathcal{X}$. A system Σ_q of the above form is called quadratic.

For a quadratic system $\Sigma_q = (f_q, g_q)$, where $f_q = f_q \frac{\partial}{\partial x}$ and $g_q = \frac{\partial}{\partial w}$, it is shown in [Appendix B](#) that any smooth f_q can be written, locally around (x_0, w_0) , as

$$f_q(x, w) = A(x) \sum_{k=0}^{+\infty} \frac{(w-w_0)^{2k+2}}{(2k+2)!} \tau_q(x)^k + B(x) \sum_{k=0}^{+\infty} \frac{(w-w_0)^{2k+1}}{(2k+1)!} \tau_q(x)^k + C(x), \quad (2)$$

where A, B, C are smooth. Since $\mathcal{G} = \text{span}\{g_q\} = \text{span}\left\{\frac{\partial}{\partial w}\right\}$ is involutive and of constant rank one, it is integrable and its integral curves can be identified with the points x of \mathcal{X} and A, B , and C can be seen as smooth vector fields on \mathcal{X} , for which we have $A \wedge B \neq 0$. The following proposition shows that Σ_q is a second extension of a quadratic submanifold \mathcal{S}_q , thus justifies to call Σ_q a quadratic system, and identifies the elliptic, hyperbolic, and parabolic subclasses by describing three normal forms of $f_q = f_q \frac{\partial}{\partial x}$ given for $\tau_q \neq 0$ and $\tau_q \equiv 0$.

Proposition 3.1 *Locally around $\xi_0 \in \mathcal{M}$ we have the followings*

- (i) Σ_q is a second extension of a conic submanifold \mathcal{S}_q and, conversely, any second extension $\Sigma_{\mathcal{S}_q}$ of a conic submanifold \mathcal{S}_q is feedback equivalent to a system of the form Σ_q .
- (ii) If $\tau_q(\xi_0) < 0$ in a neighbourhood, resp. $\tau_q(\xi_0) > 0$, resp. $\tau_q \equiv 0$, then Σ_q is locally feedback equivalent to Σ_E , resp. Σ_H , resp. Σ_P , given by, respectively, f_q of the form

$$\begin{aligned} f_E &= A(x) \cos(\tilde{w}) + B(x) \sin(\tilde{w}) + C(x), & f_H &= A(x) \cosh(\tilde{w}) + B(x) \sinh(\tilde{w}) + C(x), \\ f_P &= A(x)w^2 + B(x)w + C(x), \end{aligned}$$

where in all three cases $A \wedge B \neq 0$.

- (iii) Σ_E , resp. Σ_H , resp. Σ_P , is a second extension of a conic submanifold \mathcal{S}_q satisfying $\Delta_2 > 0$, resp. $\Delta_2 < 0$, resp. $\Delta_2 \equiv 0$.

The conic submanifold \mathcal{S}_q of item (iii) is, by [Lemma 2.1](#), equivalent to \mathcal{S}_E (if $\Delta_2 > 0$), resp. \mathcal{S}_H (if $\Delta_2 < 0$), resp. \mathcal{S}_P (if $\Delta_2 \equiv 0$). So it is natural to call Σ_E an elliptic system, Σ_H a hyperbolic system, and Σ_P a parabolic system. We will denote by Q the set $\{E, H, P\}$, and, consequently, $f_Q = \{f_E, f_H, f_P\}$, $g_Q = \{g_E, g_H, g_P\}$, $f_Q = \{f_E, f_H, f_P\}$, and $\Sigma_Q = \{\Sigma_E, \Sigma_H, \Sigma_P\}$.

Proof

- (i) Consider f_q given by (2), for simplicity of the notations we assume $w_0 = 0$. By [Appendix A](#), we choose coordinates $(z, y) = \phi(x)$ that rectify simultaneously the distributions spanned by A and B so we may assume that $A = a \frac{\partial}{\partial z}$ and $B = b \frac{\partial}{\partial y}$, where a and b are smooth functions satisfying $a(x_0)b(x_0) \neq 0$. We set $C = c_0 \frac{\partial}{\partial z} + c_1 \frac{\partial}{\partial y}$, where c_0 and c_1 are smooth functions of x . Using Cauchy products we compute

$$\begin{aligned} \left(\frac{\dot{z} - c_0}{a}\right)^2 &= \left(\sum_{k=0}^{+\infty} \frac{w^{2k+2}}{(2k+2)!} \tau_q^k\right)^2 = \sum_{k=0}^{+\infty} \frac{(8 \cdot 4^k - 2)w^{2k+4} \tau_q^k}{(2k+4)!}, \\ \left(\frac{\dot{y} - c_1}{b}\right)^2 &= \left(\sum_{k=0}^{+\infty} \frac{w^{2k+1}}{(2k+1)!} \tau_q^k\right)^2 = \sum_{k=0}^{+\infty} \frac{2 \cdot 4^k w^{2k+2} \tau_q^k}{(2k+2)!}, \\ &= w^2 + \sum_{k=1}^{+\infty} \frac{2 \cdot 4^k w^{2k+2} \tau_q^k}{(2k+2)!} = w^2 + \sum_{k=0}^{+\infty} \frac{2 \cdot 4^{k+1} w^{2k+4} \tau_q^{k+1}}{(2k+4)!}, \\ &= w^2 + \tau_q \sum_{k=0}^{+\infty} \frac{8 \cdot 4^k w^{2k+4} \tau_q^k}{(2k+4)!} = w^2 + \tau_q \left(\frac{\dot{z} - c_0}{a}\right)^2 + \tau_q \sum_{k=0}^{+\infty} \frac{2w^{2k+4} \tau_q^k}{(2k+4)!}, \\ \left(\frac{\dot{y} - c_1}{b}\right)^2 - \tau_q \left(\frac{\dot{z} - c_0}{a}\right)^2 &= w^2 + \sum_{k=0}^{+\infty} \frac{2w^{2k+4} \tau_q^{k+1}}{(2k+4)!} = \sum_{k=-1}^{+\infty} \frac{2w^{2k+4} \tau_q^{k+1}}{(2k+4)!}, \\ &= \sum_{k=0}^{+\infty} \frac{2w^{2k+2} \tau_q^k}{(2k+2)!} = 2 \left(\frac{\dot{z} - c_0}{a}\right). \end{aligned}$$

Hence, Σ_q is a second extension of the submanifold \mathcal{S}_q given by

$$\left(\frac{\dot{y} - c_1}{b}\right)^2 - \tau_q \left(\frac{\dot{z} - c_0}{a}\right)^2 - 2 \left(\frac{\dot{z} - c_0}{a}\right) = 0, \quad (3)$$

and for which we have $\Delta_1 = -\frac{1}{a^2 b^2}$ and $\Delta_2 = -\frac{\tau_q}{a^2 b^2}$.

To prove that any second extension of a conic submanifold S_q is feedback equivalent to a system of the form Σ_q , we show that any conic submanifold is equivalent via a diffeomorphism to a conic submanifold of the form given by (3). To this end, consider $S_q = \dot{x}^t \mathbf{g} \dot{x} + 2\omega \dot{x} + h = a\dot{z}^2 + 2b\dot{z}\dot{y} + c\dot{y}^2 + 2d\dot{z} + 2e\dot{y} + h = 0$, where all functions a, b, c, d, e, h depend smoothly on $x = (z, y)$. By $\Delta_1(x_0) \neq 0$, we have $\text{rk } \mathbf{g}(x_0) \geq 1$, where $\mathbf{g} = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$. If $\text{rk } \mathbf{g}(x_0) = 2$, then we can apply the results of Lemma 2.1 to get S_q in the form of (3). If $\text{rk } \mathbf{g}(x_0) = 1$, then we may assume that $c(x_0) \neq 0$. Indeed, if $a(x_0) = c(x_0) = 0$, then $b(x_0) \neq 0$ implying that $\text{rk } \mathbf{g}(x_0) = 2$ so either $a(x_0) \neq 0$ or $c(x_0) \neq 0$ and we can always suppose $c(x_0) \neq 0$ by permuting y and z , if necessary. Dividing $S_q = 0$ by c we obtain (we keep the same names for all remaining functions):

$$S_q = a\dot{z}^2 + 2b\dot{z}\dot{y} + \dot{y}^2 + 2d\dot{z} + 2e\dot{y} + h = (a - b^2)\dot{z}^2 + (b\dot{z} + \dot{y})^2 + 2d\dot{z} + 2e\dot{y} + h = 0.$$

Choose local coordinates $(\tilde{z}, \tilde{y}) = (z, \psi(z, y))$, hence $\frac{\partial \psi}{\partial y} \neq 0$, that rectify the line-distribution $\text{span} \left\{ \frac{\partial}{\partial z} - b \frac{\partial}{\partial y} \right\}$, i.e. $\frac{\partial \psi}{\partial z} - b \frac{\partial \psi}{\partial y} = 0$. Therefore, $\dot{\tilde{z}} = \dot{z}$ and $\dot{\tilde{y}} = \frac{\partial \psi}{\partial z} \dot{z} + \frac{\partial \psi}{\partial y} \dot{y} = \frac{\partial \psi}{\partial y} (b\dot{z} + \dot{y})$. Thus \tilde{S}_q , which is S_q in (\tilde{z}, \tilde{y}) -coordinates, reads

$$\tilde{S}_q = \frac{1}{\tilde{b}^2} \dot{\tilde{y}}^2 + \tilde{a} \dot{\tilde{z}}^2 + 2\tilde{d} \dot{\tilde{z}} + 2\tilde{e} \dot{\tilde{y}} + \tilde{h} = 0,$$

where $\tilde{b} = \frac{\partial \psi}{\partial y}$ and $\tilde{a}(\tilde{x}_0) = 0$ because $\text{rk } \tilde{\mathbf{g}}(\tilde{x}_0) = 1$. By $\tilde{\Delta}_1(\tilde{x}_0) \neq 0$ and $\tilde{a}(\tilde{x}_0) = 0$, we conclude that $\tilde{d}(\tilde{x}_0) \neq 0$ and thus dividing by \tilde{d} (and removing the "tildes" from the coordinates) we get

$$\bar{S}_q = \frac{\tilde{S}_q}{\tilde{d}} = \frac{1}{\bar{b}^2} \dot{y}^2 + \bar{A} \dot{z}^2 + 2\dot{z} + 2\bar{e} \dot{y} + \bar{h} = \left(\frac{\dot{y} - \bar{c}_1}{\bar{b}} \right)^2 + \bar{A} \dot{z}^2 + 2\dot{z} + \bar{H} = 0,$$

with $\bar{b} = \tilde{b} \sqrt{\tilde{d}}$ (if $\tilde{d} < 0$, then we take $z = -\tilde{z}$ to get $\tilde{d} > 0$), $\bar{A} = \frac{\tilde{a}}{\tilde{d}}$ (satisfying $\bar{A}(x_0) = 0$), $\bar{c}_1 = -\bar{e} \bar{b}^2$, and $\bar{H} = \tilde{h} - \bar{e}^2 \bar{b}^2$ (see also [4] for another proof for the smooth diagonalisation of a symmetric $(0, 2)$ -tensor in the case of rank deficiency). To get the form (3), we will now prove that there exists, locally around x_0 , functions $\bar{\tau}_q, \bar{a} \neq 0$, and \bar{c}_0 such that $\bar{A} \dot{z}^2 + 2\dot{z} + \bar{H} = -\bar{\tau}_q \left(\frac{\dot{z} - \bar{c}_0}{\bar{a}} \right)^2 - 2 \left(\frac{\dot{z} - \bar{c}_0}{\bar{a}} \right)$. We obtain

$$\frac{-\bar{\tau}_q}{\bar{a}^2} = \bar{A}, \quad \frac{\bar{\tau}_q \bar{c}_0}{\bar{a}^2} - \frac{1}{\bar{a}} = 1, \quad \text{and} \quad -\frac{\bar{\tau}_q (\bar{c}_0)^2}{\bar{a}^2} + 2 \frac{\bar{c}_0}{\bar{a}} = \bar{H}.$$

Hence, from the second equation, $\frac{\bar{c}_0}{\bar{a}} = -\bar{A}(\bar{c}_0)^2 - \bar{c}_0$ implying that $\bar{A}(\bar{c}_0)^2 + 2\bar{c}_0 + \bar{H} = 0$. The latter equation possesses smooth solutions $\bar{c}_0(z, y)$ because the discriminant (evaluated at x_0) of that degree two polynomial is $\bar{\Delta}(x_0) = 4 > 0$ (recall that $\bar{A}(x_0) = 0$). So

$$\bar{S}_q = \left(\frac{\dot{y} - \bar{c}_1}{\bar{b}} \right)^2 - \bar{\tau}_q \left(\frac{\dot{z} - \bar{c}_0}{\bar{a}} \right)^2 - 2 \left(\frac{\dot{z} - \bar{c}_0}{\bar{a}} \right) = 0,$$

where $\bar{a} = \frac{-1}{1 + \bar{A} \bar{c}_0}$, which is well defined since $\bar{A}(x_0) = 0$, and $\bar{\tau}_q = -\bar{A} \bar{a}^2$, proving that \bar{S}_q is of the desired form (3).

- (ii) If $\tau_q < 0$, then $\frac{\partial^3 f_q}{\partial w^3} = \tau_q \frac{\partial f_q}{\partial w}$ implies that $f_q = A \cos(\sqrt{-\tau_q} w) + B \sin(\sqrt{-\tau_q} w) + C$, where A, B , and C depend on x and, clearly, $\frac{\partial^2 f_q}{\partial w^2} \wedge \frac{\partial f_q}{\partial w} \neq 0$ implies $A \wedge B \neq 0$ (for this case, as well as for the next two cases). Via the change of coordinate $\tilde{w} = \sqrt{-\tau_q} w$, we get $f_q = f_E$. If $\tau_q > 0$, then $\frac{\partial^3 f_q}{\partial w^3} = \tau_q \frac{\partial f_q}{\partial w}$ implies that $f_q = A \cosh(\sqrt{\tau_q} w) + B \sinh(\sqrt{\tau_q} w) + C$ which, via the change of coordinate $\tilde{w} = \sqrt{\tau_q} w$, gives $f_q = f_H$. Finally, if $\tau_q \equiv 0$, then $\frac{\partial^3 f_q}{\partial w^3} \equiv 0$ implies that $f_q = f_P = Aw^2 + Bw + C$.
- (iii) By item (i), Σ_q is a second extension of a conic submanifold S_q for which we have $\Delta_2 = -\frac{\tau_q}{a^2 b^2}$. Thus, $\Delta_2 > 0$, resp. $\Delta_2 < 0$, resp. $\Delta_2 \equiv 0$ if and only if $\tau_q < 0$, resp. $\tau_q > 0$, resp. $\tau_q \equiv 0$, which, by item (ii), correspond to Σ_E , resp. Σ_H , resp. Σ_P . □

Notice that τ_q plays for Σ_q an analogous role to that played by Δ_2 for S_q ; indeed, we have $\text{sgn}(\Delta_2(x)) = -\text{sgn}(\tau_q(x))$. In particular, the sign of τ_q identifies the subclasses of elliptic, hyperbolic, and parabolic control-affine systems. Moreover, statement (i) shows that every second extension of a quadratic submanifold S_q is feedback equivalent to a quadratic system Σ_q and the other way around, therefore to obtain a characterisation of quadratic submanifolds, it is crucial to characterise the class of quadratizable systems.

The remaining part of this section is organised as follows. First, we will state our main theorem giving necessary and sufficient conditions characterising the class of quadratic control-affine systems Σ_q . Second, by carefully studying the conditions of that theorem, we will give a normal form of the quadratizable systems.

3.1 Characterisation of quadratzable control-affine systems

We now focus on the feedback equivalence of a general control-affine system $\Sigma : \dot{\xi} = f(\xi) + g(\xi)u$ with a quadratic control-affine system of the form Σ_q . The theorem below gives checkable necessary and sufficient conditions in terms of the vector fields f and g of Σ for the existence of a smooth feedback transformation (ϕ, α, β) that locally brings Σ into a quadratic system Σ_q . Equivalence to particular subcases of Σ_q , namely elliptic Σ_E , hyperbolic Σ_H , and parabolic Σ_P , is provided by [Corollary 3.1](#) below.

Theorem 3.1 (Feedback quadratisation) *Let $\Sigma = (f, g)$ be a control-affine system on a 3-dimensional smooth manifold with a scalar control. The system Σ is, locally around $\xi_0 \in \mathcal{M}$, feedback equivalent to a quadratic system Σ_q if and only if*

- (C1) $g \wedge \text{ad}_g f \wedge \text{ad}_g^2 f(\xi_0) \neq 0$,
- (C2) The structure functions ρ and τ in the decomposition $\text{ad}_g^3 f = \rho \text{ad}_g^2 f + \tau \text{ad}_g f \pmod{\mathcal{G}}$ satisfy, locally around ξ_0 ,

$$\mathbf{L}_g(\chi) - \frac{2}{3}\rho\chi = 0, \quad (4)$$

where $\chi = 3\mathbf{L}_g(\rho) - 2\rho^2 - 9\tau$.

Condition (C1) is a regularity condition, it ensures that the vector fields $g, \text{ad}_g f$, and $\text{ad}_g^2 f$ are locally linearly independent and thus that they form a local frame, hence the structure functions (ρ, τ) of (C2) are well defined. The main idea behind this theorem is to observe that if for Σ we have $\text{ad}_g^3 f = \tau(x) \text{ad}_g f$, modulo $\mathcal{G} = \text{span}\{g\}$, i.e. the third Lie derivative of f along g is proportional to the first Lie derivative of f along g , modulo \mathcal{G} , then with the help of a diffeomorphism we can obtain the form Σ_q , see the sufficiency part of the proof for details. Thus condition (C2) shows how that relation changes when we allow for feedback transformations (α, β) .

Proof Necessity. Consider the affine control system Σ given by two smooth vector fields f and g and recall that \mathcal{G} is the distribution $\mathcal{G} = \text{span}\{g\}$. Let (α, β) form a control-affine feedback and let ϕ be a diffeomorphism such that Σ is, locally, transformed into Σ_q via ϕ and (α, β) . In coordinates $\tilde{\xi} = \phi(\xi) = (\tilde{x}, \tilde{w}) = (\tilde{z}, \tilde{y}, \tilde{w})$, we denote $\tilde{f}_q = \tilde{f}_q \frac{\partial}{\partial \tilde{x}}$ and $\tilde{g}_q = \frac{\partial}{\partial \tilde{w}}$ the vector fields of Σ_q , $\tilde{\mathcal{G}}$ the distribution spanned by \tilde{g}_q , and $(\tilde{\rho}, \tilde{\tau})$ the structure functions of Σ_q , defined as in (C2). By definition of feedback equivalence the following relations between (f, g) and $(\tilde{f}_q, \tilde{g}_q)$ hold: $\tilde{f}_q = \phi_*(f + \alpha g)$ and $\tilde{g}_q = \phi_*(g\beta)$.

The system Σ_q is quadratic, so by [Definition 3.1](#), we have $\frac{\partial^2 \tilde{f}_q}{\partial \tilde{w}^2} \wedge \frac{\partial \tilde{f}_q}{\partial \tilde{w}}(\tilde{x}_0, \tilde{w}_0) \neq 0$, which implies that (C1) holds for Σ_q , and we also have $\frac{\partial^3 \tilde{f}_q}{\partial \tilde{w}^3} = \tilde{\tau}_q \frac{\partial \tilde{f}_q}{\partial \tilde{w}}$, thus we get $\tilde{\rho} = 0$ and $\tilde{\tau} = \tilde{\tau}_q(\tilde{z}, \tilde{y})$. Therefore for Σ_q we have $\tilde{\chi}(\tilde{z}, \tilde{y}) = -9\tilde{\tau}(\tilde{z}, \tilde{y})$ implying $\mathbf{L}_{\tilde{g}_q}(\tilde{\chi}) - \frac{2}{3}\tilde{\rho}\tilde{\chi} = \frac{\partial \tilde{\chi}}{\partial \tilde{w}} = 0$. Hence Σ_q satisfies (C1) and (C2) and we will now prove that those conditions are invariant under diffeomorphisms ϕ and feedback transformations (α, β) . Clearly, (C1) is invariant under diffeomorphisms (as $[\phi_* f, \phi_* g] = \phi_* [f, g]$) and under feedback (α, β) since $\beta \neq 0$. We have checked that (C2) holds for $\Sigma_q = (\tilde{f}_q, \tilde{g}_q)$ and, clearly, (C2) is invariant under diffeomorphisms since they conjugate structure functions. Moreover (C2) is invariant under the transformation $\tilde{f}_q \mapsto \tilde{f}_q + \alpha \tilde{g}_q$, since the expression of $\text{ad}_{\tilde{g}_q}^3 \tilde{f}_q$ is considered modulo the distribution $\tilde{\mathcal{G}}$. Finally, under the action of β the brackets, with $\tilde{g}_q = g\beta$, are transformed by

$$\begin{aligned} \text{ad}_{\tilde{g}_q} f_q &= \beta \text{ad}_g f \pmod{\tilde{\mathcal{G}}}, \\ \text{ad}_{\tilde{g}_q}^2 f_q &= \beta^2 \text{ad}_g^2 f + \beta \mathbf{L}_g(\beta) \text{ad}_g f \pmod{\tilde{\mathcal{G}}}, \\ \text{ad}_{\tilde{g}_q}^3 f_q &= (\beta^3 \rho + 3\beta^2 \mathbf{L}_g(\beta)) \text{ad}_g^2 f + (\beta^3 \tau + \beta \mathbf{L}_g(\beta \mathbf{L}_g(\beta))) \text{ad}_g f \pmod{\tilde{\mathcal{G}}}, \\ &= (\rho\beta + 3\mathbf{L}_g(\beta)) \text{ad}_{\tilde{g}_q}^2 f + \left(\tau\beta^2 + \mathbf{L}_g(\beta \mathbf{L}_g(\beta)) - \rho\beta \mathbf{L}_g(\beta) - 3(\mathbf{L}_g(\beta))^2 \right) \text{ad}_{\tilde{g}_q} f \pmod{\tilde{\mathcal{G}}} \end{aligned}$$

This implies that the structure functions $\tilde{\rho}$ and $\tilde{\tau}$ of Σ_q defined by $\text{ad}_{\tilde{g}_q}^3 f_q = \tilde{\rho} \text{ad}_{\tilde{g}_q}^2 f_q + \tilde{\tau} \text{ad}_{\tilde{g}_q} f_q \pmod{\tilde{\mathcal{G}}}$ are given in terms of the feedback transformation β and the structure functions ρ and τ of Σ by

$$\tilde{\rho} = \rho\beta + 3\mathbf{L}_g(\beta), \quad \tilde{\tau} = \tau\beta^2 + \mathbf{L}_g(\beta \mathbf{L}_g(\beta)) - \rho\beta \mathbf{L}_g(\beta) - 3(\mathbf{L}_g(\beta))^2. \quad (5)$$

Since for Σ_q the structure function $\tilde{\rho} = 0$, we have the relation $\mathbf{L}_g(\beta) = -\frac{\beta\rho}{3}$ and thus $\tilde{\chi} = -9\tilde{\tau}$, which is equal to

$$\begin{aligned} \tilde{\chi} &= -9 \left(\tau\beta^2 + \mathbf{L}_g(\beta \mathbf{L}_g(\beta)) \right) = -9 \left(\tau\beta^2 + \mathbf{L}_g \left(\frac{-\beta^2 \rho}{3} \right) \right), \\ &= -9 \left(\tau\beta^2 - \frac{1}{3} \left(\rho \mathbf{L}_g(\beta^2) + \beta^2 \mathbf{L}_g(\rho) \right) \right) = -9\beta^2 \left(\tau - \frac{1}{3} \mathbf{L}_g(\rho) + \frac{2}{9} \rho^2 \right) = \beta^2 \chi. \end{aligned}$$

And finally,

$$L_{\tilde{g}_q}(\tilde{\chi}) = \beta L_g(\beta^2 \chi) = \beta^3 L_g(\chi) + 2\beta^2 \chi L_g(\beta) = \beta^3 L_g(\chi) - \frac{2}{3} \beta^3 \chi \rho = 0,$$

showing the necessity of relation (4) and concludes the necessity part of the proof.

Sufficiency. There are two steps in the sufficiency part. The first one consists in building a vector field g such that $\text{ad}_g^3 f = \tau \text{ad}_g f \pmod{\mathcal{G}}$ with $L_g(\tau) = 0$. Then we will construct a diffeomorphism ϕ that brings Σ into the form Σ_q .

Consider the system $\Sigma : \dot{\xi} = f + gu$, for which we assume $g \wedge \text{ad}_g f \wedge \text{ad}_g^2 f(\xi_0) \neq 0$ and suppose that relation (4) holds for the structure functions ρ and τ of Σ . Choose a function $\beta \neq 0$ satisfying $L_g(\beta) = -\frac{\beta \rho}{3}$, which exists locally since $g \neq 0$ by condition (C1); to guarantee that $\beta \neq 0$, we actually may solve the equation $L_g(\ln(\beta)) = -\frac{\rho}{3}$. Define the system $\tilde{\Sigma} : \dot{\xi} = \tilde{f} + \tilde{g}u$, where $\tilde{g} = g\beta$ and $\tilde{f} = f$, then by (5) the structure function $\tilde{\rho}$ of $\tilde{\Sigma}$ vanishes. Therefore, we have $\tilde{\chi} = -9\tilde{\tau}$ and thus relation (4) implies that $L_{\tilde{g}}(\tilde{\tau}) = 0$.

Since $\tilde{g} \neq 0$, we apply a diffeomorphism $(z, y, w) = \phi(\xi)$ such that $\phi_* \tilde{g} = g_q = \frac{\partial}{\partial w}$ and denote $f_q = \phi_* \tilde{f}$, and $\tau_q = \tilde{\tau} \circ \phi$. Therefore, the decomposition $\text{ad}_{g_q}^3 f_q = \tau_q \text{ad}_{g_q} f_q \pmod{\mathcal{G}}$ implies that $f_q = f_q^1 \frac{\partial}{\partial z} + f_q^2 \frac{\partial}{\partial y} + f_q^3 \frac{\partial}{\partial w}$ satisfies

$$\frac{\partial^3 f_q^i}{\partial w^3} = \tau(z, y) \frac{\partial f_q^i}{\partial w}, \quad (6)$$

for $i = 1, 2$. Applying the feedback $u = f_q^3(z, y, w) + \tilde{u}$ we obtain the form Σ_q with $f_q = f_q^1 \frac{\partial}{\partial z} + f_q^2 \frac{\partial}{\partial y}$ and $g_q = \frac{\partial}{\partial w}$. The condition $\frac{\partial^2 f_q}{\partial w^2} \wedge \frac{\partial f_q}{\partial w} \neq 0$ follows from (C1) and feedback invariance of the latter. \square

The following corollary shows that we can test on the structure functions of Σ if the equivalent quadratic system Σ_q will be of elliptic, hyperbolic, or parabolic type.

Corollary 3.1 *Under conditions (C1) and (C2) of the previous theorem we have, locally around ξ_0 ,*

- (i) Σ is feedback equivalent to Σ_E if and only if $\chi(\xi_0) > 0$,
- (ii) Σ is feedback equivalent to Σ_H if and only if $\chi(\xi_0) < 0$,
- (iii) Σ is feedback equivalent to Σ_P if and only if $\chi \equiv 0$ in a neighbourhood of ξ_0 ,

where $\chi = 3L_g(\rho) - 2\rho^2 - 9\tau$.

Notice that Σ is locally feedback equivalent to Σ_P if and only if it satisfies (C1) and $\chi \equiv 0$, condition (C2) being satisfied automatically.

Proof From the necessity part of the proof of Theorem 3.1 we know that for Σ_q , with structure functions $\rho = 0$ and $\tau = \tau_q$, we have $\chi = -9\tau_q$ and we observed that under pure feedback transformations (α, β) we have $\tilde{\chi} = \beta^2 \chi$, thus the sign of χ is invariant as well as the locus where it vanishes. Moreover, by statement (ii) of Proposition 3.1, Σ_q is elliptic if $\tau_q > 0$, equivalently $\chi < 0$, Σ_q is hyperbolic if $\tau_q < 0$, equivalently $\chi > 0$, and Σ_q is parabolic if $\tau_q \equiv 0$, equivalently $\chi \equiv 0$. Hence the necessity of the stated conditions is established.

Conversely, in the sufficiency part of the proof of Theorem 3.1 we obtained Σ with structure functions $(\tilde{\rho}, \tilde{\tau}) = (0, \tilde{\tau}(z, y))$ via a suitable feedback transformations. Since $\tilde{\chi} = \beta^2 \chi$, we have $-9\tilde{\tau} = \beta^2 \chi$ and thus we get $\text{sgn}(\tilde{\tau}) = -\text{sgn}(\chi)$ and the conclusion follows by statement (ii) of Proposition 3.1. \square

3.2 Normal form of quadratizable control-affine system

Any control-affine system Σ , under the regularity assumption $g \wedge \text{ad}_g f(\xi_0) \neq 0$, can be written (after applying a suitable feedback transformation) locally around $0 \in \mathbb{R}^3$ as

$$\Sigma_h : \begin{cases} \dot{z} = h(z, y, w) \\ \dot{y} = w + \varepsilon \\ \dot{w} = u \end{cases},$$

with h a smooth function and $\varepsilon = 0, 1$. The parameter ε , which is invariant under local feedback transformations, is $\varepsilon = 0$ if either $(f \wedge g \wedge \text{ad}_g f)(\xi_0) \neq 0$ or $(f \wedge g)(\xi_0) = 0$ and $\varepsilon = 1$ otherwise, i.e. $(f \wedge g)(\xi_0) \neq 0$ but $(f \wedge g \wedge \text{ad}_g f)(\xi_0) = 0$. By applying Theorem 3.1, we will give in this subsection a normal form of all smooth functions $h(z, y, w)$ that describe quadratizable systems, that is, control-affine systems feedback equivalent to Σ_q . In what follows, we assume to work locally around $0 \in \mathbb{R}^3$ and all derivatives are taken with respect to w and denoted by prime, double prime etc. Whenever we apply $\ln(a)$, we assume that $a > 0$ (if not, take the absolute value).

Theorem 3.2 (Normal form of quadratzable control-affine systems) *The following statements are equivalent, locally around $0 \in \mathbb{R}^3$:*

- (i) Σ_h is feedback equivalent to a quadratic system Σ_q ;
- (ii) The function h satisfies $h''(0) \neq 0$ and, in a neighbourhood, it holds

$$9h^{(5)}(h'')^2 - 45h^{(4)}h^{(3)}h'' + 40(h^{(3)})^3 = 0, \quad (7)$$

recall that the derivatives are taken with respect to w ;

- (iii) The second derivative of h is of the following form

$$h''(x, w) = a(dw^2 + ew + 1)^{-3/2}, \quad (8)$$

where $a = a(x)$, $d = d(x)$, and $e = e(x)$ are smooth functions satisfying $a(0) \neq 0$;

- (iv) The function h is given by

$$h(x, w) = 2a \left(\frac{w^2}{(\sqrt{dw^2 + ew + 1} + 1)^2 - dw^2} \right) + bw + c \quad (9)$$

where a, b, c, d, e are any smooth functions of x such that $a(0) \neq 0$.

Proof (i) \Rightarrow (ii). It is a straightforward application of the conditions of [Theorem 3.1](#) with the structure functions of Σ_h given by $\rho = \frac{h^{(3)}}{h''}$ and $\tau = 0$ yielding $\chi = 3\rho' - 2\rho^2$. By [\(C1\)](#), we have $h''(0) \neq 0$ and then condition [\(C2\)](#) reads

$$\chi' - \frac{2}{3}\rho\chi = 3\rho'' - 6\rho\rho' + \frac{4}{3}\rho^3 = 0, \quad (10)$$

which, by plugging $\rho = \frac{h^{(3)}}{h''}$ into the last equation, gives [\(7\)](#).

(ii) \Rightarrow (iii). Assume that h satisfies $h''(0) \neq 0$ and [\(7\)](#). Set $\rho = \frac{h^{(3)}}{h''}$, then $\rho = \rho(x, w)$ fulfils $3\rho'' - 6\rho\rho' + \frac{4}{3}\rho^3 = 0$, namely, the second equation of [\(10\)](#). By a change of variable, it is easy to obtain that the solutions of [\(10\)](#) are of the following form (see [Appendix C.1](#) for the proof)

$$\rho(x, w) = -\frac{3}{2} \frac{2d(x)w + e(x)}{d(x)w^2 + e(x)w + 1}. \quad (11)$$

This form can be integrated, using $\rho = \frac{h^{(3)}}{h''} = (\ln h'')$, into $h''(x, w) = a(x) (d(x)w^2 + e(x)w + 1)^{-3/2}$ with a, d , and e any smooth functions such that $a(0) \neq 0$.

(iii) \Rightarrow (iv). To show [\(9\)](#), we integrate twice the second derivative of h given by [\(8\)](#). Denote $p = p(x, w) = d(x)w^2 + e(x)w + 1$ and $\Delta = \Delta(x) = e(x)^2 - 4d(x)$. First, we obtain (see [Appendix C.2](#) for details)

$$h'(x, w) = \frac{2aw(\sqrt{p} + 1)}{\sqrt{p}(ew + 2 + 2\sqrt{p})} + b,$$

with b an arbitrary smooth function of x . Integrate once more to get

$$\begin{aligned} h(x, w) &= \frac{2a}{\Delta\sqrt{p}} (ew\sqrt{p} - 2p) + \frac{4a}{\Delta} + bw + c = \frac{2a}{\Delta} (ew + 2 - 2\sqrt{p}) + bw + c, \\ &= \frac{2aw^2}{ew + 2 + 2\sqrt{p}} + bw + c = \frac{2aw^2}{(\sqrt{p} + 1)^2 - dw^2} + bw + c. \end{aligned}$$

(iv) \Rightarrow (i). Given Σ_h with h defined by [\(9\)](#) we will construct a feedback transformation that brings the system into Σ_q . First, we introduce coordinates, centred at $0 \in \mathbb{R}^2$, $(\tilde{z}, \tilde{y}) = \phi(z, y)$, where $\tilde{y} = y$, such that $\phi_* \left(b \frac{\partial}{\partial z} + \frac{\partial}{\partial y} \right) = \frac{\partial}{\partial \tilde{y}}$. Those coordinates transform the system Σ_h into

$$\begin{cases} \dot{\tilde{z}} = 2\tilde{a} \frac{w^2}{(\sqrt{p} + 1)^2 - \tilde{d}w^2} + \tilde{c} \\ \dot{\tilde{y}} = w + \varepsilon \\ \dot{w} = u \end{cases},$$

where $\tilde{a}, \tilde{c}, \tilde{d}$, and \tilde{p} are new functions satisfying $\tilde{a}(0) \neq 0$ and $d = \tilde{d} \circ \phi$ and $p = \tilde{p} \circ \phi$. Second, we set $\tilde{w}^2 = \frac{w^2}{(\sqrt{p} + 1)^2 - \tilde{d}w^2}$ or, equivalently, $w = \tilde{w} \left(\tilde{e}\tilde{w} \pm 2\sqrt{\tilde{d}\tilde{w}^2 + 1} \right)$, which brings the above system into (after applying a suitable feedback along the last component)

$$\tilde{\Sigma}_h : \begin{cases} \dot{\tilde{z}} = 2\tilde{a}\tilde{w}^2 + \tilde{c} \\ \dot{\tilde{y}} = \tilde{e}\tilde{w}^2 \pm 2\tilde{w}\sqrt{\tilde{d}\tilde{w}^2 + 1} + \varepsilon \\ \dot{\tilde{w}} = \tilde{u} \end{cases}. \quad (12)$$

The structure functions of $\tilde{\Sigma}_h$ are given by $\tilde{\rho} = \frac{-3\tilde{w}\tilde{d}}{d\tilde{w}^2+1}$ and $\tilde{\tau} = \frac{3\tilde{d}}{d\tilde{w}^2+1}$, then apply the feedback $\tilde{u} = \beta\tilde{w}$, where $\beta(\tilde{x}, \tilde{w}) = \sqrt{d\tilde{w}^2+1}$ (it is a solution of the equation $\frac{\partial\beta}{\partial\tilde{w}} = -\frac{\tilde{\rho}\beta}{3}$), to obtain a new vector field $\bar{g} = \beta\frac{\partial}{\partial\tilde{w}}$ and structure functions of (\bar{f}, \bar{g}) , where \bar{f} is the drift of $\tilde{\Sigma}_h$, are given by $\bar{\rho} = 0$ and $\bar{\tau} = 4\tilde{d}(x)$. To complete the form, it remains to find a new $\bar{w} = \psi(\tilde{x}, \tilde{w})$ such that the diffeomorphism $(\tilde{x}, \tilde{w}) = \Psi(\tilde{x}, \tilde{w}) = (\tilde{x}, \psi(\tilde{x}, \tilde{w}))$ satisfies $\Psi_*\bar{g} = \Psi_*\beta\frac{\partial}{\partial\tilde{w}} = \frac{\partial}{\partial\bar{w}}$. In general, ψ is given in terms of the following integral

$$\bar{w} = \psi(\tilde{x}, \tilde{w}) = \int_0^{\tilde{w}} \frac{1}{\sqrt{1+d\tilde{w}^2}} d\tilde{w}$$

and, together with a suitable feedback along $\dot{\bar{w}}$, thus provides a quadratic system Σ_q . \square

Remark 3.1 This theorem provides a normal form of submanifolds $\mathcal{S} = \{\dot{z} = s(x, \dot{y})\}$ that are equivalent to a conic submanifold \mathcal{S}_q , namely, they are represented by $s(x, \dot{y}) = h(x, \dot{y})$ with h as in (9). Moreover, system (12) leads to a normal form for all conic submanifold \mathcal{S}_q (even for those that smoothly pass through $\Delta_2(x) = 0$, i.e. from the elliptic to the hyperbolic submanifolds) and thus completes the characterisation of Lemma 2.1. Indeed, for (12) we have (tildes have been removed for more readability)

$$S_q(x, \dot{x}) = [e(\dot{z} - c) - 2a(\dot{y} - \varepsilon)]^2 - 4d(\dot{z} - c)^2 - 8a(\dot{z} - c),$$

for which $a \neq 0$ and we can compute $\Delta_1 = -64a^4 \neq 0$ and $\Delta_2 = -16a^2d$.

In the last item of the proof we saw that the function d plays an import role for the shape of the transformation, in the following corollary we show that this function is the key of the normal form of quadratzable control systems.

Corollary 3.2 *Assume that Σ_h is given by h of the form (9). Then Σ is feedback equivalent to Σ_P , resp. Σ_E , resp. Σ_H if and only if $d \equiv 0$, resp. $d < 0$, resp. $d > 0$. Moreover, the normalizing feedback transformation is given by*

$$\begin{aligned} \bar{w} &= \frac{w}{1 + \sqrt{ew + 1}} \quad \text{for } \Sigma_P, \\ \text{resp. } \sin^2(\sqrt{-d}\bar{w}) &= \frac{-dw^2}{ew + 2 + 2\sqrt{p}} \quad \text{for } \Sigma_E, \quad \text{resp. } \sinh^2(\sqrt{d}\bar{w}) = \frac{dw^2}{ew + 2 + 2\sqrt{p}} \quad \text{for } \Sigma_H, \end{aligned}$$

where $p = p(x, w) = d(x)w^2 + e(x)w + 1$.

Proof First we show that Σ_h is feedback equivalent to Σ_P , resp. Σ_E , resp. Σ_H , if and only if $d \equiv 0$, resp. $d < 0$, and resp. $d > 0$. From Corollary 3.1 we know that we have to compute the sign of χ , which is given by $\chi = -9\frac{d}{p}$ for Σ_h (this can easily be deduced from the expression of ρ given by (11)). Since $p(0) = 1 > 0$ we have $\text{sgn}(\chi) = -\text{sgn}(d)$ and thus the conclusion follows.

We now show how to explicitly transform Σ_h into Σ_P , resp. Σ_E , resp. Σ_H . From the last part of the proof of the previous theorem we know that a suitable parametrisation \bar{w} is given by the following two steps

$$\tilde{w}^2 = \frac{w^2}{ew + 2 + 2\sqrt{p}}, \quad \text{and} \quad \bar{w} = \int_0^{\tilde{w}} \frac{1}{\sqrt{1+d\tilde{w}^2}} d\tilde{w}.$$

Assume $d \equiv 0$, then the procedure reduces to the first step only and thus $\bar{w}^2 = \tilde{w}^2 = \frac{w^2}{ew+2+2\sqrt{ew+1}} = \left(\frac{w}{1+\sqrt{ew+1}}\right)^2$ and we choose $\tilde{w} = \frac{w}{1+\sqrt{ew+1}}$. Assume $d < 0$, then the second step of the procedure leads to $\bar{w} = \frac{1}{\sqrt{-d}} \arcsin(\sqrt{-d}\tilde{w})$. Hence a reparametrisation is given by $\sin^2(\sqrt{-d}\bar{w}) = \frac{-dw^2}{ew+2+2\sqrt{p}}$. Assume $d > 0$, then from the second step of the procedure we have $\bar{w} = \frac{1}{\sqrt{d}} \operatorname{arcsinh}(\sqrt{d}\tilde{w})$. Hence a reparametrisation is given by $\sinh^2(\sqrt{d}\bar{w}) = \frac{dw^2}{ew+2+2\sqrt{p}}$. \square

Remark 3.2 (Interpretation of parametrising functions) In the normal form (9), there are 5 parametrising functions¹. However, only $d = d(x)$ and $e = e(x)$ play a significant role in the shape of the submanifold \mathcal{S}_q . Indeed, a is a scaling of the submanifold, c is the value of h at $w = 0$, and by an appropriate choice of coordinates (as in (12)) we can always assume that $b \equiv 0$. From the above corollary, the role of d is clear: its sign around $x_0 = 0 \in \mathbb{R}^2$ determines the nature of the submanifold, that is, whether the submanifold is elliptic, hyperbolic, or parabolic.

¹ On <https://www.geogebra.org/m/tyb4yggpb> the reader can play with those parameters (the functions a, b, c, d , and e become real numbers when fixing $x \in \mathcal{X}$).

The role of the function e is, however, more subtle. Clearly, h is well defined whenever $p > 0$ and, for a given d , the function e determines the region in which $p > 0$ (in particular, whether h is defined globally with respect to w or not). If $d \equiv 0$, then $p > 0$ holds everywhere (h is defined globally) if and only if $e \equiv 0$ that is, h is explicitly given by $h = 2aw^2 + bw + c$. If $d < 0$, then we have $p > 0$ only between its roots and the parametrisation is never global. Finally, if $d > 0$ then the parametrisation is global if and only if $\Delta < 0$ (where Δ is the discriminant of p), that is $|e| < 2\sqrt{d}$.

4 Classification of quadratic systems

Notations.

$\Xi_Q = \{\Xi_{EH}, \Xi_P\}$, $\Xi_{EH} = \{\Xi_E, \Xi_H\}$	First extension of elliptic, hyperbolic, parabolic submanifolds, seen as a control-nonlinear system on \mathcal{X} .
$\Xi_Q = (A, B, C)$	Triple of vector fields attached to a first extension of a quadratic submanifold.
(α, β)	Reparametrisation (feedback) acting on quadratic nonlinear systems, given by Proposition 4.1 .
(μ_0, μ_1) , $\gamma = (\gamma_0, \gamma_1)$	Structure functions attached to (A, B, C) by $[A, B] = \mu_0 A + \mu_1 B$ and $C = \gamma_0 A + \gamma_1 B$; see (16) .
ι	Notation for $\iota = (\gamma_1, \mp\gamma_0)$, used in the elliptic and hyperbolic cases.
$\Gamma_E, \Gamma_H, \Gamma_P$	Functions given by $\Gamma_E = (\gamma_0)^2 + (\gamma_1)^2$, $\Gamma_H = (\gamma_0)^2 - (\gamma_1)^2$, and $\Gamma_P = \gamma_0 + (\gamma_1)^2$.
\mathfrak{g}_\pm , κ_\pm	A (pseudo-)Riemannian metric defined by $\mathfrak{g}_\pm(A, A) = 1$, $\mathfrak{g}_\pm(B, B) = \pm 1$, and $\mathfrak{g}_\pm(A, B) = 0$, and κ_\pm its Gaussian curvature; see (19) .

From [Theorem 3.1](#) we know how to characterise control-affine system equivalent to the quadratic form Σ_Q and, in particular, we know how to characterise the subclasses of elliptic, hyperbolic, and parabolic systems (see [Corollary 3.1](#)). We are now interested in classifying, under feedback transformations, the systems inside those three subclasses because, due to [Proposition 2.1](#), their classification provides a classification of the elliptic, hyperbolic, and parabolic submanifolds that they parametrise (see [Lemma 4.1](#) below). To this end, we consider the quadratic nonlinear system

$$\Xi_Q : \dot{x} = f_Q(x, w),$$

where $x \in \mathcal{X}$ is the 2-dimensional state, $w \in \mathbb{R}$ plays the role of a control that enters in a nonlinear way and f_Q is a w -parameterised vector field on \mathcal{X} given by either

$$\begin{aligned} f_E &= A(x) \cos(w) + B(x) \sin(w) + C(x), & \text{defining } \Xi_E, & \text{ or} \\ f_H &= A(x) \cosh(w) + B(x) \sinh(w) + C(x), & \text{defining } \Xi_H, & \text{ or} \\ f_P &= A(x)w^2 + B(x)w + C(x), & \text{defining } \Xi_P. & \end{aligned}$$

In each of the three cases, A , B , and C are smooth vector fields on \mathcal{X} satisfying $(A \wedge B)(x_0) \neq 0$. We call Ξ_E an elliptic system, Ξ_H a hyperbolic system, and Ξ_P a parabolic system, because in each fiber $T_x \mathcal{X}$, the system Ξ_E , resp. Ξ_H , resp. Ξ_P parametrises an ellipse, resp. a hyperbola, resp. a parabola. A quadratic nonlinear system Ξ_Q is then represented by the triple (A, B, C) of three smooth vector fields satisfying $A \wedge B \neq 0$. We call the pair (A, B) a Q -frame, and if additionally $[A, B] = 0$, then we call (A, B) a *commutative* Q -frame. We will denote by the index EH object attached to either the elliptic or the hyperbolic case, as those two are treated in a similar manner.

For quadratic submanifolds, elliptic and hyperbolic, of the form $\mathcal{S}_{EH} = \{a^2(\dot{z} - c_0)^2 \pm b^2(\dot{y} - c_1)^2 = 1\}$ and parabolic of the form $\mathcal{S}_P = \{a\dot{y}^2 - \dot{z} + b\dot{y} + c = 0\}$, we distinguished specific classes that are summarised in the following [Table 1](#).

Elliptic and hyperbolic submanifolds classification		Parabolic submanifolds classification	
$a = b$	conformally-flat	$a = 1$	weakly-flat
$a = b = 1$	flat	$a = 1, b = 0$	strongly-flat
$a = b = 1, (c_0, c_1) \in \mathbb{R}^2$	constant-form	$a = 1, b = 0, c \in \mathbb{R}$	constant-form
$a = b = 1, c_0 = c_1 = 0$	null-form	$a = 1, b = 0, \text{ and } c = 0$	null-form

Table 1 Classification of elliptic, hyperbolic, and parabolic submanifolds.

Our goal is to characterise the above types of quadratic submanifolds and we now show in the next lemma that the classification of elliptic, hyperbolic, and parabolic submanifolds \mathcal{S}_Q presented in [Table 1](#) is reflected in properties of the control system $\Xi_Q = (A, B, C)$.

Lemma 4.1 Consider a quadratic submanifold \mathcal{S}_Q together with its regular parametrisation $\Xi_Q = (A, B, C)$.

- (i) \mathcal{S}_{EH} is locally equivalent to a conformally-flat elliptic/hyperbolic submanifold if and only if Ξ_{EH} is locally feedback equivalent to Ξ_{EH} , whose EH-frame (A, B) is given by $A = r(x) \frac{\partial}{\partial z}$ and $B = r(x) \frac{\partial}{\partial y}$ for some nonvanishing function $r(x)$.
- (ii) \mathcal{S}_{EH} is locally equivalent to a flat elliptic/hyperbolic submanifold if and only if Ξ_{EH} is locally feedback equivalent to Ξ_{EH} , whose frame (A, B) is commutative.
- (iii) \mathcal{S}_{EH} is locally equivalent to a constant-form elliptic/hyperbolic submanifold if and only if Ξ_{EH} is locally feedback equivalent to Ξ_{EH} , whose EH-frame (A, B) is commutative and, additionally, $[A, C] = [B, C] = 0$.
- (iv) \mathcal{S}_{EH} is locally equivalent to a null-form elliptic/hyperbolic submanifold if and only if Ξ_{EH} is locally feedback equivalent to Ξ_{EH} , whose EH-frame (A, B) is commutative and, additionally, $C = 0$.
- (v) \mathcal{S}_P is locally equivalent to weakly-flat parabolic submanifold if and only if Ξ_P is locally feedback equivalent to Ξ_P , whose P-frame (A, B) is commutative.
- (vi) \mathcal{S}_P is locally equivalent to strongly-flat parabolic submanifold if and only if Ξ_P is locally feedback equivalent to Ξ_P , whose P-frame (A, B) is commutative and, additionally, $A \wedge C = 0$.
- (vii) \mathcal{S}_P is locally equivalent to constant-form parabolic submanifold if and only if Ξ_P is locally feedback equivalent to Ξ_P , whose P-frame (A, B) is commutative and, additionally, $[A, C] = [B, C] = 0$.
- (viii) \mathcal{S}_P is locally equivalent to null-form parabolic submanifold if and only if Ξ_P is locally feedback equivalent to Ξ_P , whose P-frame (A, B) is commutative and, additionally, $C = 0$.

Recall that a general conic submanifold \mathcal{S}_q is equivalent to an elliptic \mathcal{S}_E , resp. a hyperbolic \mathcal{S}_H , resp. a parabolic \mathcal{S}_P , submanifold if and only the determinant Δ_2 satisfies $\Delta_2 > 0$, resp. $\Delta_2 < 0$, resp. $\Delta_2 \equiv 0$; see [Lemma 2.1](#). Therefore the above lemma allows to check equivalence of \mathcal{S}_q to a submanifold of any of the subclasses listed in [Table 1](#).

Proof It is a straightforward computation to check that for the submanifolds \mathcal{S}_Q of the indicated forms, their first extensions $\Xi_Q = (A, B, C)$ have the triple (A, B, C) or the Q-frame (A, B) satisfying the above stated conditions. Conversely, in local coordinates $x = (z, y)$ in which either $A = r(x) \frac{\partial}{\partial z}$ and $B = r(x) \frac{\partial}{\partial y}$, for (i), or $A = \frac{\partial}{\partial z}$ and $B = \frac{\partial}{\partial y}$, for (ii) to (viii), the system Ξ_Q is a regular parametrisation of \mathcal{S}_Q with the desired properties. Now, all items (i) to (viii) follow from [Proposition 2.1](#). \square

The above lemma asserts that to achieve the classification of elliptic, hyperbolic, and parabolic submanifolds presented in [Table 1](#), it is crucial to classify, under feedback transformations, quadratic control systems $\Xi_Q = (A, B, C)$ with the properties presented in [Table 2](#), which will be the goal of the remaining part of this section. In particular, we will show that the characterisation of item (i), resp. item (v), is always satisfied by any Ξ_{EH} and thus by the corresponding \mathcal{S}_{EH} , resp. by any Ξ_P and thus by the corresponding \mathcal{S}_P , while the characterisations of the remaining classes of systems, and thus of the corresponding submanifolds, require non-trivial conditions.

Elliptic and hyperbolic classification		Parabolic classification	
conformally-flat	$A = r(x) \frac{\partial}{\partial z}$ and $B = r(x) \frac{\partial}{\partial y}$	weakly-flat	$[A, B] = 0$
flat	$[A, B] = 0$	strongly-flat	$[A, B] = 0$ and $A \wedge C = 0$
constant-form	$[A, B] = [A, C] = [B, C] = 0$,	constant-form	$[A, B] = [A, C] = [B, C] = 0$
null-form	$[A, B] = 0$ and $C = 0$	null-form	$[A, B] = 0$ and $C = 0$

Table 2 Reflection of the classification of elliptic, hyperbolic, and parabolic submanifolds in properties of the triple $\Xi_Q = (A, B, C)$.

Although the systems of the form Ξ_Q are nonlinear with respect to the control w , the feedback transformations that preserve this class are not as general as possible. Indeed, feedback transformations which preserve the class of quadratic system Ξ_Q are *affine* (and even of Brockett type, in the case of elliptic and hyperbolic systems) with respect to the control w , as assured by the next proposition, which also shows how feedback acts on the triple (A, B, C) .

Proposition 4.1 (Reparametrisation of quadratic nonlinear systems) Consider two quadratic systems Ξ_Q and $\tilde{\Xi}_Q$ around (x_0, w_0) and $(\tilde{x}_0, \tilde{w}_0)$, respectively.

- (i) Two elliptic systems Ξ_E and $\tilde{\Xi}_E$ are locally feedback equivalent if and only if there exists a local diffeomorphism $\tilde{x} = \phi(x)$ and a reparametrisation (feedback) $w = \psi(x, \tilde{w})$, given by $\psi = \pm \tilde{w} + \alpha(x)$, and satisfying

$$\tilde{A} = \phi_* (A \cos \alpha + B \sin \alpha), \quad \tilde{B} = \pm \phi_* (-A \sin \alpha + B \cos \alpha), \quad \tilde{C} = \phi_* (C). \quad (13)$$

- (ii) Two hyperbolic systems Ξ_H and $\tilde{\Xi}_H$ are locally feedback equivalent if and only if there exists a local diffeomorphism $\tilde{x} = \phi(x)$ and a reparametrisation (feedback) $w = \psi(x, \tilde{w})$, given by $\psi = \pm\tilde{w} + \alpha(x)$, and satisfying

$$\tilde{A} = \phi_* (A \cosh \alpha + B \sinh \alpha), \quad \tilde{B} = \pm \phi_* (A \sinh \alpha + B \cosh \alpha), \quad \tilde{C} = \phi_* (C). \quad (14)$$

- (iii) Two parabolic systems Ξ_P and $\tilde{\Xi}_P$ are locally feedback equivalent if and only if there exists a local diffeomorphism $\tilde{x} = \phi(x)$ and an invertible reparametrisation (feedback) $w = \psi(x, \tilde{w})$, given by $\psi = \alpha(x) + \beta(x)\tilde{w}$ and $\beta(\cdot) \neq 0$, and satisfying

$$\tilde{A} = \phi_* (A\beta^2), \quad \tilde{B} = \phi_* (2A\alpha\beta + B\beta), \quad \tilde{C} = \phi_* (C + A\alpha^2 + B\alpha). \quad (15)$$

Proof We show the necessity of each statement as the converse implications are immediate.

- (i) Assume that Ξ_E and $\tilde{\Xi}_E$ are locally equivalent via a diffeomorphism $\tilde{x} = \phi(x)$ and a reparametrisation $w = \psi(x, \tilde{w})$. Then we have the following relation $\phi_* f_E(x, \psi(x, \tilde{w})) = \tilde{f}_E(\tilde{x}, \tilde{w})$, which we differentiate 3 times with respect to \tilde{w} and using $\frac{\partial^3 \tilde{f}_E}{\partial \tilde{w}^3} = -\frac{\partial \tilde{f}_E}{\partial \tilde{w}}$, we conclude the relation $\phi_* \frac{\partial^3 f_E}{\partial \tilde{w}^3} = -\phi_* \frac{\partial f_E}{\partial \tilde{w}}$, which translates into

$$\begin{aligned} & A \left(-\psi''' \sin(\psi) + (\psi')^3 \sin(\psi) - 3\psi' \psi'' \cos(\psi) \right) \\ & + B \left(\psi''' \cos(\psi) - (\psi')^3 \cos(\psi) - 3\psi' \psi'' \sin(\psi) \right) = A\psi' \sin(\psi) - B\psi' \cos(\psi), \end{aligned}$$

where the derivatives are taken with respect to \tilde{w} . Since the functions \cos and \sin are linearly independent, we obtain $\psi'' = 0$ and $(\psi')^2 = 1$. Thus $\psi(x, \tilde{w}) = \pm\tilde{w} + \alpha(x)$. Applying this reparametrisation we obtain relations (13).

- (ii) Exactly the same reasoning, using f_H and the fact $\phi_* \frac{\partial^3 f_H}{\partial \tilde{w}^3} = \phi_* \frac{\partial f_H}{\partial \tilde{w}}$, implies that $\psi(x, \tilde{w}) = \pm\tilde{w} + \alpha(x)$. Applying $\phi(x)$ and $w = \psi(x, \tilde{w}) = \pm\tilde{w} + \alpha$ we obtain relations (14).
- (iii) We repeat again the same reasoning to f_P with the property $\phi_* \frac{\partial^3 f_P}{\partial \tilde{w}^3} = 0$. However, this time we obtain the conditions $\psi''' = 0$ and $\psi\psi''' + 3\psi'\psi'' = 0$ on the reparametrisation ψ , which implies $\psi'' = 0$, that is $\psi(x, \tilde{w}) = \beta(x)\tilde{w} + \alpha(x)$, with β satisfying $\beta(\cdot) \neq 0$. Applying this reparametrisation together with a diffeomorphism ϕ yields relations (15). \square

Remark 4.1 (Local character of the results) Initially, Ξ_Q was considered locally around a point x_0 and a control w_0 , however, since Ξ_Q is defined globally with respect to w and, moreover, by the last proposition, the transformations $w = \psi(x, \tilde{w})$ are *global* with respect to w , so we will consider the systems Ξ_Q and their equivalence locally in x and globally with respect to w . All results below are stated assuming this structure.

We will develop relations involving structure functions attached to any fixed triple (A, B, C) in a unique way and thus change accordingly with diffeomorphisms $\tilde{x} = \phi(x)$. So we will act on (A, B, C) by (α, β) only (β is ± 1 in the elliptic and hyperbolic cases) and we will denote by $(\tilde{A}, \tilde{B}, \tilde{C})$ the result of that action (given by (13), or (14), or (15), with $\phi = \text{id}$), called a reparametrisation.

Observe that the reparametrisations of Ξ_P depend on two smooth functions α and β while those of Ξ_E and Ξ_H depend on one smooth function α only. Therefore we expect the classification of parabolic systems to be less rich (less parametrising functions) than the classification of elliptic and hyperbolic systems. In the following subsections we will first classify elliptic and hyperbolic systems as the procedures are similar, and then we will classify parabolic systems under reparametrisation actions.

4.1 Classification of elliptic and hyperbolic systems

In this subsection, we classify elliptic and hyperbolic systems under the action of reparametrisations. Recall, that our aim is to classify elliptic and hyperbolic nonholonomic constraints (submanifolds of $T\mathcal{X}$) \mathcal{S}_E and \mathcal{S}_H , which are parameterised by systems of the form Ξ_E and Ξ_H , respectively. The classification of submanifolds given in Table 1 is reflected in special properties of the vector fields (A, B, C) , attached to the control system Ξ_E and Ξ_H , that we list in Table 2 above and summarise in Lemma 4.1. Firstly, we will give a normal form for both types of systems Ξ_E and Ξ_H showing that they actually depend on three smooth functions, that normal form corresponds to conformally-flat elliptic and hyperbolic submanifolds. Secondly, we will further develop their classification, in particular we will give conditions for the existence of commutative frames (corresponding to flat elliptic and hyperbolic submanifolds) and a complete characterisation of forms without functional parameters, corresponding to constant-form (and, in particular, null-form) elliptic and hyperbolic submanifolds.

Notations. In order to simplify and unify notations, in the following formulae the upper sign always corresponds to the elliptic case and the lower sign to the hyperbolic case, e.g. we will use the symbol \pm to design similar objects attached to the elliptic (+ case) and to the hyperbolic (− case) systems and in the case of a \mp symbol we have − for elliptic systems and + for hyperbolic ones. We denote Ξ_{EH} elliptic and hyperbolic systems, and an EH-frame stands for an E-frame or an H-frame. To avoid unnecessary computations, we assume that EH-frames (A, B) and (\tilde{A}, \tilde{B}) of two equivalent systems have the same orientation (we will come back to this simplification in [Proposition 4.4](#)), therefore we restrict reparametrisations of the control w , given by (13) and (14) to those with $\beta = 1$. Denoting by $\bar{R}_{EH}(\alpha)$ the (trigonometric or hyperbolic) rotation matrix given by

$$\bar{R}_E(\alpha) = \begin{pmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{pmatrix}, \quad \text{and} \quad \bar{R}_H(\alpha) = \begin{pmatrix} \cosh(\alpha) & -\sinh(\alpha) \\ -\sinh(\alpha) & \cosh(\alpha) \end{pmatrix},$$

respectively, we see from (13) and (14) that EH-frames are transformed by $(\tilde{A}, \tilde{B}) = (A, B)\bar{R}_{EH}(\pm\alpha)$ under reparametrisations of the form $w = \tilde{w} + \alpha$. Introduce structure functions (μ_0, μ_1) and (γ_0, γ_1) uniquely defined by

$$[A, B] = \mu_0 A + \mu_1 B \quad \text{and} \quad C = \gamma_0 A + \gamma_1 B, \quad (16)$$

respectively. We denote $\gamma = (\gamma_0, \gamma_1)$, and $\mathcal{L} = (\gamma_1, \mp\gamma_0)$, and set $\Gamma_{EH} = (\gamma_0)^2 \pm (\gamma_1)^2$.

We begin by a technical lemma showing how structure functions behave under reparametrisations of the control w .

Lemma 4.2 (Transformation of structure functions) *Consider an elliptic/hyperbolic system Ξ_{EH} with structure functions $(\mu_0, \mu_1, \gamma_0, \gamma_1)$. Then under the reparametrisation $w = \tilde{w} + \alpha(x)$ we have*

$$(\tilde{\mu}_0, \tilde{\mu}_1) = (\mu_0 \mp L_A(\alpha), \mu_1 - L_B(\alpha))\bar{R}_{EH}(\alpha), \quad \text{and} \quad \tilde{\gamma} = \gamma\bar{R}_{EH}(\alpha). \quad (17)$$

Proof Details of the computations can be found in [Appendix D](#). □

Clearly from (17), $\Gamma_{EH} = (\gamma_0)^2 \pm (\gamma_1)^2$ is invariant under reparametrisations, i.e. $\tilde{\Gamma}_{EH} = \Gamma_{EH}$.

Proposition 4.2 (Conformal form of elliptic and hyperbolic systems)

(i) *Any elliptic system Ξ_E , resp. hyperbolic system Ξ_H , always admits under a reparametrisation the following conformal form, locally around x_0 ,*

$$\Xi_E^c : \dot{x} = r(x) \begin{pmatrix} \cos(w) \\ \sin(w) \end{pmatrix} + \begin{pmatrix} c_0(x) \\ c_1(x) \end{pmatrix}, \quad \text{resp.} \quad \Xi_H^c : \dot{x} = r(x) \begin{pmatrix} \cosh(w) \\ \sinh(w) \end{pmatrix} + \begin{pmatrix} c_0(x) \\ c_1(x) \end{pmatrix},$$

with r a smooth function satisfying $r > 0$.

(ii) *Two conformal forms Ξ_E^c and $\tilde{\Xi}_E^c$, resp. Ξ_H^c and $\tilde{\Xi}_H^c$, are locally feedback equivalent if and only if there exists a local diffeomorphism $\tilde{x} = \phi(x) = (\phi_1(x), \phi_2(x))$, where $x = (z, y)$, satisfying*

$$\frac{\partial\phi_1}{\partial z} = \frac{\partial\phi_2}{\partial y}, \quad \frac{\partial\phi_1}{\partial y} = \mp \frac{\partial\phi_2}{\partial z}, \quad \left(\frac{\partial\phi_1}{\partial z}\right)^2 \pm \left(\frac{\partial\phi_1}{\partial y}\right)^2 = \left(\frac{\tilde{r}}{r}\right)^2, \quad \text{and} \quad \phi_*C = \tilde{C}. \quad (18)$$

We call Ξ_{EH}^c a conformal-form because the systems of that class parametrise elliptic and hyperbolic submanifolds for which the quadratic term, interpreted as a (pseudo-)Riemannian metric, is conformally flat.

Proof

(i) For the system $\Xi_{EH} = (A, B, C)$, define a (pseudo-)Riemannian metric \mathbf{g}_\pm on \mathcal{X} by $\mathbf{g}_\pm(A, A) = 1$, $\mathbf{g}_\pm(B, B) = \pm 1$, and $\mathbf{g}_\pm(A, B) = 0$. It is known that any non-degenerate metric on a manifold of dimension two is conformally flat (see [1, pp 15-35] or [20, Addendum 1 of chapter 9] for the elliptic case and [19, theorem 7.2] for the hyperbolic case). Therefore, there exists a local diffeomorphism $(\tilde{z}, \tilde{y}) = \tilde{x} = \phi(x)$ such that $\mathbf{g}_\pm = \phi^*\tilde{\mathbf{g}}_\pm$, where $\tilde{\mathbf{g}}_\pm = \varrho(\tilde{x})(d\tilde{z}^2 \pm d\tilde{y}^2)$, $\varrho > 0$. The vector fields $\tilde{A} = \phi_*A$ and $\tilde{B} = \phi_*B$ satisfy $\tilde{\mathbf{g}}_\pm(\tilde{A}, \tilde{A}) = 1$, $\tilde{\mathbf{g}}_\pm(\tilde{B}, \tilde{B}) = \pm 1$, and $\tilde{\mathbf{g}}_\pm(\tilde{A}, \tilde{B}) = 0$ which implies that (\tilde{A}, \tilde{B}) is a (pseudo-)orthonormal frame. Finally, using feedback α we can smoothly rotate (\tilde{A}, \tilde{B}) into $(r\frac{\partial}{\partial\tilde{z}}, r\frac{\partial}{\partial\tilde{y}})$ with $r = \frac{1}{\sqrt{\varrho}}$, which gives the desired form $\Xi_{EH}^c = (\tilde{A}, \tilde{B}, \tilde{C})$, with $\tilde{C} = \phi_*C$.

(ii) By relations (13) and (14), the reparametrisations do not act on C and thus the relation $\tilde{C} = \phi_*C$ is necessary for the equivalence of conformal forms. Consider two elliptic conformal systems Ξ_E^c and $\tilde{\Xi}_E^c$, resp. two hyperbolic conformal systems Ξ_H^c and $\tilde{\Xi}_H^c$, with frames (A, B) and (\tilde{A}, \tilde{B}) and related by a feedback $w = \tilde{w} + \alpha$ and a diffeomorphism ϕ . Thus, using relation (13), resp. (14), we obtain

$$\begin{aligned} \frac{\partial \phi_1}{\partial z} &= \frac{\tilde{r}}{r} \cos(\alpha) = \frac{\partial \phi_2}{\partial y}, & \frac{\partial \phi_1}{\partial y} &= \frac{\tilde{r}}{r} \sin(\alpha) = -\frac{\partial \phi_2}{\partial z}, \\ \text{resp. } \frac{\partial \phi_1}{\partial z} &= \frac{\tilde{r}}{r} \cosh(\alpha) = \frac{\partial \phi_2}{\partial y}, & \frac{\partial \phi_1}{\partial y} &= -\frac{\tilde{r}}{r} \sinh(\alpha) = \frac{\partial \phi_2}{\partial z}, \end{aligned}$$

from which we deduce condition (18). Conversely, applying the diffeomorphism ϕ given by (18), together with the feedback $w = \tilde{w} + \alpha$, with α being a solution of

$$\begin{aligned} \cos(\alpha) &= \frac{r}{\tilde{r}} \frac{\partial \phi_1}{\partial z}, & \sin(\alpha) &= -\frac{r}{\tilde{r}} \frac{\partial \phi_1}{\partial y}, \\ \text{resp. } \cosh(\alpha) &= \frac{r}{\tilde{r}} \frac{\partial \phi_1}{\partial z}, & \sinh(\alpha) &= -\frac{r}{\tilde{r}} \frac{\partial \phi_1}{\partial y}, \end{aligned}$$

we transform Ξ_E^c into $\tilde{\Xi}_E^c$, resp. Ξ_H^c into $\tilde{\Xi}_H^c$. □

Remark 4.2 In the above proof we used the metric \mathbf{g}_\pm on \mathcal{X} defined by

$$\mathbf{g}_\pm(A, A) = 1, \quad \mathbf{g}_\pm(B, B) = \pm 1, \quad \mathbf{g}_\pm(A, B) = 0. \quad (19)$$

This object will play a special role in the interpretation of the conditions describing the existence of a commutative EH-frame.

The above proposition shows that elliptic and hyperbolic systems Ξ_{EH} are parametrized by three smooth functions of two variables (and not by 6 functions defining the triple (A, B, C)). Moreover, relation (18) shows that the group of diffeomorphisms, conjugating the EH-frames of two given conformal forms, is parametrized by any function ϕ_1 of two variables satisfying the third equation of (18), ϕ_2 being given in terms of ϕ_1 via the first and second equations. Now we will pass to the problem of commutative frames and the following proposition gives equivalent algebraic and geometric conditions for the existence of a commutative EH-frame.

Proposition 4.3 (Existence of a commutative EH-frame) *Consider an elliptic/hyperbolic system $\Xi_{EH} = (A, B, C)$ with structure functions (μ_0, μ_1) of the EH-frame (A, B) . The following statements are equivalent locally around x_0 :*

- (i) *There exists a commutative EH-frame.*
- (ii) *The structure functions (μ_0, μ_1) attached to the EH-frame (A, B) satisfy*

$$-(\mu_0)^2 \mp (\mu_1)^2 \pm L_A(\mu_1) - L_B(\mu_0) = 0. \quad (20)$$

- (iii) *The Gaussian curvature κ_\pm of the metric \mathbf{g}_\pm vanishes.*

Notice that item (i) describes the following normal forms,

$$\Xi_E' : \begin{cases} \dot{z} = \cos(w) + c_0(x) \\ \dot{y} = \sin(w) + c_1(x) \end{cases}, \quad \text{and} \quad \Xi_H' : \begin{cases} \dot{z} = \cosh(w) + c_0(x) \\ \dot{y} = \sinh(w) + c_1(x) \end{cases},$$

whose structure functions are $\mu_0 = \mu_1 = 0$, $\gamma_0 = c_0$, and $\gamma_1 = c_1$. We call Ξ_E' a *flat elliptic system* and Ξ_H' a *flat hyperbolic system*.

Proof The equivalence between (ii) and (iii) is immediate since (20) is the Gaussian curvature κ_\pm of \mathbf{g}_\pm (details of the computations are in Appendix E). We show that (i) is equivalent to (ii). If the EH-frame (A, B) is equivalent via $w = \tilde{w} + \alpha(x)$ to a commutative EH-frame (\tilde{A}, \tilde{B}) , then by (17) we immediately have $L_A(\alpha) = \pm\mu_0$ and $L_B(\alpha) = \mu_1$; the integrability condition of this system of first order partial differential equations gives (20). Conversely, consider the system $\Xi_{EH} = (A, B, C)$ and construct α as a solution of the system $L_A(\alpha) = \pm\mu_0$ and $L_B(\alpha) = \mu_1$, whose solvability is guaranteed by the integrability condition given by (20). Then by (17) we see that the resulting EH-frame (\tilde{A}, \tilde{B}) , of the system $\tilde{\Xi}_{EH}$ obtained by the reparametrisation $w = \tilde{w} + \alpha$, is commutative. □

Notice that when proving [Proposition 4.3](#) we have shown that the Gaussian curvature κ_{\pm} of the metric g_{\pm} is given by the left hand side of [\(20\)](#). Moreover relation [\(17\)](#) implies that κ_{\pm} is invariant under reparametrisations $w = \tilde{w} + \alpha$ and is therefore an equivariant of the feedback transformations of the system Ξ_{EH} .

In the following proposition, we give first a classification of flat elliptic/hyperbolic systems, second we characterise those without functional parameters, i.e. constant-forms, and third we provide a canonical form for the latter. Recall that $\iota = (\gamma_1, \mp\gamma_0)$ and that for flat elliptic/hyperbolic systems Ξ'_{EH} we have $(\gamma_0, \gamma_1) = (c_0, c_1)$ so all statements of the proposition below are actually expressed in terms of structure functions. From now on, we will consider the group of feedback transformations consisting of $\tilde{x} = \phi(x)$ and $w = \pm\tilde{w} + \alpha(x)$. The additional transformation $w = -\tilde{w} + \alpha$ implies $(\tilde{A}, \tilde{B}) = (A, B)\bar{R}_{EH}(\pm\alpha)$, where $\bar{R}_E(\alpha) = \begin{pmatrix} \cos(\alpha) & \sin(\alpha) \\ \sin(\alpha) & -\cos(\alpha) \end{pmatrix}$ and $\bar{R}_H(\alpha) = \begin{pmatrix} \cosh(\alpha) & \sinh(\alpha) \\ -\sinh(\alpha) & -\cosh(\alpha) \end{pmatrix}$, and the corresponding structure functions change by, compare [\(17\)](#),

$$(\tilde{\mu}_0, \tilde{\mu}_1) = -(\mu_0 \mp L_A(\alpha), \mu_1 - L_B(\alpha))\bar{R}_{EH}(\alpha) \quad \text{and} \quad \tilde{\gamma} = \gamma\bar{R}_{EH}(\alpha). \quad (21)$$

Proposition 4.4 (Characterisation and classification of flat elliptic/hyperbolic systems)

(i) Two flat elliptic systems Ξ'_E and $\tilde{\Xi}'_E$, resp. two flat hyperbolic systems Ξ'_H and $\tilde{\Xi}'_H$, are locally feedback equivalent around $x_0 = 0 \in \mathbb{R}^2$ if and only if there exists a constant $\alpha \in \mathbb{R}$ satisfying

$$R_{EH}(\pm\alpha)^{-1}C(x) = \tilde{C}\left(R_{EH}(\pm\alpha)^{-1}x\right), \quad (22)$$

where R_{EH} stands for either \bar{R}_{EH} or \bar{R}_{EH} .

(ii) An elliptic/hyperbolic system Ξ_{EH} is locally feedback equivalent to a constant-form, i.e. Ξ'_{EH} with $(c_0, c_1) \in \mathbb{R}^2$, if and only if one of the equivalent conditions of [Proposition 4.3](#) holds and, additionally,

$$L_A(\gamma) + \iota\mu_0 = 0 \quad \text{and} \quad L_B(\gamma) \pm \iota\mu_1 = 0. \quad (23)$$

(iii) A constant-form elliptic system is always feedback equivalent, locally around $x_0 = 0 \in \mathbb{R}^2$, to the canonical form

$$\Xi_E^{\Gamma_E} : \begin{cases} \dot{z} = \cos(w) + \sqrt{\Gamma_E} \\ \dot{y} = \sin(w) \end{cases},$$

where $\Gamma_E = (c_0)^2 + (c_1)^2 \in \mathbb{R}$ is an invariant.

(iv) A constant-form hyperbolic system is always feedback equivalent, locally around $x_0 = 0 \in \mathbb{R}^2$, to one of the following canonical form

$$\begin{aligned} \Xi_H^{\Gamma_H, \varepsilon} : \begin{cases} \dot{z} = \cosh(w) + \varepsilon\sqrt{\Gamma_H} \\ \dot{y} = \sinh(w) \end{cases}, \quad \text{or} \quad \Xi_H^{-\Gamma_H} : \begin{cases} \dot{z} = \cosh(w) \\ \dot{y} = \sinh(w) + \sqrt{-\Gamma_H} \end{cases}, \\ \text{or} \quad \Xi_H^{0, \varepsilon} : \begin{cases} \dot{z} = \cosh(w) + \varepsilon \\ \dot{y} = \sinh(w) + 1 \end{cases}, \quad \text{or} \quad \Xi_H^{0, 0} : \begin{cases} \dot{z} = \cosh(w) \\ \dot{y} = \sinh(w) \end{cases}, \end{aligned}$$

where $\Gamma_H = (c_0)^2 - (c_1)^2 \in \mathbb{R}$ and satisfies $\Gamma_H > 0$ for the first form, $\Gamma_H < 0$ for the second form, and $\Gamma_H = 0$ for the third and fourth ones, where $\varepsilon = \text{sgn}(c_0) = \pm 1$. Moreover (Γ_H, ε) is a complete invariant.

Observe that if in items [\(i\)](#), [\(iii\)](#), and [\(iv\)](#) the considered systems are defined globally, then their feedback equivalence is also global and, in particular, the proposed canonical forms are also global.

Remark 4.3 In item [\(iv\)](#), notice that there are two orbits of the local action of feedback transformations group for $\Gamma_H > 0$, corresponding to $\text{sgn}(c_0) = \varepsilon = \pm 1$, one orbit for $\Gamma_H < 0$, and three orbits for $\Gamma_H = 0$ corresponding, respectively, to $\text{sgn}(c_0) = \varepsilon = \pm 1$ or $(c_0, c_1) = (0, 0)$. The invariant $\varepsilon = \pm 1$ corresponds to the parametrisation of one of two branches of the hyperbola $(\dot{z} - \sqrt{\Gamma_H})^2 - \dot{y}^2 = 1$.

Proof

(i) Consider, locally around $0 \in \mathbb{R}^2$, two equivalent flat elliptic/hyperbolic systems Ξ'_{EH} and $\tilde{\Xi}'_{EH}$ given by structure functions $(\mu_0, \mu_1, \gamma_0, \gamma_1) = (0, 0, c_0, c_1)$ and $(\tilde{\mu}_0, \tilde{\mu}_1, \tilde{\gamma}_0, \tilde{\gamma}_1) = (0, 0, \tilde{c}_0, \tilde{c}_1)$, respectively. Since they both have a commutative EH-frame, by [\(17\)](#) (and [\(21\)](#)) they differ by a reparametrisation $w = \pm\tilde{w} + \alpha$ satisfying $L_A(\alpha) = L_B(\alpha) = 0$ and thus $\alpha \in \mathbb{R}$. Applying this reparametrisation together with a diffeomorphism ϕ satisfying $\phi_* = R_{EH}(\pm\alpha)^{-1}$, that is $\tilde{x} = \phi(x) = R_{EH}(\pm\alpha)^{-1}x$, transforms Ξ'_{EH} into $\tilde{\Xi}'_{EH}$ if and only if

$$\begin{pmatrix} \tilde{c}_0(\tilde{x}) \\ \tilde{c}_1(\tilde{x}) \end{pmatrix} = R_{EH}(\pm\alpha)^{-1} \begin{pmatrix} c_0(x) \\ c_1(x) \end{pmatrix},$$

which is [\(23\)](#).

- (ii) Assume that Ξ_{EH} , given by structure functions $(\mu_0, \mu_1, \gamma_0, \gamma_1)$, is equivalent via $\tilde{x} = \phi(x)$ and $w = \tilde{w} + \alpha$ to Ξ'_{EH} with structure functions $(\tilde{\mu}_0, \tilde{\mu}_1, \tilde{\gamma}_0, \tilde{\gamma}_1) = (0, 0, c_0, c_1)$, where $(c_0, c_1) \in \mathbb{R}^2$. Necessity of one (and thus any) of the conditions of [Proposition 4.3](#) is clear, and by (17) and (21) we have first, $L_A(\alpha) = \pm\mu_0$ and $L_B(\alpha) = \mu_1$ and second, $\gamma R_{EH}(\alpha) = \tilde{\gamma} = (c_0, c_1)$. By differentiating this last relation along A and B we obtain

$$\begin{aligned} 0 &= L_A(\gamma) R_{EH}(\alpha) + \gamma L_A(R_{EH}(\alpha)) = L_A(\gamma) R_{EH}(\alpha) + \gamma \begin{pmatrix} 0 & \mp 1 \\ 1 & 0 \end{pmatrix} R_{EH}(\alpha), \\ 0 &= L_A(\gamma) + \iota\mu_0, \quad \text{and} \\ 0 &= L_B(\gamma) R_{EH}(\alpha) + \gamma L_B(R_{EH}(\alpha)) = L_B(\gamma) R_{EH}(\alpha) + \gamma \begin{pmatrix} 0 & \mp 1 \\ 1 & 0 \end{pmatrix} R_{EH}(\alpha), \\ 0 &= L_B(\gamma) \pm \iota\mu_1, \end{aligned}$$

thus proving that (23) holds. Conversely, assume that (20) and (23) hold for Ξ_{EH} . By [Proposition 4.3](#), Ξ_{EH} is equivalent to Ξ'_{EH} with a commutative EH-frame (A, B) , and applying (23) to the latter we get $L_A(\gamma) = L_B(\gamma) = 0$ and therefore, we have $(c_0, c_1) \in \mathbb{R}^2$.

- (iii) Consider a flat elliptic system Ξ'_E with $(c_0, c_1) \in \mathbb{R}^2$, then relation (22) reads

$$\begin{pmatrix} \tilde{c}_0 \\ \tilde{c}_1 \end{pmatrix} = \begin{pmatrix} \cos(\alpha) & \sin(\alpha) \\ \mp \sin(\alpha) & \pm \cos(\alpha) \end{pmatrix} \begin{pmatrix} c_0 \\ c_1 \end{pmatrix}. \quad (22')$$

Take α as a solution of $-\sin(\alpha)c_0 + \cos(\alpha)c_1 = 0$, then we have $\tilde{c}_1 = 0$ and $\tilde{c}_0 = \pm\sqrt{\Gamma_E}$, with $\Gamma_E = (c_0)^2 + (c_1)^2$. If necessary, apply $\alpha = \pi$ to send $(\tilde{c}_0, \tilde{c}_1) = (-\sqrt{\Gamma_E}, 0)$ into $(\sqrt{\Gamma_E}, 0)$. The proof that $\Xi_E^{\Gamma_E}$ is equivalent to $\tilde{\Xi}_E^{\Gamma_E}$ if and only if $\Gamma_E = \tilde{\Gamma}_E$ is immediate from (22').

- (iv) Consider a flat hyperbolic system Ξ'_H with $(c_0, c_1) \in \mathbb{R}^2$ and denote $\Gamma_H = (c_0)^2 - (c_1)^2$, then relation (22) reads

$$\begin{pmatrix} \tilde{c}_0 \\ \tilde{c}_1 \end{pmatrix} = \begin{pmatrix} \cosh(\alpha) & -\sinh(\alpha) \\ \mp \sinh(\alpha) & \pm \cosh(\alpha) \end{pmatrix} \begin{pmatrix} c_0 \\ c_1 \end{pmatrix}. \quad (22'')$$

We consider four cases. First, assume that $\Gamma_H > 0$, that is $c_0 \neq 0$ and $-1 < \frac{c_1}{c_0} < 1$, and take α as the solution of $\tanh(\alpha) = \frac{c_1}{c_0}$ yielding $\tilde{c}_1 = 0$ and $\tilde{c}_0 = \text{sgn}(c_0)\sqrt{\Gamma_H}$, which gives the canonical form $\Xi_H^{\Gamma_H, \varepsilon}$. Second, assume that $\Gamma_H < 0$, that is $c_1 \neq 0$ and $-1 < \frac{c_0}{c_1} < 1$, and take α as the solution of $\tanh(\alpha) = \frac{c_0}{c_1}$ yielding $\tilde{c}_0 = 0$ and $\tilde{c}_1 = \text{sgn}(c_1)\sqrt{-\Gamma_H}$. If $\text{sgn}(c_1) = -1$, then by applying (22'') with $\alpha = 0$ and the bottom sign, we can always normalize $\text{sgn}(c_1)$ yielding the canonical form $\Xi_H^{-\Gamma_H}$. Third, assume that $\Gamma_H = 0$ and $c_0 = 0$ thus $c_1 = 0$ and therefore we immediately have the canonical form $\Xi_H^{0,0}$. Fourth, and finally, assume that $\Gamma_H = 0$ and $c_0 \neq 0$, thus $c_1 = \varepsilon c_0$ with $\varepsilon = \pm 1$. If necessary, apply (22'') with $\alpha = 0$ and the bottom sign to obtain $c_1 > 0$. Take $\alpha = \varepsilon \ln c_1$ and apply (22'') with the upper sign to obtain $\tilde{c}_1 = 1$ and $\tilde{c}_0 = \varepsilon$. To show that (Γ_H, ε) is a complete invariant is trivial by applying (22'') to the canonical forms $\Xi_H^{\Gamma_H, \varepsilon}$, $\Xi_H^{-\Gamma_H}$, $\Xi_H^{0, \varepsilon}$, and $\Xi_H^{0,0}$. \square

We summarise the results of this subsection, we started from a general elliptic, resp. hyperbolic, system Ξ_E , resp. Ξ_H , which parametrises a elliptic, resp. a hyperbolic, submanifold \mathcal{S}_E , resp. \mathcal{S}_H . We show that reparametrisations (pure feedback transformations) acting on the class of elliptic/hyperbolic systems Ξ_{EH} is affine (actually it is of the Brockett type $w = \pm\tilde{w} + \alpha$) with respect to the control and thus depend on one function α only. By transforming into the the conformal form, we show that elliptic/hyperbolic systems Ξ_{EH} are given by three arbitrary smooth functions. Next, we show that the vanishing of the Gaussian curvature of a (pseudo-)Riemannian metric, associated with the EH-frame of Ξ_{EH} , characterises the flat elliptic/hyperbolic systems Ξ'_{EH} , which depend on two arbitrary smooth functions only. Finally, we gave conditions characterising the constant-form systems, i.e. elliptic/hyperbolic systems without functional parameters. In the elliptic case, equivalent constant-form systems correspond to the circles $\Gamma_E = (c_0)^2 + (c_1)^2 = \text{const.}$, and their canonical forms are parametrized by a closed half-line of real constants. On the other hand, in the hyperbolic case the structure is richer because equivalent systems correspond to connected branches of the hyperbolas $\Gamma_H = (c_0)^2 - (c_1)^2 = \text{const.}$; two connected components for $\Gamma_H > 0$, one for $\Gamma_H < 0$, and three for $\Gamma_H = 0$. Thus canonical forms of hyperbolic systems are parametrized by a real line of constants (the value of Γ_H) and by a discrete invariant $\varepsilon = \pm 1$ (if $\Gamma_H > 0$ or $\Gamma_H = 0$ and $c_0 \neq 0$). Our characterisation of elliptic/hyperbolic systems gives an equivalent classification for elliptic/hyperbolic submanifolds that is summarised in [Lemma 4.1](#) and explicitly given in [18].

4.2 Classification of parabolic systems

We now turn to the classification of parabolic systems $\Xi_P = (A, B, C)$, of the form

$$\Xi_P : \dot{x} = A(x)w^2 + B(x)w + C(x),$$

which is expected to be different from that of elliptic and hyperbolic systems because the allowed reparametrisations of the control w depend on 2 smooth functions (α, β) , see [Proposition 4.1](#). Our aim is to get a classification of parabolic submanifolds \mathcal{S}_P , as the one presented in [Table 1](#), via the properties of the triple (A, B, C) presented in [Table 2](#), see the beginning of [Section 4](#) for both tables. In particular, existence of a commutative P-frame (A, B) corresponds to weakly-flat parabolic submanifolds; existence of a commutative P-frame, which additionally satisfies $A \wedge C = 0$, corresponds to strongly-flat submanifolds; finally, existence of a commutative P-frame, which additionally satisfies C constant (in the coordinates where (A, B) is rectified), corresponds to constant-form parabolic submanifolds; in particular, the case of $C = 0$ is called null-form. As in the elliptic/hyperbolic cases, we introduce the structure functions $(\mu_0, \mu_1, \gamma_0, \gamma_1)$ uniquely defined for any triple (A, B, C) by

$$[A, B] = \mu_0 A + \mu_1 B \quad \text{and} \quad C = \gamma_0 A + \gamma_1 B.$$

By a direct computation, we obtain that under reparametrisations of the form $w = \beta\tilde{w} + \alpha$ the structure functions $(\mu_0, \mu_1, \gamma_0, \gamma_1)$ of (A, B, C) are transformed into the structure functions $(\tilde{\mu}_0, \tilde{\mu}_1, \tilde{\gamma}_0, \tilde{\gamma}_1)$ of $(\tilde{A}, \tilde{B}, \tilde{C})$ via the following relations:

$$\tilde{\gamma}_0 = \frac{1}{\beta^2} (\gamma_0 - 2\alpha\gamma_1 - \alpha^2), \quad \tilde{\gamma}_1 = \frac{1}{\beta} (\gamma_1 + \alpha), \quad (24)$$

$$\begin{aligned} \tilde{\mu}_0 &= \beta\mu_0 - 2\alpha L_A(\beta) + 2\beta L_A(\alpha) - 2L_B(\beta) - 2\alpha(L_A(\beta) + \beta\mu_1), \\ \tilde{\mu}_1 &= \beta^2\mu_1 + \beta L_A(\beta). \end{aligned} \quad (25)$$

There are two main questions that we will answer. First, when does a commutative P-frame (\tilde{A}, \tilde{B}) exist, i.e. $\tilde{\mu}_0 = \tilde{\mu}_1 = 0$? Second, provided that a commutative P-frame (A, B) has been normalised, how can we additionally simplify C ? Contrary to the elliptic and hyperbolic cases the answer to the first question is always positive without any additional assumption, as assured by the next result.

Proposition 4.5 (Existence of a commutative P-frame)

- (i) For any P-frame (A, B) there exists, locally around x_0 , a reparametrisation (α, β) such that (\tilde{A}, \tilde{B}) is a commutative P-frame.
- (ii) If (A, B) is a commutative P-frame, then (\tilde{A}, \tilde{B}) is also a commutative P-frame if and only if the reparametrisation (α, β) satisfies

$$L_A(\beta) = 0 \quad \text{and} \quad \frac{1}{\beta} L_B(\beta) = L_A(\alpha). \quad (26)$$

Proof

- (i) Consider a P-frame (A, B) whose structure functions are (μ_0, μ_1) . Apply a reparametrisation (α, β) , $\beta \neq 0$, given by a solution of the following system of equations (we solve the first equation for β , and plug in β into the second equation to solve it for α)

$$\begin{cases} L_A(\beta) = -\beta\mu_1 \\ L_A(\alpha) + \alpha\mu_1 = \frac{1}{2\beta} (2L_B(\beta) - \beta\mu_0) \end{cases}.$$

Then formula (25) implies $\tilde{\mu}_0 = \tilde{\mu}_1 = 0$, i.e. $[\tilde{A}, \tilde{B}] = 0$. Notice that to ensure $\beta \neq 0$, we may actually solve $L_A(\ln \beta) = -\mu_1$.

- (ii) Using relation (25) with $\mu_i = \tilde{\mu}_i = 0$, for $i = 0, 1$, we see that all reparametrisations (α, β) have to satisfy relation (26). Conversely, if (A, B) is a commutative P-frame ($\mu_0 = \mu_1 = 0$) and (α, β) is any solution of (26), with $\beta \neq 0$, then by (25) we obtain that $\tilde{\mu}_0 = \tilde{\mu}_1 = 0$, i.e. (\tilde{A}, \tilde{B}) is also a commutative P-frame. □

Immediately, item (i) of [Proposition 4.5](#) gives the following prenormal forms of parabolic systems Ξ_P . Recall that systems and equivalence are considered locally with respect to the state x and globally with respect to the control $w \in \mathbb{R}$.

Corollary 4.1 (Prenormal forms of Ξ_P) *The parabolic system Ξ_P is always feedback equivalent to the following prenormal forms, locally around x_0 :*

$$\Xi'_P : \begin{cases} \dot{z} = w^2 + c_0(x) \\ \dot{y} = w + c_1(x) \end{cases}, \quad \Xi''_P : \begin{cases} \dot{z} = w^2 + b(x)w + \Gamma_P(x) \\ \dot{y} = w \end{cases},$$

whose structure functions are $(\mu'_0, \mu'_1, \gamma'_0, \gamma'_1) = (0, 0, c_0, c_1)$, and $(\mu''_0, \mu''_1, \gamma''_0, \gamma''_1) = (\frac{\partial b}{\partial z}, 0, \Gamma_P, 0)$, respectively.

A parabolic system of the form Ξ'_P is called weakly-flat, this terminology is justified by generalisations of the previous result in higher dimensions that we will present in a future paper.

Remark 4.4 Since any parabolic system can be brought into Ξ'_P (and into Ξ''_P), it follows that all parabolic systems are locally parametrized by two functions of two variables (c_0 and c_1 , or, equivalently, b and Γ_P). This is in contrast with elliptic/hyperbolic systems Ξ_{EH} parametrized by three functions of two variables (compare [Proposition 4.2](#)).

Proof Apply to Ξ_P a reparametrisation (α, β) transforming its P-frame into a commutative P-frame (\tilde{A}, \tilde{B}) and let ϕ be a local diffeomorphism introducing coordinates $x = (z, y)$ such that $\phi_*\tilde{A} = \frac{\partial}{\partial z}$ and $\phi_*\tilde{B} = \frac{\partial}{\partial y}$. In this system of coordinates, Ξ_P takes the form Ξ'_P . Then apply to Ξ'_P the reparametrisation $\tilde{w} = w + c_1(x)$ to obtain the form Ξ''_P with $b = -2c_1$ and $\Gamma_P = c_0 + (c_1)^2$. The computation of the structure functions is straightforward. \square

Notice that the normal forms Ξ'_P and Ξ''_P are related by the reparametrisation $\tilde{w} = w + c_1(x)$. The function Γ_P (appearing in Ξ''_P) will be of special importance in the remaining part of this section, and in any P-frame (A, B) , commutative or not, we define it by setting

$$\Gamma_P = \gamma_0 + (\gamma_1)^2.$$

Recall that (γ_0, γ_1) are defined by $C = \gamma_0 A + \gamma_1 B$. Clearly, diffeomorphisms act on Γ_P by conjugation and reparametrisations (α, β) act by $\beta^2 \tilde{\Gamma}_P = \Gamma_P$ (as it can be computed from formula (24)).

The remaining part of this section shows how to additionally normalize Ξ'_P and Ξ''_P while preserving the commutativity of the P-frame (A, B) . Although for a parabolic system Ξ_P , there always exists a commutative P-frame (A, B) , its explicit construction can be difficult or even impossible, as it requires to solve a system of first order PDEs. For this reason, we will state our results for a general, not necessarily commutative, P-frame (A, B) .

Theorem 4.1 (Normalisation of parabolic systems) *Let $\Xi_P = (A, B, C)$ be a parabolic system with structure functions $(\mu_0, \mu_1, \gamma_0, \gamma_1)$. Then the following statements hold, locally around x_0 :*

(i) Ξ_P is strongly-flat, i.e. feedback equivalent to Ξ'_P with $c_1 \equiv 0$, we will denote that form Ξ'''_P , if and only if

$$L_A^2(\gamma_1) + \gamma_1 \left(L_A(\mu_1) - (\mu_1)^2 \right) = \frac{\mu_0 \mu_1}{2} + \frac{1}{2} L_A(\mu_0) + L_B(\mu_1). \quad (27)$$

(ii) Ξ_P is a constant-form, i.e. feedback equivalent to Ξ'_P with $(c_0, c_1) \in \mathbb{R}^2$, satisfying $c_0 + (c_1)^2 \in \mathbb{R}^*$, if and only if $\Gamma_P \neq 0$ and

$$\begin{aligned} L_A(\Gamma_P) + 2\mu_1 \Gamma_P &= 0, \\ L_B(\Gamma_P) + 2\Gamma_P L_A(\gamma_1) &= \Gamma_P \mu_0 - 2\Gamma_P \gamma_1 \mu_1. \end{aligned} \quad (28)$$

Moreover, in this case we can always normalize $c_0 = \pm 1$ and $c_1 = 0$ and we will denote that form Ξ_P^\pm .
(iii) Ξ_P is a null-form, i.e. feedback equivalent to Ξ'_P with $c_0 \equiv c_1 \equiv 0$, which we will denote Ξ_P^0 , if and only if (27) holds and, additionally, $\Gamma_P \equiv 0$.

Notice that items (i), (ii), and (iii) of the above theorem characterise, respectively, the following normal forms around x_0 (recall that we work globally with respect to w)

$$\Xi'''_P : \begin{cases} \dot{z} = w^2 + c_0(x) \\ \dot{y} = w \end{cases}, \quad \Xi_P^\pm : \begin{cases} \dot{z} = w^2 \pm 1 \\ \dot{y} = w \end{cases}, \quad \text{and} \quad \Xi_P^0 : \begin{cases} \dot{z} = w^2 \\ \dot{y} = w \end{cases}.$$

A parabolic system of the form Ξ'''_P is called strongly-flat (the terminology is justified by generalisations of the previous result to higher dimensions). Equivalent statements can be formulated to obtain special structures of the second prenormal form Ξ''_P , e.g. the normal form Ξ'''_P corresponds to Ξ''_P with $b \equiv 0$ and thus describes the intersection of the prenormal forms Ξ'_P and Ξ''_P . The constant-forms Ξ_P^\pm and Ξ_P^0

are clearly special cases of Ξ_P''' where c_0 is constant. Observe that contrary to the elliptic/hyperbolic case, where canonical forms of constant-form systems are parametrised by real-continuous parameters, in the parabolic case, there are only three distinct canonical forms of constant-form systems. The difference between the normal form Ξ_P^\pm and Ξ_P^0 lies in the existence or not of an equilibrium point: the control $w = 0$ defines an equilibrium of Ξ_P^0 while there are no equilibria for Ξ_P^\pm .

Proof

(i) **Sufficiency:** Consider a parabolic system $\Xi_P = (A, B, C)$, with structure functions $(\mu_0, \mu_1, \gamma_0, \gamma_1)$, and assume that relation (27) holds. Introduce the reparametrisation $\alpha = -\gamma_1$ and β given as a solution of the following system of equations

$$\begin{cases} \frac{1}{\beta} L_A(\beta) = -\mu_1 \\ \frac{1}{\beta} L_B(\beta) = \frac{1}{2}(\mu_0 - 2\gamma_1\mu_1 - 2L_A(\gamma_1)) \end{cases} \quad (29)$$

This system, rewritten for $\ln(\beta)$, admits solutions since the integrability condition

$$L_A(L_B(\ln \beta)) - L_B(L_A(\ln \beta)) = L_{[A, B]}(\ln \beta) = \mu_0 L_A(\ln \beta) + \mu_1 L_B(\ln \beta) \quad (29')$$

takes the form

$$\frac{1}{2} L_A(\mu_0) - \gamma_1 L_A(\mu_1) - L_A^2(\gamma_1) - \mu_1 L_A(\gamma_1) + L_B(\mu_1) = -\frac{1}{2} \mu_0 \mu_1 - \gamma_1 (\mu_1)^2 - \mu_1 L_A(\gamma_1) \quad (29'')$$

and is guaranteed by condition (27). Consider the system $\tilde{\Xi}_P = (\tilde{A}, \tilde{B}, \tilde{C})$ obtained by the above defined reparametrisation (α, β) and using relations (24) and (25) we deduce that the structure functions $(\tilde{\mu}_0, \tilde{\mu}_1, \tilde{\gamma}_0, \tilde{\gamma}_1)$ of $\tilde{\Xi}_P$ satisfy $\tilde{\mu}_0 = \tilde{\mu}_1 = 0$ and $\tilde{\gamma}_1 = 0$. By choosing local coordinates (z, y) such that $(\tilde{A}, \tilde{B}) = \left(\frac{\partial}{\partial z}, \frac{\partial}{\partial y}\right)$, we obtain the system Ξ_P' with $c_1 = \tilde{\gamma}_1 = 0$, i.e. a strongly-flat parabolic system Ξ_P''' .

(i) **Necessity:** Assume that $\Xi_P = (A, B, C)$ is feedback equivalent to $\tilde{\Xi}_P$ of the form Ξ_P' with $c_1 \equiv 0$, via ϕ and (α, β) . Then for $\tilde{\Xi}_P$ we have $\tilde{\mu}_0 = \tilde{\mu}_1 = \tilde{\gamma}_1 = 0$. First, by (24) we obtain $\alpha = -\gamma_1$ and by (25) we obtain that β satisfies the relations given by system (29) above. Therefore, by computing the integrability condition (29'), equivalently given by (29''), we conclude relation (27).

(ii) **Sufficiency:** Consider a parabolic system $\Xi_P = (A, B, C)$, with structure functions $(\mu_0, \mu_1, \gamma_0, \gamma_1)$, and assume that $\Gamma_P = \gamma_0 + (\gamma_1)^2 \neq 0$ and (28) holds. We follow the same reasoning as in the sufficiency of statement (i). Introduce a reparametrisation (α, β) , where $\alpha = -\gamma_1$ and β is a solution of the system of equations (29). To assure the existence of β , we have to fulfil the integrability condition of (29), which is (29''), equivalently (29'''). To this end, we differentiate the second condition of (28) along A and use $L_A(L_B(\cdot)) = L_B(L_A(\cdot)) + \mu_0 L_A(\cdot) + \mu_1 L_B(\cdot)$ to conclude (29'''). Consider the system $\tilde{\Xi}_P = (\tilde{A}, \tilde{B}, \tilde{C})$ obtained by the above defined reparametrisation (α, β) and using relations (24) and (25), we deduce that the structure functions $(\tilde{\mu}_0, \tilde{\mu}_1, \tilde{\gamma}_0, \tilde{\gamma}_1)$ of $\tilde{\Xi}_P = (\tilde{A}, \tilde{B}, \tilde{C})$ satisfy $\tilde{\mu}_0 = \tilde{\mu}_1 = \tilde{\gamma}_1 = 0$. Therefore $\tilde{\Gamma}_P = \tilde{\gamma}_0$, for which condition (28) yields $L_{\tilde{A}}(\tilde{\Gamma}_P) = L_{\tilde{B}}(\tilde{\Gamma}_P) = 0$, implying that $\tilde{\Gamma}_P$ is constant (we still have $\tilde{\Gamma}_P \neq 0$ since $\beta^2 \tilde{\Gamma}_P = \Gamma_P$). Introduce coordinates (\tilde{z}, \tilde{y}) such that $\tilde{A} = \frac{\partial}{\partial \tilde{z}}$ and $\tilde{B} = \frac{\partial}{\partial \tilde{y}}$, in which the system takes the form (recall that $\tilde{\gamma}_1 = 0$)

$$\begin{cases} \dot{\tilde{z}} = \tilde{w}^2 + c_0 \\ \dot{\tilde{y}} = \tilde{w} \end{cases},$$

with $c_0 \in \mathbb{R}^*$. Finally, defining new coordinates (z, y) by $z = \frac{\tilde{z}}{|c_0|}$ and $y = \frac{\tilde{y}}{\sqrt{|c_0|}}$, and reparametrising by $w = \frac{\tilde{w}}{\sqrt{|c_0|}}$, yields the normal form Ξ_P^\pm .

(ii) **Necessity:** Assume that Ξ_P , whose structure functions are $(\mu_0, \mu_1, \gamma_0, \gamma_1)$ and $\Gamma_P = \gamma_0 + \gamma_1^2$, is feedback equivalent, via ϕ and (α, β) , to $\tilde{\Xi}_P$ of the form Ξ_P' with $(c_0, c_1) \in \mathbb{R}^2$ satisfying $c_0 + c_1^2 \neq 0$. For $\tilde{\Xi}_P$ we have $\tilde{\mu}_0 = \tilde{\mu}_1 = 0$ and $\tilde{\gamma}_0 = c_0$, $\tilde{\gamma}_1 = c_1$, hence $\tilde{\Gamma}_P = c_0 + c_1^2 \neq 0$ implying $\Gamma_P \neq 0$ since $\beta^2 \tilde{\Gamma}_P = \Gamma_P$. By relation (25), we obtain that β satisfies the relations of system (29). Differentiating $\Gamma_P = \beta^2 \tilde{\Gamma}_P$ along A we deduce

$$L_A(\Gamma_P) = L_A(\beta^2 \tilde{\Gamma}_P) = 2\tilde{\Gamma}_P \beta L_A(\beta) = -2\tilde{\Gamma}_P \beta^2 \mu_1 = -2\mu_1 \Gamma_P,$$

giving the first relation of (28). A similar computation, by taking the derivative of $\Gamma_P = \beta^2 \tilde{\Gamma}_P$ along B , implies the second relation of (28).

- (iii) The proof of that statement is a special case of the proof of item (i) with the additional condition $\Gamma_P \equiv 0$.

Sufficiency: Use the proof of the sufficiency of item (i) to bring the system Ξ_P into Ξ_P''' . For the latter form we have $\Gamma_P = c_0(x)$, hence $c_0(x) \equiv 0$ (due to $\beta^2 \tilde{\Gamma}_P = \Gamma_P$ and assumption $\Gamma_P \equiv 0$) and we obtain the normal form Ξ_P^0 .

Necessity: Assume that Ξ_P , whose structure functions are $(\mu_0, \mu_1, \gamma_0, \gamma_1)$ and $\Gamma_P = \gamma_0 + \gamma_1^2$, is feedback equivalent, via ϕ and (α, β) , to $\tilde{\Xi}_P$ of the form Ξ_P^0 (which is, actually, Ξ_P' with $c_0 \equiv c_1 \equiv 0$). For that system we have $\tilde{\mu}_0 = \tilde{\mu}_1 = 0$ and $\tilde{\Gamma}_P \equiv 0$ and since Γ_P is transformed under (α, β) by $\beta^2 \tilde{\Gamma}_P = \Gamma_P$, we obtain the necessity of $\Gamma_P \equiv 0$. The necessity of (27) is deduced from the necessity part of statement (i). □

Observe that item (ii) of the above theorem does not explicitly require condition (27), while the normal form Ξ_P^\pm satisfies $c_1 \equiv 0$ and hence that condition has to be hidden in (28). Indeed, this can be observed by differentiating Γ_P along $[A, B]$ and using the constraint (28), which after a short computation gives condition (27).

Remark 4.5 (Interpretation of the conditions) We now give a tangible interpretation of our conditions. To this end, consider the system Ξ_P' for which we have $\mu_0 = \mu_1 = 0$, $\gamma_0 = c_0(x)$, $\gamma_1 = c_1(x)$ and thus $\Gamma_P(x) = c_0(x) + (c_1(x))^2$. First, condition (27) implies $\frac{\partial^2 c_1}{\partial z^2} = 0$, that is, c_1 is affine with respect to z , namely, $c_1(x) = c_1^0(y)z + c_1^1(y)$, and thus, $\Gamma_P(x)$ is given by $c_0(x) + (c_1^0(y)z + c_1^1(y))^2$. This means that if a weakly-flat system Ξ_P' is feedback equivalent to a strongly-flat system Ξ_P''' , then it is parametrized by 3 smooth functions, two of them being functions of y only, and it has the following form

$$\begin{cases} \dot{z} = w^2 + \Gamma_P(x) - (c_1^0(y)z + c_1^1(y))^2 \\ \dot{y} = w + c_1^0(y)z + c_1^1(y) \end{cases}.$$

By additionally applying the first equation of (28), we obtain $\frac{\partial c_0}{\partial z} + 2c_1^0(y)^2 z + 2c_1^0(y)c_1^1(y) = 0$ and thus $c_0(x)$ is a polynomial of degree 2 in z , related to $c_1(x)$ by $c_0(x) = -(c_1(x))^2 + c_2(y)$, for an arbitrary smooth function $c_2(y)$. We now have $\Gamma_P = \Gamma_P(y) = c_2(y)$ and we use the second equation of (28). Thus, we get $\Gamma_P(y) = G \exp(-2 \int c_1^1(y) dy)$, with $G \in \mathbb{R}$. To summarise, any system Ξ_P' satisfying (27) and (28), i.e. equivalent to a constant-form parabolic system, is parametrized by two arbitrary smooth functions of y and a constant $G \in \mathbb{R}$ and is expressed by the form

$$\begin{cases} \dot{z} = w^2 + \Gamma_P(y) - (c_1^0(y)z + c_1^1(y))^2 \\ \dot{y} = w + c_1^0(y)z + c_1^1(y) \end{cases},$$

where $\Gamma_P(y) = G \exp(-2 \int c_1^1(y) dy)$. Finally, Ξ_P' (satisfying (27) and (28)) is feedback equivalent to Ξ_P^0 if and only if $G = 0$, and is feedback equivalent to Ξ_P^+ , respectively to Ξ_P^- , if $G > 0$, resp. $G < 0$. The distinction between the three normal forms comes from the sign of Γ_P , which is thus a discrete invariant (and that sign is dictated by the value of the constant G).

5 Conclusions

In this paper, we studied the characterisation and the classification problem of three dimensional submanifolds of the tangent bundle of a smooth surface. We showed that the equivalence of submanifolds is reflected in the equivalence, under feedback transformations, of their first and second extensions treated as control-nonlinear and control-affine systems, respectively. We gave a complete characterisation of non-degenerate quadratic submanifolds and proposed a classification of the regular ones, namely the classes of elliptic, hyperbolic, and parabolic, submanifolds. To achieve our characterisation results, we introduced the novel class of quadratic control-affine systems and we gave a characterisation of that class in terms of relations between structure functions. Using our characterisation, we identified the subclasses of elliptic, hyperbolic, and parabolic control-affine systems (second extensions of elliptic, hyperbolic, and parabolic submanifolds). Moreover, we constructed a normal-form for all quadratisable systems and thus gave a normal form of quadratic submanifolds, that may smoothly pass from the elliptic to the hyperbolic classes. Finally, working within the class of control-nonlinear systems subject to a regular quadratic nonholonomic constraint, we exhibited several normal forms of elliptic, hyperbolic, and parabolic systems. In particular we highlighted a connection between the Gaussian curvature of a well-defined metric and the existence of a commutative frame for elliptic and hyperbolic systems. Normal forms include systems without functional parameters. As a consequence of our classification of quadratic systems, we obtained a classification of elliptic, hyperbolic, and parabolic submanifolds.

In future works we plan to extend our results to higher dimensional quadratic nonholonomic constraints, in particular, we will generalise our results for parabolic systems to the case of control-nonlinear systems (with the state-space being an $n \geq 3$ dimensional manifold) subject to paraboloid nonholonomic constraints. Another interesting problem is to characterise control-nonlinear systems subject to any algebraic nonholonomic constraint. For instance, in order to generalise our results for parabolic systems, one can study polynomial systems, that is, systems subject to the nonholonomic constraint $\dot{z} - \sum_{i=0}^d a_i(x)y^i = 0$.

Acknowledgements The authors would like to thank anonymous reviewers whose suggestions and remarks have substantially improved the presentation of the paper.

References

1. Bers, L.: Riemann Surfaces: Lectures given at the Institute of Mathematical Sciences, New York University 1957-1958. Stevens, Scientific books (1958)
2. Bogaevsky, I.A.: Implicit ordinary differential equations: Bifurcations and sharpening of equivalence. *Izvestiya: Mathematics* **78**(6), 1063–1078 (2014). DOI 10.1070/IM2014v078n06ABEH002720
3. Bonnard, B.: Quadratic control systems. *Mathematics of Control, Signals and Systems* **4**(2), 139–160 (1991). DOI 10.1007/BF02551263
4. Bryant, R.: Local diagonalisation of a degenerated 2d metric tensor. *MathOverflow*. URL <https://mathoverflow.net/q/427307>
5. Cartan, E.: Sur la théorie des systèmes en involution et ses applications à la relativité. *Bulletin de la Société mathématique de France* **2**, 88–118 (1931). DOI 10.24033/bsmf.1175
6. Darboux, G.: Sur le probleme de Pfaff. *Bulletin des sciences mathématiques et astronomiques* **6**(1), 14–36 (1882). URL <https://gallica.bnf.fr/ark:/12148/bpt6k68005v/f3.item>
7. Dubins, L.E.: On curves of minimal length with a constraint on average curvature, and with prescribed initial and terminal positions and tangents. *American Journal of Mathematics* **79**(3), 497 (1957). DOI 10.2307/2372560
8. Elkin, V.I.: *Reduction of Nonlinear Control Systems: A Differential Geometric Approach*, vol. 472. Springer Science & Business Media (2012)
9. Jakubczyk, B.: Equivalence and invariants of nonlinear control systems. In: *NonLinear Controllability and Optimal Control*, vol. 133, pp. 177–218. Routledge (1990)
10. Jakubczyk, B., Zhitomirskii, M.: Local reduction theorems and invariants for singular contact structures. *Annales de l'institut Fourier* **51**(1), 237–295 (2001). DOI 10.5802/aif.1823
11. Kang, W.: Extended controller form and invariants of nonlinear control systems with a single input. *Journal of Mathematical Systems Estimation and Control* **6**, 27–52 (1996)
12. Kang, W., Krener, A.J.: Extended quadratic controller normal form and dynamic state feedback linearization of nonlinear systems. *SIAM Journal on Control and Optimization* **30**(6), 1319–1337 (1992)
13. Martinet, J.: Sur les singularités des formes différentielles. *Annales de l'institut Fourier* **20**(1), 95–178 (1970). DOI 10.5802/aif.340
14. Monroy-Pérez, F.: Non-Euclidean Dubins' problem. *Journal of Dynamical and Control Systems* **4**(2), 249–272 (1998). DOI 10.1023/A:1022842019374
15. Pfaff, J.F.: *Methodus Generalis, Aequationes Differentiarum Partialium, Nec Non Aequationes Differentiales Vulgares, Utrasque Primi Ordinis Inter Quotcunque Variabiles, Complete Integrandi*. *Abhandlungen der Königlich Akademie der Wissenschaften in Berlin* (1815)
16. Respondek, W.: Feedback classification of nonlinear control systems in \mathbb{R}^2 and \mathbb{R}^3 . In: B. Jakubczyk, W. Respondek (eds.) *Geometry of Feedback and Optimal Control*, pp. 347–382. Marcel Dekker, New York (1998)
17. Respondek, W., Zhitomirskii, M.: Feedback classification of nonlinear control systems on 3-manifolds. *Mathematics of Control, Signals, and Systems* **8**(4), 299–333 (1995). DOI 10.1007/BF01209688
18. Schmoderer, T.: Study of control systems under quadratic nonholonomic constraints. Motion planning, introduction to the regularised continuation method. Ph.D. thesis, INSA Rouen Normandie (2022). URL <http://theses.fr/s221665>
19. Schottenloher, M.: *A Mathematical Introduction to Conformal Field Theory*, vol. 759. Springer (2008)
20. Spivak, M.: *A Comprehensive Introduction to Differential Geometry, Volume 4, 3rd Edition*, vol. 4, 3rd edn. Publish or Perish (1999)
21. Tall, I.A., Respondek, W.: Feedback classification of nonlinear single-input control systems with controllable linearization: normal forms, canonical forms, and invariants. *SIAM Journal on Control and Optimization* **41**(5), 1498–1531 (2002). DOI 10.1137/S0363012900381753
22. van der Schaft, A.: Observability and Controllability for Smooth Nonlinear Systems. *SIAM Journal on Control and Optimization* **20**(3), 338–354 (1982). DOI 10.1137/0320026
23. van der Schaft, A.: Linearization and input-output decoupling for general nonlinear systems. *Systems & Control Letters* **5**(1), 27–33 (1984). DOI 10.1016/0167-6911(84)90005-7
24. Zhitomirskii, M.: *Typical Singularities of Differential 1-Forms and Pfaffian Equations*, vol. 113. American Mathematical Soc. (1992)
25. Zhitomirskii, M.: Singularities and normal forms of smooth distributions. *Banach Center Publications* **32**(1), 395–409 (1995). DOI 10.4064/-32-1-395-409
26. Zhitomirskii, M., Respondek, W.: Simple germs of corank one affine distributions. *Banach Center Publications* **44**(1), 269–276 (1998). DOI 10.4064/-44-1-269-276
27. Zhu, J., Trélat, E., Cerf, M.: Planar tilting maneuver of a spacecraft: Singular arcs in the minimum time problem and chattering. arXiv:1504.06219 [math] (2015). URL <http://arxiv.org/abs/1504.06219>

A Simultaneous rectification of two rank 1 distributions in \mathbb{R}^2

We show that locally on \mathbb{R}^2 we can simultaneously rectify two involutive complementary distributions. To this end, consider $\mathcal{A} = \text{span}\{A\}$ and $\mathcal{B} = \text{span}\{B\}$, two distributions of rank 1 satisfying $\mathcal{A}(x_0) \oplus \mathcal{B}(x_0) = T_{x_0}\mathbb{R}^2$. Choose two functions ϕ and ψ satisfying $d\phi(x_0) \neq 0$, $d\psi(x_0) \neq 0$, $d\phi \perp \mathcal{B}$, and $d\psi \perp \mathcal{A}$. Obviously, $(d\phi \wedge d\psi)(x_0) \neq 0$ and set $(z, y) = (\phi, \psi)$. So in the (z, y) -coordinates, we have $\mathcal{A} = \text{span}\left\{\frac{\partial}{\partial z}\right\}$ and $\mathcal{B} = \text{span}\left\{\frac{\partial}{\partial y}\right\}$.

B Resolution of equation (6)

Consider f a smooth function of (z, y, w) , we will abbreviate (z, y) in x to shorten the notation. We show how to construct all smooth solutions around (x_0, w_0) of the equation

$$\frac{\partial^3 f}{\partial w^3} = \tau(x) \frac{\partial f}{\partial w}.$$

First, we will find an expression for $\frac{\partial f}{\partial w}$ that will be then integrated to obtain the desired form. Consider the following linear system of PDEs:

$$\begin{pmatrix} \frac{\partial f_1}{\partial w} \\ \frac{\partial f_2}{\partial w} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ \tau(x) & 0 \end{pmatrix} \begin{pmatrix} f_1 \\ f_2 \end{pmatrix}$$

given for the functions $f_1 = \frac{\partial f}{\partial w}$ and $f_2 = \frac{\partial^2 f}{\partial w^2}$. Solutions of this system (interpreted as a system of ODEs with x being a parameter) are expressed by the exponential of the matrix $\begin{pmatrix} 0 & w \\ \tau(x) & 0 \end{pmatrix}$ given by the formula

$$\exp \begin{pmatrix} 0 & w \\ \tau(x) & 0 \end{pmatrix} = \sum_{k=0}^{+\infty} \frac{w^{2k+1} \tau(x)^k}{(2k+1)!} \begin{pmatrix} 0 & 1 \\ \tau(x) & 0 \end{pmatrix} + \sum_{k=0}^{+\infty} \frac{w^{2k} \tau(x)^k}{(2k)!} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

obtained by expressing the power series of the exponential and by regrouping the terms of odd and even degree. Denote $b(x) = f_1(x, w_0)$ and $a(x) = f_2(x, w_0)$, thus we obtain

$$\frac{\partial f}{\partial w} = f_1(x, w) = a(x) \sum_{k=0}^{+\infty} \frac{(w - w_0)^{2k+1}}{(2k+1)!} \tau(x)^k + b(x) \sum_{k=0}^{+\infty} \frac{(w - w_0)^{2k}}{(2k)!} \tau(x)^k.$$

Integration of this expression yields

$$f(x, w) = a(x) \sum_{k=0}^{+\infty} \frac{(w - w_0)^{2k+2}}{(2k+2)!} \tau(x)^k + b(x) \sum_{k=0}^{+\infty} \frac{(w - w_0)^{2k+1}}{(2k+1)!} \tau(x)^k + c(x).$$

C Detailed computation of the proof of Theorem 3.2

We extend the computation made in the proof of Theorem 3.2.

C.1 Resolution of equation (10)

We show how to, locally around $0 \in \mathbb{R}^3$, solve the equation $\rho'' - 2\rho\rho' + \frac{4}{9}\rho^3 = 0$, where $\rho = \rho(x, w)$ and the derivatives are taken with respect to w . Take the new unknown function $R(x, w) = \exp\left(-\frac{2}{3} \int_0^w \rho(x, t) dt\right)$ which satisfies $R(x, 0) \neq 0$ and

$$R' = -\frac{2}{3}\rho R, \quad R'' = -\frac{2}{3}R\left(\rho' - \frac{2}{3}\rho^2\right), \quad R''' = -\frac{2}{3}R\left(\rho'' - 2\rho\rho' + \frac{4}{9}\rho^3\right) = 0.$$

Thus, $R(x, w) = a(x)w^2 + b(x)w + c(x)$ yielding $\rho = -\frac{3}{2} \frac{R'}{R} = -\frac{3}{2} \frac{2aw+b}{aw^2+bw+c}$. Since $R(x, 0) = c(x_0) \neq 0$, thus taking $d(x) = \frac{a}{c}$ and $e(x) = \frac{b}{c}$ we obtain

$$\rho(x, w) = -\frac{3}{2} \frac{2d(x)w + e(x)}{d(x)w^2 + e(x)w + 1}.$$

C.2 Smooth form of h'

We integrate $h''(x, w) = a(x)(d(x)w^2 + e(x)w + 1)^{-3/2}$ to obtain a smooth expression of $h'(x, w)$ around $0 \in \mathbb{R}^3$. Recall that we denote $p = p(x, w) = dw^2 + ew + 1$ and $\Delta = \Delta(x) = e^2 - 4d(x)$. We have

$$h'(x, w) = a(x) \int \frac{1}{p(x, w)^{3/2}} dw = \frac{-2a(2dw + e)}{\Delta\sqrt{p}} + \bar{b}(x).$$

Since $h'(x, 0) = b(x)$ is smooth, we have $\frac{-2ae}{\Delta} + \bar{b} = b(x)$, where $b(x)$ is a smooth function around 0. We can then derive a smooth closed form expression of $h'(x, w)$:

$$\begin{aligned} h'(x, w) &= \frac{-2a(2dw + e)}{\Delta\sqrt{p}} + \frac{2ae}{\Delta} + b = \frac{-2a}{\Delta\sqrt{p}} (2dw + e - e\sqrt{p}) + b, \\ &= \frac{-2a}{\Delta\sqrt{p}(ew + 2 + 2\sqrt{p})} (2dw + e - e\sqrt{p})(ew + 2 + 2\sqrt{p}) + b, \\ &= \frac{-2a}{\Delta\sqrt{p}(ew + 2 + 2\sqrt{p})} (2dew^2 + 4dw + 4dw\sqrt{p} + e^2w + 2e + 2e\sqrt{p} - e^2w\sqrt{p} - 2e\sqrt{p} - 2ep) + b, \\ &= \frac{-2a}{\Delta\sqrt{p}(ew + 2 + 2\sqrt{p})} (w\sqrt{p}(4d - e^2) + 4dw + 2dew^2 + 2e + e^2w - 2edw^2 - 2e^2w - 2e) + b, \\ &= \frac{-2a}{\Delta\sqrt{p}(ew + 2 + 2\sqrt{p})} (w\sqrt{p}(4d - e^2) + w(4d - e^2)) + b, \\ &= \frac{2aw}{\sqrt{p}(ew + 2 + 2\sqrt{p})} (\sqrt{p} + 1) + b. \end{aligned}$$

D Details of the computations of Lemma 4.2

Denoting $c_E(x) = \cos(x)$, $c_H(w) = \cosh(w)$, $s_E(w) = \sin(w)$, and $s_H(w) = \sinh(w)$ and starting from the system

$$\dot{x} = (A, B) \begin{pmatrix} c_{EH}(w) \\ s_{EH}(w) \end{pmatrix} + (A, B) \begin{pmatrix} \gamma_0 \\ \gamma_1 \end{pmatrix},$$

we apply a reparametrisation $w = \tilde{w} + \alpha(x)$:

$$\begin{aligned} \dot{x} &= (A, B) \bar{R}_{EH}(\pm\alpha) \begin{pmatrix} c_{EH}(\tilde{w}) \\ s_{EH}(\tilde{w}) \end{pmatrix} + (A, B) \begin{pmatrix} \gamma_0 \\ \gamma_1 \end{pmatrix}, \\ &= (\tilde{A}, \tilde{B}) \begin{pmatrix} c_{EH}(\tilde{w}) \\ s_{EH}(\tilde{w}) \end{pmatrix} + (\tilde{A}, \tilde{B}) \bar{R}_{EH}(\pm\alpha)^{-1} \begin{pmatrix} \gamma_0 \\ \gamma_1 \end{pmatrix}. \end{aligned}$$

This yields $(\tilde{\gamma}_0, \tilde{\gamma}_1) = (\gamma_0, \gamma_1) \bar{R}_{EH}(\pm\alpha)^{-T}$, and by the definition of $\bar{R}_{EH}(\alpha)$ we have,

$$\bar{R}_E(\alpha)^{-T} = \bar{R}_E(-\alpha)^T = \bar{R}_E(\alpha), \quad \text{and} \quad \bar{R}_H(-\alpha)^{-T} = \bar{R}_H(-\alpha)^{-1} = \bar{R}_H(\alpha).$$

Next, computing separately in the elliptic and hyperbolic cases, we have

$$\begin{aligned} [\tilde{A}, \tilde{B}] &= (\mu_0 \mp L_A(\alpha))A + (\mu_1 - L_B(\alpha))B, \\ (A, B) \bar{R}_E(\pm\alpha) \begin{pmatrix} \tilde{\mu}_0 \\ \tilde{\mu}_1 \end{pmatrix} &= (A, B) \begin{pmatrix} \mu_0 \mp L_A(\alpha) \\ \mu_1 - L_B(\alpha) \end{pmatrix}. \end{aligned}$$

Thus

$$(\tilde{\mu}_0, \tilde{\mu}_1) = (\mu_0 \mp L_A(\alpha), \mu_1 - L_B(\alpha)) \bar{R}_E(\pm\alpha)^{-T}$$

and relation (17) follows.

E Gaussian curvature for metric given in terms of vector fields

Consider a 2-dimensional manifold \mathcal{X} and two smooth vector fields A , and B satisfying $A \wedge B \neq 0$. Construct the (pseudo)-Riemannian metric \mathbf{g}_\pm defined by

$$\mathbf{g}_\pm(A, A) = 1, \quad \mathbf{g}_\pm(B, B) = \pm 1, \quad \text{and} \quad \mathbf{g}_\pm(A, B) = 0.$$

We will give a formula for the Gaussian curvature of \mathbf{g}_\pm in terms of the structure functions (μ_0, μ_1) uniquely defined by $[A, B] = \mu_0 A + \mu_1 B$. We will use the following formula for the covariant derivative

$$\nabla_{E_i} E_j = \frac{1}{2} \sum_k (\mathbf{g}_\pm([E_i, E_j], E_k) - \mathbf{g}_\pm([E_i, E_k], E_j) - \mathbf{g}_\pm([E_j, E_k], E_i)) E_k$$

for $E_i, E_j, E_k \in \{A, B\}$, and the following formula for the Gaussian curvature of a 2-dimensional manifold

$$\kappa_{\pm} = \frac{\mathfrak{g}_{\pm}((\nabla_B \nabla_A - \nabla_A \nabla_B + \nabla_{[A,B]})A, B)}{\det(\mathfrak{g}_{\pm})}.$$

Computing, we have

$$\nabla_A A = -\mu_0 B, \quad \nabla_A B = \mu_0 A, \quad \nabla_B A = \mp \mu_1 B, \quad \nabla_B B = \pm \mu_1 A.$$

Then we can deduce

$$\begin{aligned} \nabla_B \nabla_A A &= -L_B(\mu_0) B \mp \mu_0 \mu_1 A, & \nabla_A \nabla_B A &= \mp L_A(\mu_1) B \mp \mu_0 \mu_1 A, \\ \nabla_{[A,B]} A &= \mu_0 \nabla_A A + \mu_1 \nabla_B A = -(\mu_0)^2 B \mp (\mu_1)^2 B. \end{aligned}$$

Thus

$$\begin{aligned} \kappa_{\pm} &= \pm \mathfrak{g}_{\pm}((-L_B(\mu_0) \pm L_A(\mu_1) - (\mu_0)^2 \mp (\mu_1)^2) B, B), \\ &= -L_B(\mu_0) \pm L_A(\mu_1) - (\mu_0)^2 \mp (\mu_1)^2. \end{aligned}$$

Therefore, we have obtained the expression of relation (20).